



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 4, 2024 – 09:49 PM EST

PDB ID : 1XOM  
Title : Catalytic Domain Of Human Phosphodiesterase 4D In Complex With Cilomilast  
Authors : Card, G.L.; England, B.P.; Suzuki, Y.; Fong, D.; Powell, B.; Lee, B.; Luu, C.; Tabrizizad, M.; Gillette, S.; Ibrahim, P.N.; Artis, D.R.; Bollag, G.; Milburn, M.V.; Kim, S.-H.; Schlessinger, J.; Zhang, K.Y.J.  
Deposited on : 2004-10-06  
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1483 (1.56-1.56)
Clashscore	141614	1529 (1.56-1.56)
Ramachandran outliers	138981	1498 (1.56-1.56)
Sidechain outliers	138945	1495 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Mol	Chain	Length	Quality of chain
1	A	349	88% (green), 5% (yellow), 7% (grey)
1	B	349	88% (green), 8% (yellow), 4% (grey)

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	EDO	A	716	-	-	X	-
5	EDO	B	713	-	-	X	-

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6068 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cAMP-specific 3',5'-cyclic phosphodiesterase 4D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	326	2655	1678	450	511	16	4	6	0
1	B	321	2633	1668	446	503	16	0	9	0

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	65	MET	-	initiating methionine	UNP Q08499
A	66	GLY	-	cloning artifact	UNP Q08499
A	67	SER	-	cloning artifact	UNP Q08499
A	68	SER	-	cloning artifact	UNP Q08499
A	69	HIS	-	expression tag	UNP Q08499
A	70	HIS	-	expression tag	UNP Q08499
A	71	HIS	-	expression tag	UNP Q08499
A	72	HIS	-	expression tag	UNP Q08499
A	73	HIS	-	expression tag	UNP Q08499
A	74	HIS	-	expression tag	UNP Q08499
A	75	SER	-	cloning artifact	UNP Q08499
A	76	SER	-	cloning artifact	UNP Q08499
A	77	GLY	-	cloning artifact	UNP Q08499
A	78	LEU	-	cloning artifact	UNP Q08499
A	79	VAL	-	cloning artifact	UNP Q08499
A	80	PRO	-	cloning artifact	UNP Q08499
A	81	ARG	-	cloning artifact	UNP Q08499
A	82	GLY	-	cloning artifact	UNP Q08499
A	83	SER	-	cloning artifact	UNP Q08499
A	84	HIS	-	cloning artifact	UNP Q08499
A	85	MET	-	cloning artifact	UNP Q08499
B	65	MET	-	initiating methionine	UNP Q08499
B	66	GLY	-	cloning artifact	UNP Q08499
B	67	SER	-	cloning artifact	UNP Q08499
B	68	SER	-	cloning artifact	UNP Q08499

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	69	HIS	-	expression tag	UNP Q08499
B	70	HIS	-	expression tag	UNP Q08499
B	71	HIS	-	expression tag	UNP Q08499
B	72	HIS	-	expression tag	UNP Q08499
B	73	HIS	-	expression tag	UNP Q08499
B	74	HIS	-	expression tag	UNP Q08499
B	75	SER	-	cloning artifact	UNP Q08499
B	76	SER	-	cloning artifact	UNP Q08499
B	77	GLY	-	cloning artifact	UNP Q08499
B	78	LEU	-	cloning artifact	UNP Q08499
B	79	VAL	-	cloning artifact	UNP Q08499
B	80	PRO	-	cloning artifact	UNP Q08499
B	81	ARG	-	cloning artifact	UNP Q08499
B	82	GLY	-	cloning artifact	UNP Q08499
B	83	SER	-	cloning artifact	UNP Q08499
B	84	HIS	-	cloning artifact	UNP Q08499
B	85	MET	-	cloning artifact	UNP Q08499


- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0
2	B	1	Total Zn 1 1	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	B	1	Total Mg 1 1	0	0

- Molecule 4 is CILOMILAST (three-letter code: CIO) (formula: C<sub>20</sub>H<sub>25</sub>NO<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
4	A	1	Total	C	N	O	0	1
			50	40	2	8		
4	B	1	Total	C	N	O	0	1
			50	40	2	8		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	299	Total O 299 299	0	0

*Continued on next page...*

*Continued from previous page...*


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
6	B	297	Total 297	O 297	0	0



### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: cAMP-specific 3',5'-cyclic phosphodiesterase 4D

Chain A: 



- Molecule 1: cAMP-specific 3',5'-cyclic phosphodiesterase 4D

Chain B: 



SER

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	60.49Å 78.97Å 164.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	81.65 – 1.55 44.98 – 1.55	Depositor EDS
% Data completeness (in resolution range)	99.1 (81.65-1.55) 99.1 (44.98-1.55)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.35 (at 1.55Å)	Xtrriage
Refinement program	REFMAC 5.1.25	Depositor
R, $R_{free}$	0.183 , 0.205 0.238 , 0.258	Depositor DCC
$R_{free}$ test set	5683 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	15.2	Xtrriage
Anisotropy	0.310	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 56.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	6068	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.89% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, EDO, CIO, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.34	1/2735 (0.0%)	0.72	8/3716 (0.2%)
1	B	0.32	0/2728	0.71	6/3704 (0.2%)
All	All	0.33	1/5463 (0.0%)	0.71	14/7420 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	87	GLU	CB-CG	-9.32	1.34	1.52

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	87	GLU	CA-CB-CG	8.07	131.16	113.40
1	B	151	ASP	CB-CG-OD2	6.54	124.19	118.30
1	A	151	ASP	CB-CG-OD2	6.35	124.01	118.30
1	B	266	ASP	CB-CG-OD2	6.34	124.00	118.30
1	A	156	ASP	CB-CG-OD2	5.99	123.69	118.30
1	A	301	ASP	CB-CG-OD2	5.94	123.65	118.30
1	B	90	ASP	CB-CG-OD2	5.92	123.62	118.30
1	B	140	ASP	CB-CG-OD2	5.52	123.27	118.30
1	B	156	ASP	CB-CG-OD2	5.43	123.18	118.30
1	A	266	ASP	CB-CG-OD2	5.37	123.13	118.30
1	B	301	ASP	CB-CG-OD2	5.16	122.95	118.30
1	A	130	ASP	CB-CG-OD2	5.12	122.91	118.30
1	A	140	ASP	CB-CG-OD2	5.12	122.90	118.30
1	A	394	ASP	CB-CG-OD2	5.06	122.85	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2655	0	2606	9	2
1	B	2633	0	2589	8	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	50	0	48	0	0
4	B	50	0	48	0	0
5	A	36	0	54	6	0
5	B	44	0	66	7	0
6	A	299	0	0	0	2
6	B	297	0	0	2	0
All	All	6068	0	5411	19	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:406:TYR:OH	5:B:713:EDO:H22	1.77	0.84
1:A:395:ILE:HD12	5:A:716:EDO:H22	1.68	0.74
1:A:178[B]:THR:HG23	5:A:716:EDO:H11	1.78	0.64
1:B:88:GLN:N	6:B:2066:HOH:O	2.38	0.56
1:A:219:LEU:HD23	1:A:222[B]:MET:CE	2.39	0.53
1:A:392:ALA:CA	5:A:716:EDO:H12	2.43	0.49
1:A:175:LEU:O	1:A:178[A]:THR:HG23	2.13	0.47
1:B:330:ARG:HB3	5:B:713:EDO:H11	1.98	0.45
1:B:175:LEU:O	1:B:178[B]:THR:HG23	2.17	0.45
5:B:701:EDO:H22	5:B:713:EDO:H21	1.99	0.45
1:B:334:ASP:OD1	5:B:713:EDO:H12	2.17	0.45
1:A:392:ALA:HB2	5:A:716:EDO:H12	1.98	0.44
1:A:154:HIS:O	1:A:157:VAL:HG22	2.16	0.44
1:B:238:PHE:CD1	5:B:717:EDO:H22	2.53	0.44
1:A:178[B]:THR:CG2	5:A:716:EDO:H11	2.50	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:275[A]:LYS:NZ	6:B:2268:HOH:O	2.29	0.40
1:B:330:ARG:CB	5:B:713:EDO:H11	2.50	0.40
1:A:395:ILE:CD1	5:A:716:EDO:H22	2.46	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:87:GLU:OE1	6:A:1133:HOH:O[4_455]	1.24	0.96
1:A:87:GLU:CD	6:A:1133:HOH:O[4_455]	1.99	0.21

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	330/349 (95%)	326 (99%)	4 (1%)	0	100	100
1	B	326/349 (93%)	322 (99%)	4 (1%)	0	100	100
All	All	656/698 (94%)	648 (99%)	8 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	304/318 (96%)	301 (99%)	3 (1%)	76	57
1	B	303/318 (95%)	300 (99%)	3 (1%)	76	57
All	All	607/636 (95%)	601 (99%)	6 (1%)	81	57

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	87	GLU
1	A	127	GLN
1	A	182	GLU
1	B	178[A]	THR
1	B	178[B]	THR
1	B	302	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	88	GLN
1	A	245	ASN
1	A	308	GLN
1	B	88	GLN
1	B	123	HIS
1	B	242	GLN
1	B	245	ASN
1	B	308	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

Of 28 ligands modelled in this entry, 4 are monoatomic - leaving 24 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	EDO	A	702	-	3,3,3	0.25	0	2,2,2	0.41	0
4	CIO	B	601[A]	-	25,27,27	0.69	0	29,38,38	1.23	3 (10%)
5	EDO	B	721	-	3,3,3	0.29	0	2,2,2	0.32	0
5	EDO	A	716	-	3,3,3	0.28	0	2,2,2	0.15	0
5	EDO	B	711	-	3,3,3	0.28	0	2,2,2	0.34	0
5	EDO	B	718	-	3,3,3	0.28	0	2,2,2	0.35	0
4	CIO	A	603[A]	-	25,27,27	0.66	0	29,38,38	1.21	3 (10%)
5	EDO	A	715	-	3,3,3	0.25	0	2,2,2	0.39	0
5	EDO	B	713	-	3,3,3	0.30	0	2,2,2	0.49	0
5	EDO	B	714	-	3,3,3	0.29	0	2,2,2	0.29	0
5	EDO	A	712	-	3,3,3	0.27	0	2,2,2	0.31	0
4	CIO	B	601[B]	-	25,27,27	0.70	1 (4%)	29,38,38	1.08	4 (13%)
5	EDO	B	705	-	3,3,3	0.28	0	2,2,2	0.33	0
5	EDO	B	706	-	3,3,3	0.29	0	2,2,2	0.36	0
5	EDO	B	701	-	3,3,3	0.25	0	2,2,2	0.36	0
5	EDO	B	710	-	3,3,3	0.27	0	2,2,2	0.32	0
5	EDO	B	717	-	3,3,3	0.27	0	2,2,2	0.38	0
5	EDO	A	703	-	3,3,3	0.27	0	2,2,2	0.39	0
4	CIO	A	603[B]	-	25,27,27	0.75	1 (4%)	29,38,38	1.10	4 (13%)
5	EDO	A	704	-	3,3,3	0.28	0	2,2,2	0.28	0
5	EDO	A	719	-	3,3,3	0.26	0	2,2,2	0.44	0
5	EDO	B	709	-	3,3,3	0.27	0	2,2,2	0.31	0
5	EDO	A	707	-	3,3,3	0.29	0	2,2,2	0.54	0
5	EDO	A	720	-	3,3,3	0.26	0	2,2,2	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	A	702	-	-	0/1/1/1	-
4	CIO	B	601[A]	-	-	1/16/38/38	0/3/3/3
5	EDO	B	721	-	-	0/1/1/1	-
5	EDO	A	716	-	-	0/1/1/1	-
5	EDO	B	711	-	-	1/1/1/1	-
5	EDO	B	718	-	-	1/1/1/1	-
4	CIO	A	603[A]	-	-	1/16/38/38	0/3/3/3
5	EDO	A	715	-	-	0/1/1/1	-
5	EDO	B	713	-	-	1/1/1/1	-
5	EDO	B	714	-	-	0/1/1/1	-
5	EDO	A	712	-	-	0/1/1/1	-
4	CIO	B	601[B]	-	-	3/16/38/38	0/3/3/3
5	EDO	B	705	-	-	0/1/1/1	-
5	EDO	B	706	-	-	0/1/1/1	-
5	EDO	B	701	-	-	0/1/1/1	-
5	EDO	B	710	-	-	1/1/1/1	-
5	EDO	B	717	-	-	0/1/1/1	-
5	EDO	A	703	-	-	0/1/1/1	-
4	CIO	A	603[B]	-	-	0/16/38/38	0/3/3/3
5	EDO	A	704	-	-	1/1/1/1	-
5	EDO	A	719	-	-	0/1/1/1	-
5	EDO	B	709	-	-	1/1/1/1	-
5	EDO	A	707	-	-	1/1/1/1	-
5	EDO	A	720	-	-	0/1/1/1	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	603[B]	CIO	C1-C7	-2.40	1.51	1.53
4	B	601[B]	CIO	C1-C7	-2.14	1.51	1.53

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	601[A]	CIO	C8-C7-C1	3.76	124.78	121.07
4	A	603[A]	CIO	C8-C7-C1	3.56	124.58	121.07
4	A	603[A]	CIO	C12-O11-C10	3.17	122.32	117.53
4	A	603[B]	CIO	C12-O11-C10	3.05	122.13	117.53
4	B	601[A]	CIO	C12-O11-C10	2.96	122.00	117.53
4	B	601[B]	CIO	O11-C10-C13	2.92	119.48	115.41
4	A	603[B]	CIO	O11-C10-C13	2.37	118.72	115.41
4	A	603[A]	CIO	O25-C23-C4	2.31	120.24	114.21
4	B	601[B]	CIO	O25-C23-C4	2.21	120.00	114.21

Continued on next page...



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	601[B]	CIO	O24-C23-C4	-2.17	117.63	122.93
4	A	603[B]	CIO	O24-C23-C4	-2.16	117.64	122.93
4	B	601[A]	CIO	O25-C23-C4	2.04	119.56	114.21
4	B	601[B]	CIO	C12-O11-C10	2.03	120.60	117.53
4	A	603[B]	CIO	O25-C23-C4	2.01	119.46	114.21

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	601[B]	CIO	C13-C10-O11-C12
5	B	709	EDO	O1-C1-C2-O2
5	B	710	EDO	O1-C1-C2-O2
5	B	713	EDO	O1-C1-C2-O2
5	B	718	EDO	O1-C1-C2-O2
4	B	601[B]	CIO	C9-C10-O11-C12
5	B	711	EDO	O1-C1-C2-O2
5	A	707	EDO	O1-C1-C2-O2
4	B	601[A]	CIO	O25-C23-C4-C5
5	A	704	EDO	O1-C1-C2-O2
4	A	603[A]	CIO	O25-C23-C4-C5
4	B	601[B]	CIO	O25-C23-C4-C5



There are no ring outliers.


5 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	716	EDO	6	0
5	B	713	EDO	5	0
5	B	701	EDO	1	0
5	B	717	EDO	1	0
5	B	709	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.


### 6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands [i](#)




Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around CIO A 603 (B):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around CIO B 601 (A):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.