

# Full wwPDB X-ray Structure Validation Report (i)

#### Dec 23, 2024 – 04:25 PM JST

PDB ID : 8XO4	
Title : Crystal structure of measles virus fusion inhibitor M1EK comple	exed with F
protein HR1 (HR1-42) (P21 space group)	
Authors : Oishi, S.; Takahara, A.; Nakatsu, T.	
Deposited on : 2023-12-31	
Resolution : $2.36 \text{ Å}(\text{reported})$	

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.21
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.40

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 2.36 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
$R_{free}$	164625	1460 (2.36-2.36)
Clashscore	180529	$1571 \ (2.36-2.36)$
Ramachandran outliers	177936	1559 (2.36-2.36)
Sidechain outliers	177891	1559 (2.36-2.36)
RSRZ outliers	164620	1460 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	44	86%	11% •
1	С	44	77%	23%
1	Е	44	86%	14%
1	G	44	86%	14%
1	Ι	44	93%	7%
1	K	44	84%	16%



Mol	Chain	Length	Quality of chain	
1	М	44	95%	5%
1	Ο	44	91%	9%
1	Q	44	95%	5%
1	S	44	93%	7%
1	U	44	80%	20%
1	W	44	93%	7%
2	В	37	78%	22%
2	D	37	84%	16%
2	F	37	84%	16%
2	Н	37	89%	8% •
2	J	37	92%	8%
2	L	37	73%	27%
2	Ν	37	89%	11%
2	Р	37	81%	19%
2	R	37	86%	14%
2	Т	37	97%	•
2	V	37	86%	14%
2	Х	37	95%	5%



# 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 7732 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Δ	4.4	Total	С	Ν	0	S	0	1	1
1	A	44	330	198	61	70	1	0	L	1
1	C	4.4	Total	С	Ν	0	S	0	1	1
1	U	44	331	199	61	70	1	0	T	1
1	F	4.4	Total	С	Ν	Ο	S	0	0	1
1	Ľ	44	325	195	60	69	1	0	0	1
1	С	4.4	Total	С	Ν	Ο	S	0	0	1
1	G	44	325	195	60	69	1	0	0	1
1	Т	44	Total	С	Ν	Ο	$\mathbf{S}$	0	0	1
1	I	44	325	195	60	69	1	0	0	1
1	K	44	Total	С	Ν	Ο	$\mathbf{S}$	0	1	1
1	17	44	330	198	61	70	1	0	T	1
1	М	44	Total	С	Ν	Ο	$\mathbf{S}$	0	1	1
1	111	44	331	199	61	70	1	0	T	1
1	0	44	Total	С	Ν	Ο	$\mathbf{S}$	0	0	1
1	U	44	325	195	60	69	1	0	0	
1	0	44	Total	С	Ν	Ο	$\mathbf{S}$	0	0	1
1	Q	44	325	195	60	69	1	0	0	1
1	q	4.4	Total	С	Ν	Ο	$\mathbf{S}$	0	0	1
1	U U	44	325	195	60	69	1	0	0	1
1	II	44	Total	С	Ν	Ο	S	0	0	1
	0	<u>'1'1</u>	325	195	60	69	1		0	
1	W	44	Total	С	Ν	Ο	S	0	1	1
	v v	<u>'</u> ±'±	331	199	61	70	1		1	1 

• Molecule 1 is a protein called Fusion glycoprotein F1.

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	142	ACE	-	acetylation	UNP P69353
А	185	NH2	-	amidation	UNP P69353
С	142	ACE	-	acetylation	UNP P69353
С	185	NH2	-	amidation	UNP P69353
Е	142	ACE	-	acetylation	UNP P69353



Chain	Decidue	Medelled	Actual	Comment	Deference
Chain	Residue	Modelled	Actual	Comment	Reference
E	185	NH2	-	amidation	UNP P69353
G	142	ACE	-	acetylation	UNP P69353
G	185	NH2	-	amidation	UNP P69353
Ι	142	ACE	-	acetylation	UNP P69353
Ι	185	NH2	-	amidation	UNP P69353
K	142	ACE	-	acetylation	UNP P69353
K	185	NH2	-	amidation	UNP P69353
М	142	ACE	-	acetylation	UNP P69353
М	185	NH2	-	amidation	UNP P69353
0	142	ACE	-	acetylation	UNP P69353
0	185	NH2	-	amidation	UNP P69353
Q	142	ACE	-	acetylation	UNP P69353
Q	185	NH2	-	amidation	UNP P69353
S	142	ACE	-	acetylation	UNP P69353
S	185	NH2	-	amidation	UNP P69353
U	142	ACE	-	acetylation	UNP P69353
U	185	NH2	-	amidation	UNP P69353
W	142	ACE	-	acetylation	UNP P69353
W	185	NH2	-	amidation	UNP P69353

• Molecule 2 is a protein called Measles virus fusion inhibitor M1EK.

Mol	Chain	Residues		Aton	ns		ZeroOcc	AltConf	Trace
2	В	37	Total	С	Ν	0	0	1	1
	D	57	306	193	51	62	0	T	T
2	О	37	Total	$\mathbf{C}$	Ν	Ο	0	0	1
		01	300	189	51	60	0	0	1
2	F	37	Total	$\mathbf{C}$	Ν	Ο	0	2	1
	1	01	312	197	51	64	0		1
2	Н	37	Total	С	Ν	Ο	0	2	1
			314	198	54	62	Ŭ	-	-
2	J	37	Total	С	Ν	Ο	0	0	1
			300	189	51	60	0		-
2	L	37	Total	С	Ν	Ο	0	1	1
			306	194	52	60	Ŭ	-	-
2	N	37	Total	С	Ν	0	0	3	1
			318	201	51	66	Ŭ		-
2	Р	37	Total	С	Ν	Ο	0	0	1
	-		300	189	51	60	Ŭ		-
2	R	37	Total	С	Ν	Ο	0	1	1
	10		306	193	51	62	Ŭ	*	*
2	Т	37	Total	$\mathbf{C}$	Ν	Ο	0	1	1
	-		305	194	51	60		*	-



$\alpha$ $\cdot$ $\cdot$ $\cdot$	C	•	
Continued	trom	nremous	naae
Contentaca	JION	precedus	page

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace	
0	V	37	Total	С	Ν	0	0	2	1
	51	312	197	51	64	0		1	
0	V 97	Total	С	Ν	0	0	2	1	
	51	314	198	54	62	0			

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	10	Total         O           10         10	0	0
3	В	8	Total O 8 8	0	0
3	С	5	Total O 5 5	0	0
3	D	6	Total O 6 6	0	0
3	Е	5	Total O 5 5	0	0
3	F	7	Total O 7 7	0	1
3	G	1	Total O 1 1	0	0
3	Н	3	Total O 3 3	0	0
3	Ι	5	Total O 5 5	0	0
3	J	4	Total O 4 4	0	0
3	К	3	Total O 3 3	0	0
3	${ m L}$	3	Total O 3 3	0	0
3	М	4	$\begin{array}{cc} \text{Total} & \text{O} \\ 4 & 4 \end{array}$	0	0
3	Ν	8	Total O 8 8	0	0
3	Ο	2	Total O 2 2	0	0
3	Р	4	Total O 4 4	0	0
3	Q	2	Total O 2 2	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	R	4	Total O 4 4	0	0
3	S	6	Total O 6 6	0	0
3	Т	6	Total O 6 6	0	0
3	U	4	Total O 4 4	0	0
3	V	3	Total O 3 3	0	0
3	W	2	Total O 2 2	0	0
3	Х	6	Total O 6 6	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain A: 86% 11% • Molecule 1: Fusion glycoprotein F1 Chain C: 77% 23% • Molecule 1: Fusion glycoprotein F1 Chain E: 86% 14% • Molecule 1: Fusion glycoprotein F1 Chain G: 86% 14% • Molecule 1: Fusion glycoprotein F1 Chain I: 7% 93%
- Molecule 1: Fusion glycoprotein F1

• Molecule 1: Fusion glycoprotein F1



Chain K:	84%	16%
ACE142 N149 1161 1161 1161 1161 1161 1161 1161 1		
• Molecule 1: Fusion g	glycoprotein F1	
Chain M:	95%	5%
ACE142 1156 9176 NH2185		
• Molecule 1: Fusion g	glycoprotein F1	
Chain O:	91%	9%
ACE142 E162 A167 A167 A167 D180 V181 V181 V181		
• Molecule 1: Fusion	glycoprotein F1	
Chain Q:	95%	5%
ACE142 916 1170 112185		
• Molecule 1: Fusion	glycoprotein F1	
Chain S:	93%	7%
ACE142 N143 S1144 S114 A151 A179 N12185 N12185		
• Molecule 1: Fusion	glycoprotein F1	
Chain U:	80%	20%
ACE142 N143 S1144 Q145 Q145 Q159 Q159 R162 R165 R165 Y181	111-21-85 - 1-1-21-85 - 1-1-21-85 - 1-1-21-85 - 1-1-21-85 - 1-1-21-85 - 1-1-21-85 - 1-1-21-85 - 1-1-21-85 - 1-2 21-21-21-21-21-21-21-21-21-21-21-21-21-2	
• Molecule 1: Fusion	glycoprotein F1	
Chain W:	93%	7%
ACE142 T157 1161 N124 NH2185		
	R L D W I D E PROTEIN DATA BANK	

• Molecule 2: Measles virus fusion inhibitor M1EK

Chain B:	78%	22%
1452 1452 1455 1455 1454 1454 1454 1454	K486	
Molecule 2: Me	asles virus fusion inhibitor M1EK	
Chain D:	84%	16%
E455 E455 L463 E473 E473 E473 E473 E477	284CHN	
Molecule 2: Me	asles virus fusion inhibitor M1EK	
Chain F:	84%	16%
1452 1452 1455 1477 1477 1477 1477 1477 1477 1481 1481		
Molecule 2: Me	asles virus fusion inhibitor M1EK	
Chain H:	89%	8%
E455 E456 L477 K486 K486 NH2487		
Molecule 2: Me	asles virus fusion inhibitor M1EK	
Chain J:	92%	8%
E455 R456 E474 8174 NH2487		
Molecule 2: Me	asles virus fusion inhibitor M1EK	
Chain L:	73%	27%
E455 E455 E455 R457 K457 K456 K456 K456 V459 E450 E461 E461 L463 L463	L 476 L 477 K 496 NH2487	
Molecule 2: Me	asles virus fusion inhibitor M1EK	
Chain N:	89%	11%



#### ACE451 1452 E455 L476 L476 K479 K479 NH2487

• Molecule 2: Measles virus fusion inhibitor M1EK

Chain P:	81%	19%
ACE461 1452 1453 2454 2454 2455 2455 2455 2455 2455 2	K469 E474 M428 NH2487	
• Molecule 2:	Measles virus fusion inhibitor M1EK	
Chain R:	86%	14%
ACE451 E455 E455 E474 S480 K486 K486 NH2487		
• Molecule 2:	Measles virus fusion inhibitor M1EK	
Chain T:	97%	•
ACE461 1462 E453 MH2487		
• Molecule 2:	Measles virus fusion inhibitor M1EK	
Chain V:	86%	14%
ACE461 E455 R456 K458 K458 N462 N462	MH2487	
• Molecule 2:	Measles virus fusion inhibitor M1EK	
Chain X:	95%	5%



# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	55.84Å $35.58$ Å $237.65$ Å	Deperitor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $93.98^{\circ}$ $90.00^{\circ}$	Depositor
$\mathbf{P}_{\text{oscolution}}(\hat{\mathbf{A}})$	49.10 - 2.36	Depositor
Resolution (A)	49.10 - 2.36	EDS
% Data completeness	99.5 (49.10-2.36)	Depositor
(in resolution range)	99.5 (49.10-2.36)	EDS
R <sub>merge</sub>	(Not available)	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.42 (at 2.37 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
D D.	0.215 , $0.301$	Depositor
$\Pi, \Pi_{free}$	0.223 , $0.301$	DCC
$R_{free}$ test set	1965 reflections $(5.00%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	55.1	Xtriage
Anisotropy	0.694	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.31,44.0	EDS
L-test for twinning <sup>2</sup>	$< L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	0.026 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7732	wwPDB-VP
Average B, all atoms $(Å^2)$	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 8.29% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

# 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ACE, NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles	
	Unam	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.77	0/330	0.86	0/447
1	С	0.80	0/331	0.85	0/448
1	Е	0.79	0/322	0.87	0/436
1	G	0.73	0/322	0.82	0/436
1	Ι	0.71	0/322	0.82	0/436
1	Κ	0.70	0/330	0.81	0/447
1	М	0.71	0/331	0.83	0/448
1	0	0.72	0/322	0.81	0/436
1	Q	0.68	0/322	0.86	0/436
1	S	0.72	0/322	0.83	0/436
1	U	0.74	0/322	0.81	0/436
1	W	0.70	0/331	0.80	0/448
2	В	0.81	0/305	0.94	1/398~(0.3%)
2	D	0.75	0/296	0.84	0/386
2	F	0.77	0/314	0.91	0/410
2	Н	0.70	0/316	0.86	0/412
2	J	0.68	0/296	0.81	0/386
2	L	0.67	0/305	0.83	0/397
2	Ν	0.72	0/323	0.83	0/422
2	Р	0.73	0/296	0.83	0/386
2	R	0.74	0/305	0.86	0/398
2	Т	0.69	0/304	0.85	0/397
2	V	0.78	0/314	0.86	0/410
2	Х	0.74	0/316	0.91	0/412
All	All	0.73	0/7597	0.85	$1/1\overline{0104}\ (0.0\%)$

There are no bond length outliers.

All (1) bond angle outliers are listed below:



 $Observed(^{o}) | Ideal(^{o}) |$ 

Conti	nued from	n previe	ous page		
Mol	Chain	Res	Type	Atoms	

Mol	Chain	$\mathbf{Res}$	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	В	475	GLU	CB-CA-C	-5.31	99.78	110.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	330	0	323	5	0
1	С	331	0	325	8	0
1	Е	325	0	317	7	0
1	G	325	0	317	3	0
1	Ι	325	0	317	3	0
1	Κ	330	0	323	6	0
1	М	331	0	325	3	0
1	0	325	0	317	4	0
1	Q	325	0	317	1	0
1	S	325	0	317	2	0
1	U	325	0	317	7	0
1	W	331	0	325	2	0
2	В	306	0	338	6	0
2	D	300	0	332	6	0
2	F	312	0	344	7	0
2	Н	314	0	351	3	0
2	J	300	0	332	4	0
2	L	306	0	345	9	0
2	Ν	318	0	350	3	0
2	Р	300	0	332	6	0
2	R	306	0	338	3	0
2	Т	305	0	343	1	0
2	V	312	0	344	4	0
2	Х	314	0	351	3	0
3	A	10	0	0	1	0
3	В	8	0	0	0	0



8XO4
------

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	С	5	0	0	0	0
3	D	6	0	0	0	0
3	Ε	5	0	0	0	0
3	F	7	0	0	0	0
3	G	1	0	0	0	0
3	Н	3	0	0	0	0
3	Ι	5	0	0	0	0
3	J	4	0	0	1	0
3	Κ	3	0	0	0	0
3	L	3	0	0	0	0
3	М	4	0	0	0	0
3	Ν	8	0	0	0	0
3	Ο	2	0	0	0	0
3	Р	4	0	0	1	0
3	Q	2	0	0	0	0
3	R	4	0	0	0	0
3	S	6	0	0	0	0
3	Т	6	0	0	0	0
3	U	4	0	0	0	0
3	V	3	0	0	0	0
3	W	2	0	0	0	0
3	Х	6	0	0	0	0
All	All	7732	0	7940	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:181:TYR:HB2	2:D:455:GLU:HG3	1.73	0.69
1:O:162:GLU:OE2	2:R:474:GLU:OE1	2.11	0.69
2:B:474[A]:GLU:OE1	1:E:165:ARG:NE	2.31	0.63
1:W:157:THR:O	1:W:161:ILE:HG13	2.00	0.62
2:D:459:VAL:O	2:D:463:LEU:HG	2.00	0.61
1:G:165:ARG:NE	2:J:474:GLU:OE1	2.42	0.52
1:U:162:GLU:OE1	2:X:474:GLU:OE2	2.28	0.52
2:N:452:ILE:O	2:N:455[B]:GLU:HG3	2.10	0.52
1:O:179:GLN:OE1	2:R:455:GLU:OE2	2.27	0.52
1:C:165:ARG:NH1	1:C:169[B]:GLN:OE1	2.43	0.51
2:B:452:ILE:CG1	2:B:452:ILE:O	2.58	0.51



Atom 1 Atom 2		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:C:155:GLU:OE1	2:F:481[A]:GLU:OE1	2.29	0.51
2:P:453:GLU:OE2	2:X:456[B]:ARG:NH2	2.46	0.48
2:H:455[A]:GLU:OE2	1:K:179:GLN:OE1	2.30	0.48
2:H:477:LEU:HD11	1:K:161:ILE:HG22	1.95	0.48
2:L:454:GLU:O	2:L:458[B]:LYS:HG2	2.14	0.48
1:C:181:TYR:CB	2:D:455:GLU:HG3	2.42	0.48
1:E:181:TYR:CB	2:F:455:GLU:HG3	2.43	0.48
1:G:162:GLU:HG3	1:G:166:GLN:HE21	1.78	0.47
2:B:474[A]:GLU:OE1	1:E:162:GLU:OE1	2.33	0.47
1:U:181:TYR:HB2	2:V:455[B]:GLU:HG3	1.97	0.47
2:P:478:LYS:NZ	3:P:501:HOH:O	2.48	0.47
1:K:166:GLN:O	1:K:170:GLU:HG2	2.15	0.46
1:G:146:ALA:O	1:G:149:ASN:HB3	2.16	0.46
1:O:181:TYR:CB	2:P:455:GLU:HG3	2.45	0.46
1:C:142:ACE:O	1:C:145:GLN:HG2	2.16	0.46
1:C:166:GLN:O	1:C:170:GLU:HG2	2.15	0.46
2:T:453:GLU:HA	2:T:453:GLU:OE1	2.16	0.46
2:H:456[B]:ARG:HE	1:K:183:ASN:ND2	2.13	0.45
2:R:486:LYS:HG3	2:R:487:NH2:N	2.32	0.45
2:L:459:VAL:O	2:L:463:LEU:HG	2.17	0.45
1:S:179:GLN:NE2	2:V:456:ARG:HA	2.31	0.44
1:Q:166:GLN:O	1:Q:170:GLU:HG2	2.18	0.44
1:U:165:ARG:NE	2:X:474:GLU:OE1	2.49	0.44
2:B:455:GLU:OE2	2:B:459:VAL:HG21	2.17	0.44
1:C:172:ILE:HD11	2:F:466:ALA:HB1	1.99	0.44
1:S:151:ARG:HG2	1:U:150:LEU:HD21	2.00	0.44
1:C:163:ALA:HB3	2:D:473:ALA:HB2	1.98	0.44
2:J:456:ARG:HD3	3:J:501:HOH:O	2.18	0.44
2:L:457:LYS:O	2:L:461:GLU:HB3	2.18	0.44
1:I:179:GLN:OE1	2:L:455:GLU:OE2	2.36	0.43
1:A:161:ILE:HG22	2:D:477:LEU:HD22	2.00	0.43
1:U:165:ARG:NH1	1:U:169:GLN:OE1	2.49	0.43
1:A:153:SER:OG	1:E:154:LEU:HD21	2.17	0.43
2:B:452:ILE:O	2:B:452:ILE:HG12	2.19	0.43
2:V:458:LYS:O	2:V:462:ASN:ND2	2.47	0.43
1:M:176[B]:GLN:HG3	2:P:463:LEU:HD13	1.99	0.43
1:A:166:GLN:O	1:A:170:GLU:HG2	2.18	0.43
1:I:181:TYR:HB2	2:J:455:GLU:HG3	2.01	0.42
2:B:484:LEU:HD13	1:E:155:GLU:HG3	2.01	0.42
1:O:167:ALA:HB2	2:P:469:LYS:HB2	2.01	0.42
2:L:457:LYS:O	2:L:461:GLU:CB	2.68	0.42



Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
2:F:452:ILE:HG21	2:J:456:ARG:HH21	1.84	0.42
1:U:143:ASN:N	1:U:143:ASN:HD22	2.16	0.42
1:U:159:GLN:HB3	2:V:476:LEU:HD13	2.02	0.42
1:E:181:TYR:HB2	2:F:455:GLU:HG3	2.02	0.41
1:K:149[A]:ASN:ND2	2:L:486:LYS:O	2.54	0.41
2:P:474:GLU:OE2	2:P:474:GLU:HA	2.19	0.41
1:A:148:ASP:HB2	3:A:206:HOH:O	2.20	0.41
1:I:161:ILE:HG22	2:L:477:LEU:HD11	2.03	0.41
1:K:160:ALA:HB2	2:L:476:LEU:HB3	2.03	0.41
2:F:477:LEU:HD23	2:F:477:LEU:HA	1.89	0.40
1:E:181:TYR:HB2	2:F:455:GLU:CG	2.51	0.40
1:M:156:THR:HG21	2:N:479:LYS:HB3	2.03	0.40
1:M:156:THR:HG22	2:N:476:LEU:O	2.22	0.40
1:A:162:GLU:OE1	2:D:474:GLU:OE1	2.40	0.40
2:L:477:LEU:HD23	2:L:477:LEU:HA	1.84	0.40
1:W:184:ASN:HD22	1:W:184:ASN:HA	1.75	0.40

There are no symmetry-related clashes.

# 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	43/44~(98%)	43 (100%)	0	0	100 100
1	С	43/44~(98%)	42 (98%)	1 (2%)	0	100 100
1	Ε	42/44~(96%)	39 (93%)	2(5%)	1 (2%)	5 3
1	G	42/44~(96%)	42 (100%)	0	0	100 100
1	Ι	42/44~(96%)	41 (98%)	1 (2%)	0	100 100
1	К	43/44~(98%)	41 (95%)	2(5%)	0	100 100
1	М	43/44~(98%)	42 (98%)	1 (2%)	0	100 100



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	Ο	42/44~(96%)	41 (98%)	1 (2%)	0	100	100
1	Q	42/44~(96%)	40 (95%)	2 (5%)	0	100	100
1	S	42/44~(96%)	41 (98%)	1 (2%)	0	100	100
1	U	42/44~(96%)	41 (98%)	0	1 (2%)	5	3
1	W	43/44~(98%)	42 (98%)	1 (2%)	0	100	100
2	В	36/37~(97%)	36 (100%)	0	0	100	100
2	D	35/37~(95%)	35 (100%)	0	0	100	100
2	F	37/37~(100%)	35~(95%)	1 (3%)	1 (3%)	4	2
2	Н	37/37~(100%)	33 (89%)	3~(8%)	1 (3%)	4	2
2	J	35/37~(95%)	34 (97%)	1 (3%)	0	100	100
2	L	36/37~(97%)	35~(97%)	1 (3%)	0	100	100
2	Ν	38/37~(103%)	37~(97%)	1 (3%)	0	100	100
2	Р	35/37~(95%)	35 (100%)	0	0	100	100
2	R	36/37~(97%)	31 (86%)	5(14%)	0	100	100
2	Т	36/37~(97%)	35~(97%)	1 (3%)	0	100	100
2	V	37/37~(100%)	32 (86%)	5 (14%)	0	100	100
2	Х	37/37~(100%)	34 (92%)	3~(8%)	0	100	100
All	All	944/972~(97%)	907 (96%)	33 (4%)	4 (0%)	30	34

All (4) Ramachandran outliers are listed below:

Mol	Chain	$\operatorname{Res}$	Type
2	Н	486	LYS
2	F	486	LYS
1	U	144	SER
1	Е	144	SER

### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	А	35/34~(103%)	34~(97%)	1 (3%)	37	48
1	С	35/34~(103%)	35~(100%)	0	100	100
1	Ε	34/34~(100%)	34 (100%)	0	100	100
1	G	34/34~(100%)	33~(97%)	1 (3%)	37	48
1	Ι	34/34~(100%)	34 (100%)	0	100	100
1	Κ	35/34~(103%)	35~(100%)	0	100	100
1	М	35/34~(103%)	35~(100%)	0	100	100
1	Ο	34/34~(100%)	34 (100%)	0	100	100
1	Q	34/34~(100%)	34 (100%)	0	100	100
1	S	34/34~(100%)	33~(97%)	1 (3%)	37	48
1	U	34/34~(100%)	33~(97%)	1 (3%)	37	48
1	W	35/34~(103%)	35~(100%)	0	100	100
2	В	34/33~(103%)	32~(94%)	2~(6%)	16	18
2	D	33/33~(100%)	33~(100%)	0	100	100
2	F	35/33~(106%)	35~(100%)	0	100	100
2	Н	35/33~(106%)	33~(94%)	2~(6%)	17	19
2	J	33/33~(100%)	33 (100%)	0	100	100
2	L	34/33~(103%)	34 (100%)	0	100	100
2	Ν	36/33~(109%)	36 (100%)	0	100	100
2	Р	33/33~(100%)	32~(97%)	1 (3%)	36	46
2	R	34/33~(103%)	33 (97%)	1 (3%)	37	48
2	Т	34/33~(103%)	34 (100%)	0	100	100
2	V	35/33~(106%)	35 (100%)	0	100	100
2	Х	35/33~(106%)	35 (100%)	0	100	100
All	All	824/804~(102%)	814 (99%)	10 (1%)	70	80

All (10) residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	А	153	SER
2	В	453	GLU
2	В	486	LYS
1	G	154	LEU
2	Н	456[A]	ARG
2	Н	456[B]	ARG



Continued from previous page...

Mol	Chain	Res	Type
2	Р	458	LYS
2	R	480	SER
1	S	144	SER
1	U	145	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	145	GLN
1	С	145	GLN
1	Е	145	GLN
1	Е	169	GLN
1	Е	184	ASN
1	Ι	179	GLN
1	Κ	179	GLN
1	Κ	183	ASN
1	М	179	GLN
1	0	184	ASN
1	S	143	ASN
1	S	179	GLN
1	U	145	GLN
1	U	179	GLN
1	W	166	GLN
1	W	184	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.



## 5.6 Ligand geometry (i)

There are no ligands in this entry.

## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

# 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	7	#RSF	RZ>2	$OWAB(Å^2)$	Q < 0.9
1	А	42/44~(95%)	-0.43	0	100	100	37, 54, 82, 96	1 (2%)
1	С	42/44~(95%)	-0.37	0	100	100	31,60,74,92	1 (2%)
1	Е	42/44~(95%)	-0.48	0	100	100	38,51,75,92	0
1	G	42/44~(95%)	-0.22	0	100	100	52, 70, 93, 114	0
1	Ι	42/44~(95%)	-0.22	0	100	100	56, 74, 115, 125	0
1	K	42/44~(95%)	-0.06	0	100	100	59, 77, 119, 125	1 (2%)
1	М	42/44~(95%)	-0.28	0	100	100	28, 57, 83, 104	1 (2%)
1	Ο	42/44~(95%)	-0.34	0	100	100	52, 63, 97, 103	0
1	Q	42/44~(95%)	-0.35	0	100	100	43, 65, 85, 91	0
1	S	42/44~(95%)	-0.46	0	100	100	48, 64, 99, 116	0
1	U	42/44~(95%)	-0.37	0	100	100	44, 60, 88, 121	0
1	W	42/44~(95%)	-0.59	0	100	100	27, 56, 79, 105	1 (2%)
2	В	35/37~(94%)	-0.38	0	100	100	29, 59, 96, 99	1 (2%)
2	D	35/37~(94%)	-0.41	0	100	100	48, 66, 87, 101	0
2	F	35/37~(94%)	-0.42	0	100	100	33, 62, 89, 99	2 (5%)
2	Н	35/37~(94%)	-0.17	0	100	100	34, 78, 105, 113	2(5%)
2	J	35/37~(94%)	-0.29	0	100	100	64, 78, 97, 110	0
2	L	35/37~(94%)	-0.01	0	100	100	50, 91, 114, 120	1 (2%)
2	N	35/37~(94%)	-0.35	0	100	100	28, 63, 89, 91	3 (8%)
2	Р	35/37~(94%)	-0.32	0	100	100	55, 73, 92, 103	0
2	R	35/37~(94%)	-0.16	0	100	100	38, 74, 98, 103	1 (2%)
2	Т	35/37~(94%)	-0.21	0	100	100	43, 70, 94, 106	1 (2%)
2	V	35/37~(94%)	-0.21	0	100	100	34, 72, 98, 100	2(5%)
2	X	$\overline{35/37}\ (94\%)$	-0.40	0	100	100	31, 66, 89, 94	2(5%)
							Continued on ne	ext page



Continued from previous page...

Mol	Chain	Analysed	< <b>RSRZ</b> >	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
All	All	924/972~(95%)	-0.32	0 100 100	27, 67, 101, 125	20 (2%)

There are no RSRZ outliers to report.

# 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

## 6.4 Ligands (i)

There are no ligands in this entry.

# 6.5 Other polymers (i)

There are no such residues in this entry.

