



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2024 – 12:00 AM EST

PDB ID : 1XSE
Title : Crystal Structure of Guinea Pig 11beta-Hydroxysteroid Dehydrogenase Type 1
Authors : Ogg, D.; Elleby, B.; Norstrom, C.; Stefansson, K.; Abrahmsen, L.; Oppermann, U.; Svensson, S.
Deposited on : 2004-10-19
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	295	 69% 19% • 7%
1	B	295	 68% 20% • 7%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4655 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 11beta-hydroxysteroid dehydrogenase type 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	274	2105	1351	351	388	15	0	0	0
1	B	274	2105	1351	351	388	15	0	0	0

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	3	MET	-	cloning artifact	UNP Q6QLL4
A	4	GLY	-	cloning artifact	UNP Q6QLL4
A	5	SER	-	cloning artifact	UNP Q6QLL4
A	6	SER	-	cloning artifact	UNP Q6QLL4
A	7	HIS	-	cloning artifact	UNP Q6QLL4
A	8	HIS	-	cloning artifact	UNP Q6QLL4
A	9	HIS	-	cloning artifact	UNP Q6QLL4
A	10	HIS	-	cloning artifact	UNP Q6QLL4
A	11	HIS	-	cloning artifact	UNP Q6QLL4
A	12	HIS	-	cloning artifact	UNP Q6QLL4
A	13	SER	-	cloning artifact	UNP Q6QLL4
A	14	SER	-	cloning artifact	UNP Q6QLL4
A	15	GLY	-	cloning artifact	UNP Q6QLL4
A	16	LEU	-	cloning artifact	UNP Q6QLL4
A	17	VAL	-	cloning artifact	UNP Q6QLL4
A	18	PRO	-	cloning artifact	UNP Q6QLL4
A	19	ARG	-	cloning artifact	UNP Q6QLL4
A	20	GLY	-	cloning artifact	UNP Q6QLL4
A	21	SER	-	cloning artifact	UNP Q6QLL4
A	22	HIS	-	cloning artifact	UNP Q6QLL4
A	23	MET	-	cloning artifact	UNP Q6QLL4
B	3	MET	-	cloning artifact	UNP Q6QLL4
B	4	GLY	-	cloning artifact	UNP Q6QLL4
B	5	SER	-	cloning artifact	UNP Q6QLL4
B	6	SER	-	cloning artifact	UNP Q6QLL4

Continued on next page...

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	235	Total 235	O 235	0	0
3	B	114	Total 114	O 114	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 11beta-hydroxysteroid dehydrogenase type 1

Chain A: 



- Molecule 1: 11beta-hydroxysteroid dehydrogenase type 1

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, α , β , γ	118.39Å 118.39Å 184.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.50 9.98 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.2 (10.00-2.50) 99.7 (9.98-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.26 (at 2.50Å)	Xtrriage
Refinement program	REFMAC 5.2.0001	Depositor
R, R_{free}	0.192 , 0.267 0.255 , 0.330	Depositor DCC
R_{free} test set	1161 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å ²)	56.4	Xtrriage
Anisotropy	0.032	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.44 , 78.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4655	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.75	0/2145	0.84	5/2897 (0.2%)
1	B	0.64	1/2145 (0.0%)	0.77	3/2897 (0.1%)
All	All	0.70	1/4290 (0.0%)	0.81	8/5794 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	80	GLU	CG-CD	6.45	1.61	1.51

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	295	LEU	CA-CB-CG	7.42	132.37	115.30
1	A	252	ARG	NE-CZ-NH2	-7.11	116.75	120.30
1	A	204	ASN	N-CA-C	7.05	130.04	111.00
1	B	204	ASN	N-CA-C	6.35	128.14	111.00
1	A	198	ARG	NE-CZ-NH2	-6.17	117.22	120.30
1	A	252	ARG	NE-CZ-NH1	5.76	123.18	120.30
1	B	180	ILE	N-CA-C	5.67	126.30	111.00
1	A	198	ARG	NE-CZ-NH1	5.37	122.98	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2105	0	2135	63	0
1	B	2105	0	2135	56	0
2	A	48	0	26	4	0
2	B	48	0	26	3	0
3	A	235	0	0	5	0
3	B	114	0	0	6	0
All	All	4655	0	4322	106	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (106) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:171:VAL:HB	3:A:1518:HOH:O	1.25	1.30
1:B:122:LEU:H	1:B:143:ASN:HD21	1.06	1.00
1:A:140:MET:HE1	1:A:144:PHE:CD2	2.12	0.84
1:A:122:LEU:H	1:A:143:ASN:HD21	1.28	0.80
1:B:220:THR:HG22	1:B:223:ALA:H	1.49	0.77
1:A:276:MET:CE	1:B:267:LEU:HD12	2.15	0.76
1:B:122:LEU:H	1:B:143:ASN:ND2	1.84	0.75
1:A:125:ARG:CG	1:A:125:ARG:HH11	2.01	0.74
1:A:220:THR:HG22	1:A:223:ALA:H	1.53	0.72
1:A:140:MET:HE3	1:A:144:PHE:HB3	1.74	0.70
1:A:162:GLN:O	1:A:252:ARG:NH2	2.25	0.69
1:A:162:GLN:HG2	1:A:207:ASN:HB3	1.74	0.69
1:A:125:ARG:HH11	1:A:125:ARG:HG3	1.56	0.69
1:A:140:MET:CE	1:A:144:PHE:CD2	2.75	0.69
1:B:220:THR:HG21	2:B:2300:NDP:O2N	1.93	0.68
1:A:207:ASN:HA	3:A:1342:HOH:O	1.94	0.68
1:A:63:VAL:HG13	1:A:71:LEU:HD22	1.75	0.68
1:A:87:HIS:HD2	3:A:1306:HOH:O	1.76	0.68
1:A:206:VAL:O	1:A:207:ASN:HB2	1.94	0.67
1:A:40:THR:OG1	1:A:120:HIS:HD2	1.79	0.64
1:A:273:ARG:HD3	1:B:175:ILE:HG13	1.80	0.63
1:A:125:ARG:HH11	1:A:125:ARG:CB	2.11	0.62
1:A:120:HIS:HE1	1:A:146:SER:OG	1.83	0.62
1:A:175:ILE:HD13	1:A:268:LEU:CD2	2.30	0.61
1:A:276:MET:HE1	1:B:267:LEU:HD12	1.83	0.60
1:B:119:ASN:HD22	1:B:168:VAL:HG21	1.67	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:40:THR:O	1:B:119:ASN:HB3	2.01	0.59
1:A:270:ASN:HB3	1:B:270:ASN:CB	2.33	0.58
1:A:267:LEU:HD12	1:B:276:MET:CE	2.34	0.57
1:B:292:ASN:O	1:B:295:LEU:HD12	2.04	0.57
1:A:137:ARG:HH22	1:B:141:GLU:CD	2.08	0.57
1:A:239:GLU:HB3	3:A:1313:HOH:O	2.03	0.57
1:A:232:LEU:HD11	1:B:285:ASN:HB2	1.86	0.57
1:B:40:THR:HB	1:B:120:HIS:HD2	1.70	0.57
1:A:169:SER:O	2:A:1300:NDP:H6N	2.05	0.56
1:B:295:LEU:HD13	1:B:296:TYR:N	2.21	0.56
1:A:200:GLU:O	1:A:204:ASN:HB2	2.07	0.55
1:B:174:LYS:O	1:B:175:ILE:HD12	2.07	0.54
1:A:175:ILE:CD1	1:A:268:LEU:HD23	2.39	0.53
1:A:125:ARG:HG3	1:A:125:ARG:NH1	2.24	0.52
1:A:122:LEU:H	1:A:143:ASN:ND2	2.04	0.52
1:A:216:GLY:O	2:A:1300:NDP:H42N	2.10	0.52
1:B:40:THR:CB	1:B:120:HIS:HD2	2.23	0.52
1:A:178:PRO:O	1:A:179:LEU:HB2	2.10	0.52
1:B:296:TYR:HD2	1:B:297:GLY:N	2.08	0.52
1:A:175:ILE:CD1	1:A:268:LEU:CD2	2.88	0.51
1:A:220:THR:HG21	2:A:1300:NDP:O2N	2.11	0.51
1:B:63:VAL:HG12	1:B:71:LEU:HD22	1.93	0.51
1:A:206:VAL:O	1:A:207:ASN:CB	2.55	0.50
1:A:231:TYR:CE1	1:B:283:GLU:HB3	2.46	0.50
1:B:295:LEU:HD13	1:B:296:TYR:H	1.75	0.50
1:A:126:LEU:CD2	1:A:180:ILE:CG2	2.90	0.50
1:A:53:HIS:O	1:A:57:MET:HE3	2.12	0.49
1:B:85:SER:HB3	3:B:2414:HOH:O	2.12	0.49
1:A:162:GLN:HG2	1:A:207:ASN:CB	2.41	0.49
1:B:273:ARG:HG2	3:B:2334:HOH:O	2.12	0.49
1:B:40:THR:HG1	1:B:119:ASN:H	1.60	0.49
1:B:63:VAL:O	1:B:88:TYR:HA	2.13	0.49
1:B:64:THR:HB	1:B:102:PHE:CE1	2.48	0.48
1:B:36:LYS:HG2	1:B:114:ASP:OD2	2.13	0.48
1:B:37:VAL:HG22	1:B:115:MET:HB3	1.95	0.48
1:A:175:ILE:HD13	1:A:268:LEU:HD23	1.94	0.48
1:B:273:ARG:HD2	1:B:277:GLU:OE2	2.13	0.48
1:B:40:THR:HA	1:B:64:THR:HG22	1.96	0.47
1:A:54:LEU:HA	1:A:57:MET:HE3	1.95	0.47
1:B:169:SER:O	2:B:2300:NDP:H6N	2.14	0.47
1:B:290:LEU:HA	3:B:2340:HOH:O	2.14	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:107:GLY:O	1:B:111:GLY:N	2.48	0.47
1:A:63:VAL:O	1:A:88:TYR:HA	2.14	0.47
1:B:296:TYR:CD2	1:B:297:GLY:N	2.83	0.47
1:A:195:SER:HB2	1:B:176:THR:HG21	1.96	0.46
1:A:141:GLU:OE2	1:B:137:ARG:NH2	2.49	0.46
1:A:246:ILE:HD12	1:A:246:ILE:HG21	1.56	0.46
1:A:94:GLU:OE2	1:A:138:LYS:HE3	2.16	0.46
1:A:126:LEU:HD22	1:A:180:ILE:CG2	2.47	0.45
1:A:107:GLY:O	1:A:111:GLY:N	2.49	0.45
1:A:280:SER:O	1:A:283:GLU:HB2	2.16	0.45
1:B:121:VAL:HB	2:B:2300:NDP:H3D	1.97	0.45
1:A:270:ASN:HB3	1:B:270:ASN:HB2	1.98	0.45
1:B:63:VAL:CG1	1:B:71:LEU:HD22	2.47	0.45
1:B:270:ASN:HA	1:B:271:PRO:HD2	1.60	0.44
1:B:293:GLU:HG2	3:B:2340:HOH:O	2.18	0.44
1:B:264:VAL:N	1:B:265:PRO:HD2	2.33	0.44
1:B:242:ALA:O	1:B:246:ILE:HG12	2.18	0.44
1:B:133:ILE:HD11	3:B:2316:HOH:O	2.17	0.44
1:B:206:VAL:HA	3:B:2406:HOH:O	2.17	0.43
1:A:30:GLU:CD	1:A:30:GLU:H	2.21	0.43
1:B:174:LYS:C	1:B:175:ILE:HD12	2.39	0.43
1:B:127:THR:HG23	1:B:128:PHE:O	2.19	0.43
1:A:143:ASN:HB2	1:A:186:SER:HB2	2.01	0.42
3:A:1500:HOH:O	1:B:145:HIS:CD2	2.73	0.42
1:B:56:LYS:HG3	1:B:81:LEU:HG	2.02	0.42
1:A:143:ASN:N	1:A:143:ASN:HD22	2.17	0.42
1:B:200:GLU:O	1:B:204:ASN:HB2	2.19	0.42
1:A:196:THR:O	1:A:199:SER:HB2	2.20	0.41
1:A:215:LEU:HB2	2:A:1300:NDP:C5N	2.49	0.41
1:A:246:ILE:HG23	1:A:246:ILE:HD13	1.66	0.41
1:A:63:VAL:HG13	1:A:71:LEU:CD2	2.47	0.41
1:A:270:ASN:HA	1:A:271:PRO:HD2	1.77	0.41
1:B:175:ILE:HD13	1:B:268:LEU:CD2	2.51	0.41
1:A:126:LEU:HD22	1:A:180:ILE:HG23	2.03	0.41
1:A:175:ILE:HG22	1:A:177:TYR:CE1	2.56	0.41
1:A:175:ILE:HG13	1:B:273:ARG:HB2	2.03	0.41
1:A:176:THR:HG21	1:B:195:SER:HB2	2.03	0.40
1:B:128:PHE:O	1:B:130:HIS:CD2	2.73	0.40
1:B:180:ILE:O	1:B:180:ILE:HG13	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/295 (92%)	260 (96%)	10 (4%)	2 (1%)	22	39
1	B	272/295 (92%)	263 (97%)	9 (3%)	0	100	100
All	All	544/590 (92%)	523 (96%)	19 (4%)	2 (0%)	34	54

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	65	ALA
1	A	270	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/242 (93%)	204 (91%)	20 (9%)	9	19
1	B	224/242 (93%)	199 (89%)	25 (11%)	6	11
All	All	448/484 (93%)	403 (90%)	45 (10%)	7	15

All (45) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	63	VAL
1	A	69	GLU
1	A	72	GLN
1	A	81	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	125	ARG
1	A	133	ILE
1	A	158	LEU
1	A	175	ILE
1	A	179	LEU
1	A	199	SER
1	A	220	THR
1	A	225	LYS
1	A	239	GLU
1	A	246	ILE
1	A	251	LEU
1	A	252	ARG
1	A	262	ARG
1	A	273	ARG
1	A	295	LEU
1	A	296	TYR
1	B	36	LYS
1	B	61	VAL
1	B	63	VAL
1	B	69	GLU
1	B	81	LEU
1	B	122	LEU
1	B	126	LEU
1	B	127	THR
1	B	133	ILE
1	B	158	LEU
1	B	162	GLN
1	B	175	ILE
1	B	199	SER
1	B	203	VAL
1	B	220	THR
1	B	246	ILE
1	B	251	LEU
1	B	259	VAL
1	B	261	SER
1	B	262	ARG
1	B	273	ARG
1	B	274	LYS
1	B	294	LYS
1	B	295	LEU
1	B	296	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10)

such sidechains are listed below:

Mol	Chain	Res	Type
1	A	87	HIS
1	A	119	ASN
1	A	120	HIS
1	A	143	ASN
1	B	72	GLN
1	B	119	ASN
1	B	120	HIS
1	B	130	HIS
1	B	143	ASN
1	B	270	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NDP	B	2300	-	45,52,52	2.02	6 (13%)	53,80,80	1.33	6 (11%)
2	NDP	A	1300	-	45,52,52	2.04	6 (13%)	53,80,80	1.45	9 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	B	2300	-	-	5/30/77/77	0/5/5/5
2	NDP	A	1300	-	-	9/30/77/77	0/5/5/5

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1300	NDP	O7N-C7N	9.21	1.46	1.24
2	B	2300	NDP	O7N-C7N	8.72	1.45	1.24
2	B	2300	NDP	C7N-N7N	5.31	1.47	1.33
2	A	1300	NDP	C4N-C3N	-5.12	1.39	1.49
2	B	2300	NDP	C4N-C3N	-5.05	1.40	1.49
2	A	1300	NDP	C7N-N7N	4.03	1.44	1.33
2	A	1300	NDP	C4N-C5N	-3.97	1.38	1.48
2	B	2300	NDP	C6N-C5N	3.92	1.40	1.33
2	A	1300	NDP	C6N-C5N	3.46	1.39	1.33
2	B	2300	NDP	C4N-C5N	-3.43	1.39	1.48
2	A	1300	NDP	C5A-C4A	2.74	1.48	1.40
2	B	2300	NDP	C5A-C4A	2.17	1.46	1.40

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1300	NDP	N3A-C2A-N1A	-5.03	120.82	128.68
2	B	2300	NDP	O4D-C1D-N1N	3.95	115.77	108.06
2	B	2300	NDP	N3A-C2A-N1A	-3.70	122.90	128.68
2	B	2300	NDP	C4A-C5A-N7A	-2.93	106.35	109.40
2	B	2300	NDP	C3N-C2N-N1N	-2.88	118.98	123.10
2	A	1300	NDP	C2A-N1A-C6A	2.88	123.68	118.75
2	B	2300	NDP	PN-O3-PA	-2.79	123.25	132.83
2	A	1300	NDP	C3N-C2N-N1N	-2.66	119.29	123.10
2	A	1300	NDP	O4D-C1D-N1N	2.57	113.09	108.06
2	A	1300	NDP	O2B-C2B-C1B	2.49	119.08	110.10
2	A	1300	NDP	PN-O3-PA	-2.48	124.30	132.83
2	B	2300	NDP	O3X-P2B-O2X	2.45	117.00	107.64
2	A	1300	NDP	C4A-C5A-N7A	-2.39	106.91	109.40
2	A	1300	NDP	C3B-C2B-C1B	-2.04	99.05	102.89
2	A	1300	NDP	O2D-C2D-C1D	2.02	116.78	110.02

There are no chirality outliers.

All (14) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1300	NDP	C5D-O5D-PN-O1N
2	A	1300	NDP	O4D-C1D-N1N-C6N
2	B	2300	NDP	O4D-C1D-N1N-C6N
2	A	1300	NDP	C5D-O5D-PN-O3
2	B	2300	NDP	PN-O3-PA-O2A
2	A	1300	NDP	C5D-O5D-PN-O2N
2	B	2300	NDP	PN-O3-PA-O1A
2	A	1300	NDP	C1B-C2B-O2B-P2B
2	A	1300	NDP	C3B-C2B-O2B-P2B
2	B	2300	NDP	O4B-C4B-C5B-O5B
2	B	2300	NDP	C2B-O2B-P2B-O3X
2	A	1300	NDP	PN-O3-PA-O1A
2	A	1300	NDP	PN-O3-PA-O2A
2	A	1300	NDP	O4B-C4B-C5B-O5B


There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	2300	NDP	3	0
2	A	1300	NDP	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.


6.3 Carbohydrates [i](#)


Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.