



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 15, 2024 – 06:52 PM EDT

PDB ID : 1XT3  
Title : Structure Basis of Venom Citrate-Dependent Heparin Sulfate-Mediated Cell Surface Retention of Cobra Cardiotoxin A3  
Authors : Lee, S.-C.; Guan, H.-H.; Wang, C.-H.; Huang, W.-N.; Chen, C.-J.; Wu, W.-G.  
Deposited on : 2004-10-21  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	60	52% 47% .
1	B	60	42% 52% 7%
2	C	6	33% 67%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	IDS	C	4	-	-	X	-
2	IDS	C	6	X	-	-	-

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 1058 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytotoxin 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	60	Total 465	C 299	N 79	O 77	S 10	0	0	0
1	B	60	Total 465	C 299	N 79	O 77	S 10	0	0	0

- Molecule 2 is an oligosaccharide called 2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	6	Total 105	C 36	N 3	O 57	S 9	0	0	0

- Molecule 3 is CITRIC ACID (three-letter code: CIT) (formula: C<sub>6</sub>H<sub>8</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 13 6 7	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	9	Total O 9 9	0	0
4	B	1	Total O 1 1	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: Cytotoxin 3

Chain A: 



- Molecule 1: Cytotoxin 3

Chain B: 



- Molecule 2: 2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose-(1-4)-2-O-sulfo-alpha-L-idopyranuronic acid-(1-4)-2-deoxy-6-O-sulfo-2-(sulfoamino)-alpha-D-glucopyranose

Chain C: 



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	43.90Å 59.45Å 98.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 2.40	Depositor
% Data completeness (in resolution range)	(Not available) (25.00-2.40)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.04	Depositor
Refinement program	CNS	Depositor
R, $R_{free}$	0.227 , 0.250	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	1058	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	55.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IDS, CIT, SGN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.33	0/474	0.56	0/637
1	B	0.29	0/474	0.56	0/637
All	All	0.31	0/948	0.56	0/1274

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	465	0	494	26	0
1	B	465	0	494	38	0
2	C	105	0	32	9	0
3	A	13	0	5	1	1
4	A	9	0	0	0	0
4	B	1	0	0	0	0
All	All	1058	0	1025	66	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:6:LEU:HB2	2:C:6:IDS:O2S	1.77	0.83
1:A:33:PRO:HG3	1:A:36:ARG:HH12	1.43	0.81
2:C:3:SGN:O3	2:C:4:IDS:H5	1.83	0.78
1:B:18:LYS:HZ2	1:B:40:ASP:H	1.33	0.77
3:A:101:CIT:O4	1:B:31:LYS:HD3	1.85	0.74
1:B:33:PRO:HG3	1:B:36:ARG:NH1	2.04	0.72
1:A:37:GLY:HA3	2:C:6:IDS:O6B	1.91	0.70
1:B:20:LEU:HD12	1:B:40:ASP:O	1.96	0.66
1:A:33:PRO:HG3	1:A:36:ARG:NH1	2.12	0.64
1:A:24:MET:HB3	1:A:35:LYS:HB2	1.79	0.64
1:A:36:ARG:HH11	1:B:31:LYS:NZ	1.96	0.63
1:B:33:PRO:HG3	1:B:36:ARG:HH12	1.62	0.63
1:B:56:THR:HG22	1:B:57:ASP:H	1.64	0.62
1:B:45:SER:HB3	1:B:49:VAL:O	2.01	0.61
1:B:56:THR:HG22	1:B:57:ASP:N	2.20	0.56
1:B:18:LYS:NZ	1:B:40:ASP:H	2.02	0.55
1:A:43:PRO:HB2	1:A:51:TYR:CD2	2.42	0.54
1:B:23:LYS:HE3	1:B:25:PHE:HD2	1.73	0.54
1:B:25:PHE:CD1	1:B:30:PRO:HA	2.43	0.54
2:C:4:IDS:O2	2:C:5:SGN:N2	2.41	0.53
1:B:56:THR:HB	1:B:59:CYS:HB3	1.91	0.53
1:A:5:LYS:O	1:A:6:LEU:C	2.47	0.53
1:B:26:MET:O	1:B:28:ALA:N	2.42	0.52
1:B:2:LYS:HB3	1:B:58:ARG:HG2	1.90	0.52
1:A:39:ILE:HG13	1:A:40:ASP:N	2.24	0.52
1:B:26:MET:O	1:B:29:THR:N	2.29	0.52
1:A:2:LYS:HD3	1:A:11:TYR:CE2	2.44	0.52
1:B:23:LYS:HE3	1:B:25:PHE:CD2	2.45	0.52
1:A:56:THR:HG22	1:A:57:ASP:N	2.26	0.51
1:A:60:ASN:O	1:B:31:LYS:HE2	2.09	0.51
1:A:37:GLY:HA3	2:C:6:IDS:C6	2.41	0.50
1:A:43:PRO:HB2	1:A:51:TYR:CG	2.46	0.50
1:A:25:PHE:HD1	1:A:26:MET:O	1.95	0.50
1:B:40:ASP:CG	1:B:41:VAL:H	2.15	0.50
1:B:24:MET:O	1:B:34:VAL:HG22	2.12	0.49
1:B:55:ASN:H	1:B:55:ASN:HD22	1.61	0.49
1:B:39:ILE:HG12	1:B:43:PRO:HD3	1.94	0.49
1:B:24:MET:HB3	1:B:35:LYS:HB3	1.96	0.48
1:A:5:LYS:O	1:A:36:ARG:HD2	2.14	0.47
1:A:60:ASN:O	1:B:31:LYS:HG2	2.14	0.47
1:A:55:ASN:C	1:A:55:ASN:HD22	2.17	0.47

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:3:SGN:O3	2:C:4:IDS:C5	2.59	0.47
1:A:36:ARG:HH11	1:B:31:LYS:HZ3	1.60	0.46
1:B:32:VAL:HG12	1:B:32:VAL:O	2.16	0.46
1:B:27:VAL:HG12	1:B:48:LEU:O	2.16	0.46
1:B:18:LYS:HD2	1:B:39:ILE:C	2.36	0.45
2:C:4:IDS:H3	2:C:5:SGN:O3S	2.17	0.45
1:A:40:ASP:N	1:A:40:ASP:OD2	2.42	0.45
2:C:4:IDS:H3	2:C:5:SGN:S1	2.58	0.44
1:B:5:LYS:HE2	1:B:12:LYS:HB3	2.00	0.44
1:A:27:VAL:HG13	1:A:28:ALA:N	2.33	0.43
1:A:25:PHE:CD1	1:A:30:PRO:HA	2.53	0.43
1:B:41:VAL:O	1:B:41:VAL:HG13	2.18	0.42
1:B:18:LYS:HD2	1:B:39:ILE:CA	2.49	0.42
1:A:17:GLY:O	1:A:19:ASN:N	2.52	0.42
1:A:56:THR:O	1:A:59:CYS:SG	2.78	0.42
2:C:3:SGN:O4	2:C:4:IDS:O3	2.35	0.42
1:B:18:LYS:HZ3	1:B:40:ASP:HB3	1.85	0.42
1:A:15:PRO:O	1:A:16:ALA:C	2.58	0.41
1:B:26:MET:O	1:B:27:VAL:C	2.59	0.41
1:B:22:TYR:OH	1:B:35:LYS:HG2	2.21	0.41
1:A:24:MET:O	1:A:34:VAL:HG22	2.20	0.41
1:B:14:CYS:HA	1:B:15:PRO:HD2	1.95	0.41
1:B:31:LYS:O	1:B:33:PRO:HD2	2.21	0.41
1:B:55:ASN:HD22	1:B:55:ASN:N	2.18	0.40
1:B:5:LYS:C	1:B:7:VAL:N	2.73	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:101:CIT:O2	3:A:101:CIT:O2[3_555]	2.19	0.01

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	58/60 (97%)	48 (83%)	9 (16%)	1 (2%)	9	11
1	B	58/60 (97%)	46 (79%)	9 (16%)	3 (5%)	2	1
All	All	116/120 (97%)	94 (81%)	18 (16%)	4 (3%)	3	3

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	27	VAL
1	B	40	ASP
1	A	18	LYS
1	B	32	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	56/56 (100%)	55 (98%)	1 (2%)	59	76
1	B	56/56 (100%)	55 (98%)	1 (2%)	59	76
All	All	112/112 (100%)	110 (98%)	2 (2%)	59	76

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	55	ASN
1	B	55	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	55	ASN
1	B	55	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SGN	C	1	2	20,20,20	4.13	5 (25%)	25,31,31	1.21	2 (8%)
2	IDS	C	2	2	16,16,17	2.39	4 (25%)	16,24,26	2.69	4 (25%)
2	SGN	C	3	2	19,19,20	3.99	3 (15%)	23,29,31	1.57	4 (17%)
2	IDS	C	4	2	16,16,17	2.46	6 (37%)	16,24,26	1.23	1 (6%)
2	SGN	C	5	2	19,19,20	3.44	4 (21%)	23,29,31	1.68	5 (21%)
2	IDS	C	6	2	15,15,17	3.59	6 (40%)	14,22,26	2.93	5 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SGN	C	1	2	-	2/11/31/31	0/1/1/1
2	IDS	C	2	2	-	3/9/26/29	0/1/1/1
2	SGN	C	3	2	-	4/11/28/31	0/1/1/1
2	IDS	C	4	2	-	2/9/26/29	0/1/1/1
2	SGN	C	5	2	-	2/11/28/31	0/1/1/1
2	IDS	C	6	2	1/1/5/7	2/9/22/29	0/1/1/1

All (28) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1	SGN	S1-N2	15.25	1.78	1.59
2	C	3	SGN	S1-N2	14.33	1.77	1.59
2	C	5	SGN	S1-N2	12.28	1.75	1.59
2	C	6	IDS	C4-C5	-9.75	1.36	1.52
2	C	3	SGN	O6-S2	8.20	1.78	1.56
2	C	1	SGN	O6-S2	8.08	1.78	1.56
2	C	2	IDS	O2-S	7.16	1.78	1.57
2	C	6	IDS	O2-S	7.14	1.78	1.57
2	C	5	SGN	O6-S2	6.85	1.75	1.56
2	C	4	IDS	O2-S	6.66	1.77	1.57
2	C	6	IDS	C5-C6	4.93	1.58	1.52
2	C	4	IDS	O2-C2	-4.34	1.40	1.47
2	C	1	SGN	O1S-S1	3.44	1.46	1.42
2	C	3	SGN	O1S-S1	3.33	1.45	1.42
2	C	2	IDS	O2-C2	-3.29	1.42	1.47
2	C	1	SGN	O4-C4	3.22	1.50	1.43
2	C	4	IDS	O5-C5	3.02	1.49	1.43
2	C	1	SGN	O2S-S1	2.93	1.45	1.42
2	C	2	IDS	C1-C2	2.88	1.56	1.51
2	C	5	SGN	O1S-S1	2.75	1.45	1.42
2	C	4	IDS	C4-C3	2.73	1.59	1.52
2	C	5	SGN	C2-N2	-2.64	1.43	1.47
2	C	4	IDS	C1-C2	2.58	1.55	1.51
2	C	2	IDS	C4-C5	2.54	1.57	1.53
2	C	6	IDS	O5-C5	-2.39	1.39	1.43
2	C	4	IDS	O6B-C6	-2.18	1.23	1.30
2	C	6	IDS	O3-C3	-2.14	1.38	1.43
2	C	6	IDS	O2-C2	-2.09	1.44	1.47

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	IDS	O4-C4-C3	9.01	131.61	110.38
2	C	6	IDS	O5-C5-C4	8.52	122.22	111.16
2	C	3	SGN	O1S-S1-O2S	-4.20	111.03	120.36
2	C	1	SGN	O1S-S1-O2S	-3.98	111.50	120.36
2	C	6	IDS	C1-O5-C5	3.85	120.06	113.81
2	C	5	SGN	C1-O5-C5	3.64	117.07	112.19
2	C	4	IDS	O3S-S-O2	3.63	114.71	106.37
2	C	5	SGN	O1S-S1-O2S	-3.57	112.43	120.36
2	C	2	IDS	O4-C4-C5	-3.44	101.91	109.76
2	C	5	SGN	C3-C2-N2	-3.33	105.94	110.32

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	5	SGN	C3-C4-C5	-2.82	105.11	110.23
2	C	6	IDS	O6B-C6-C5	2.78	119.93	112.71
2	C	2	IDS	C3-C4-C5	2.64	113.84	109.30
2	C	6	IDS	C1-C2-C3	2.56	113.45	109.84
2	C	3	SGN	O2S-S1-N2	-2.54	104.64	108.88
2	C	2	IDS	O5-C1-C2	2.53	114.63	109.51
2	C	5	SGN	C4-C3-C2	-2.42	107.48	111.02
2	C	1	SGN	O5-C5-C6	2.20	111.05	106.69
2	C	3	SGN	O6S-S2-O6	2.17	111.35	106.37
2	C	6	IDS	O3-C3-C2	-2.12	105.67	110.08
2	C	3	SGN	O6-C6-C5	2.06	111.23	107.57

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	C	6	IDS	C5

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	1	SGN	C1-C2-N2-S1
2	C	2	IDS	C2-O2-S-O1S
2	C	2	IDS	C2-O2-S-O3S
2	C	3	SGN	C4-C5-C6-O6
2	C	3	SGN	O5-C5-C6-O6
2	C	4	IDS	C1-C2-O2-S
2	C	4	IDS	C3-C2-O2-S
2	C	5	SGN	C4-C5-C6-O6
2	C	5	SGN	O5-C5-C6-O6
2	C	6	IDS	C1-C2-O2-S
2	C	2	IDS	C2-O2-S-O2S
2	C	3	SGN	C2-N2-S1-O1S
2	C	3	SGN	C5-C6-O6-S2
2	C	1	SGN	C5-C6-O6-S2
2	C	6	IDS	C4-C5-C6-O6A

There are no ring outliers.

4 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	6	IDS	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	3	SGN	3	0
2	C	5	SGN	3	0
2	C	4	IDS	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	CIT	A	101	-	12,12,12	3.64	5 (41%)	17,17,17	1.64	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CIT	A	101	-	-	5/16/16/16	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	101	CIT	C4-C3	9.17	1.65	1.54
3	A	101	CIT	C3-C6	-7.29	1.45	1.53
3	A	101	CIT	C4-C5	2.40	1.58	1.50
3	A	101	CIT	O4-C5	-2.19	1.23	1.30
3	A	101	CIT	C2-C3	-2.07	1.51	1.54

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	101	CIT	O7-C3-C6	-3.30	104.27	108.96
3	A	101	CIT	O6-C6-C3	3.05	118.98	113.14
3	A	101	CIT	O5-C6-C3	-2.54	117.17	122.09
3	A	101	CIT	O7-C3-C4	-2.18	104.40	109.38
3	A	101	CIT	C3-C4-C5	2.12	119.71	113.92

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	101	CIT	C2-C3-C6-O5
3	A	101	CIT	C2-C3-C6-O6
3	A	101	CIT	O7-C3-C6-O5
3	A	101	CIT	O7-C3-C6-O6
3	A	101	CIT	C6-C3-C4-C5

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	101	CIT	1	1

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.