



Full wwPDB NMR Structure Validation Report ⓘ

May 28, 2020 – 09:05 pm BST

PDB ID : 1XU0
Title : Solution structure of Xenopus leavis prion protein
Authors : Perez, D.R.; Wuthrich, K.
Deposited on : 2004-10-25

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	130	

2 Ensemble composition and analysis

This entry contains 20 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:125-A:226 (102)	0.63	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 4 single-model clusters were found.

Cluster number	Models
1	1, 2, 7, 12, 13, 14, 15, 19
2	3, 5, 8, 9, 20
3	4, 11, 17
Single-model clusters	6; 10; 16; 18

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1653 atoms, of which 803 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called prion protein.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	102	1653	527	803	149	165	9	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	97	GLY	-	CLONING ARTIFACT	UNP Q5S1W7

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: prion protein




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1 (medoid)


- Molecule 1: prion protein



4.2.2 Score per residue for model 2


- Molecule 1: prion protein






4.2.3 Score per residue for model 3

- Molecule 1: prion protein




4.2.4 Score per residue for model 4

- Molecule 1: prion protein




4.2.5 Score per residue for model 5

- Molecule 1: prion protein




4.2.6 Score per residue for model 6

- Molecule 1: prion protein




4.2.7 Score per residue for model 7

- Molecule 1: prion protein




4.2.8 Score per residue for model 8

- Molecule 1: prion protein




4.2.9 Score per residue for model 9

- Molecule 1: prion protein




4.2.10 Score per residue for model 10

- Molecule 1: prion protein




4.2.11 Score per residue for model 11

- Molecule 1: prion protein




4.2.12 Score per residue for model 12

- Molecule 1: prion protein




4.2.13 Score per residue for model 13

- Molecule 1: prion protein




4.2.14 Score per residue for model 14

- Molecule 1: prion protein




4.2.15 Score per residue for model 15

- Molecule 1: prion protein




4.2.16 Score per residue for model 16

- Molecule 1: prion protein




4.2.17 Score per residue for model 17

- Molecule 1: prion protein




4.2.18 Score per residue for model 18

- Molecule 1: prion protein




4.2.19 Score per residue for model 19

- Molecule 1: prion protein



4.2.20 Score per residue for model 20

- Molecule 1: prion protein



5 Refinement protocol and experimental data overview i

The models were refined using the following method: *torsion angle dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DYANA	structure solution	6.01
OPALP	refinement	

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

COVALENT-GEOMETRY INFOmissingINFO

5.1 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	850	803	804	3±2
All	All	17000	16060	16080	55

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:186:VAL:HG13	1:A:190:ILE:HD12	0.94	1.38	17	6
1:A:205:LEU:O	1:A:209:VAL:HG13	0.57	2.00	6	1
1:A:130:LEU:HD22	1:A:160:ARG:HD2	0.56	1.77	11	9
1:A:186:VAL:CG2	1:A:190:ILE:HD12	0.55	2.30	11	6
1:A:139:TYR:CE2	1:A:216:GLU:HB3	0.51	2.41	15	1
1:A:175:ASP:HA	1:A:178:VAL:HG12	0.50	1.83	10	2
1:A:149:TYR:CE2	1:A:157:MET:CE	0.50	2.94	8	1
1:A:205:LEU:O	1:A:209:VAL:HG12	0.49	2.07	11	1
1:A:181:CYS:SG	1:A:218:CYS:SG	0.49	3.11	5	1
1:A:141:PHE:CE1	1:A:209:VAL:HG23	0.48	2.42	11	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:130:LEU:HD22	1:A:160:ARG:CD	0.48	2.38	19	2
1:A:178:VAL:HG23	1:A:218:CYS:HB2	0.46	1.87	9	1
1:A:139:TYR:CE2	1:A:216:GLU:CB	0.45	3.00	12	1
1:A:158:PRO:HB3	1:A:189:TYR:CE2	0.44	2.48	20	2
1:A:182:TYR:O	1:A:186:VAL:HG13	0.44	2.13	18	1
1:A:137:MET:SD	1:A:139:TYR:CD2	0.44	3.10	20	1
1:A:182:TYR:O	1:A:186:VAL:HG12	0.44	2.13	13	2
1:A:134:VAL:HG11	1:A:217:MET:SD	0.43	2.54	14	1
1:A:158:PRO:HB3	1:A:189:TYR:CZ	0.43	2.48	8	1
1:A:149:TYR:CD2	1:A:209:VAL:HG11	0.43	2.49	11	1
1:A:149:TYR:CZ	1:A:206:ASP:HA	0.43	2.49	5	2
1:A:183:ASN:HA	1:A:186:VAL:HG22	0.42	1.90	16	1
1:A:149:TYR:CE1	1:A:157:MET:CE	0.42	3.02	15	1
1:A:137:MET:CE	1:A:217:MET:HG3	0.42	2.45	9	1
1:A:151:ASN:HD22	1:A:152:ASP:N	0.42	2.13	19	1
1:A:137:MET:CE	1:A:217:MET:HG2	0.41	2.46	4	2
1:A:132:ASN:ND2	1:A:224:ARG:HH12	0.41	2.14	8	1
1:A:149:TYR:CE1	1:A:206:ASP:HA	0.40	2.51	5	1
1:A:177:PHE:CD1	1:A:222:TYR:CE1	0.40	3.09	8	1
1:A:178:VAL:HG23	1:A:218:CYS:CB	0.40	2.46	10	1
1:A:130:LEU:HD13	1:A:160:ARG:CG	0.40	2.46	17	1

5.2 Torsion angles [i](#)

5.2.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	100/130 (77%)	81±2 (81±2%)	14±2 (14±2%)	5±2 (5±2%)	4	27
All	All	2000/2600 (77%)	1620 (81%)	288 (14%)	92 (5%)	4	27

All 23 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	200	SER	12

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	196	GLY	11
1	A	168	GLY	7
1	A	170	GLU	7
1	A	137	MET	6
1	A	197	LYS	5
1	A	132	ASN	5
1	A	130	LEU	5
1	A	139	TYR	5
1	A	140	GLN	4
1	A	172	VAL	3
1	A	201	GLU	3
1	A	134	VAL	3
1	A	141	PHE	3
1	A	171	TYR	3
1	A	166	TYR	2
1	A	198	ASN	2
1	A	186	VAL	1
1	A	128	TYR	1
1	A	131	GLY	1
1	A	169	GLU	1
1	A	136	ARG	1
1	A	138	SER	1

5.2.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	93/112 (83%)	76±3 (82±3%)	17±3 (18±3%)	4 38
All	All	1860/2240 (83%)	1529 (82%)	331 (18%)	4 38

All 69 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	215	ARG	18
1	A	220	THR	16
1	A	130	LEU	15

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	224	ARG	14
1	A	136	ARG	13
1	A	203	ASN	12
1	A	184	MET	11
1	A	140	GLN	10
1	A	200	SER	9
1	A	152	ASP	9
1	A	132	ASN	9
1	A	222	TYR	9
1	A	204	GLN	9
1	A	148	ARG	8
1	A	167	ARG	8
1	A	160	ARG	7
1	A	226	SER	6
1	A	163	ARG	6
1	A	159	ASN	6
1	A	156	GLN	6
1	A	180	ASP	6
1	A	161	VAL	6
1	A	138	SER	6
1	A	223	ARG	6
1	A	169	GLU	5
1	A	199	ASN	5
1	A	137	MET	5
1	A	179	ARG	4
1	A	153	TYR	4
1	A	134	VAL	4
1	A	173	SER	4
1	A	142	ASN	4
1	A	139	TYR	4
1	A	186	VAL	3
1	A	185	SER	3
1	A	212	GLN	3
1	A	189	TYR	3
1	A	201	GLU	3
1	A	221	GLU	3
1	A	187	THR	3
1	A	210	LYS	3
1	A	157	MET	2
1	A	208	THR	2
1	A	174	GLU	2
1	A	198	ASN	2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	205	LEU	2
1	A	175	ASP	2
1	A	147	SER	2
1	A	213	ILE	2
1	A	143	ASN	2
1	A	151	ASN	2
1	A	206	ASP	2
1	A	207	THR	2
1	A	202	LEU	2
1	A	129	MET	2
1	A	146	GLU	2
1	A	155	ASN	1
1	A	181	CYS	1
1	A	172	VAL	1
1	A	219	ILE	1
1	A	188	GLU	1
1	A	177	PHE	1
1	A	197	LYS	1
1	A	176	ARG	1
1	A	171	TYR	1
1	A	170	GLU	1
1	A	218	CYS	1
1	A	145	MET	1
1	A	183	ASN	1

5.2.3 RNA [i](#)

There are no RNA molecules in this entry.

5.3 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.4 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.5 Ligand geometry [i](#)

There are no ligands in this entry.

5.6 Other polymers [i](#)

There are no such molecules in this entry.

5.7 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Chemical shift validation

No chemical shift data were provided