

# Full wwPDB X-ray Structure Validation Report (i)

Jan 8, 2025 – 04:13 PM JST

PDB ID : 8XU7

Title: Crystal structure of Lsd18 in complex with a product

Authors : Deng, Y.M.; Chen, X.

Deposited on : 2024-01-12

Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.21

EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.004 (Gargrove)

Density-Fitness : 1.0.11

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.40

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}({\rm \AA})) \end{array}$
$R_{free}$	164625	5791 (2.20-2.20)
Clashscore	180529	6634 (2.20-2.20)
Ramachandran outliers	177936	6560 (2.20-2.20)
Sidechain outliers	177891	6561 (2.20-2.20)
RSRZ outliers	164620	5791 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	A	488	87%	8%	•	-
1	В	488	86%	9%	·	•

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	LSB	A	702	X	-	-	-
3	LSB	В	702	X	-	-	-
3	LSB	В	703	X	-	_	-



## 2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 7482 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Putative epoxidase LasC.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	468	Total 3485	C 2168	N 660	O 644	S 13	0	0	0
1	В	468	Total 3492	C 2172	N 663	O 645	S 12	0	1	0

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP B5M9L6
A	2	ASN	_	expression tag	UNP B5M9L6
A	3	HIS	_	expression tag	UNP B5M9L6
A	4	LYS	_	expression tag	UNP B5M9L6
A	5	VAL	-	expression tag	UNP B5M9L6
A	6	HIS	-	expression tag	UNP B5M9L6
A	7	HIS	-	expression tag	UNP B5M9L6
A	8	HIS	-	expression tag	UNP B5M9L6
A	9	HIS	-	expression tag	UNP B5M9L6
A	10	HIS	-	expression tag	UNP B5M9L6
A	11	HIS	-	expression tag	UNP B5M9L6
A	12	ILE	-	expression tag	UNP B5M9L6
A	13	GLU	-	expression tag	UNP B5M9L6
A	14	GLY	-	expression tag	UNP B5M9L6
A	15	ARG	-	expression tag	UNP B5M9L6
A	16	HIS	-	expression tag	UNP B5M9L6
В	1	MET	-	initiating methionine	UNP B5M9L6
В	2	ASN	-	expression tag	UNP B5M9L6
В	3	HIS	-	expression tag	UNP B5M9L6
В	4	LYS	-	expression tag	UNP B5M9L6
В	5	VAL	-	expression tag	UNP B5M9L6
В	6	HIS	-	expression tag	UNP B5M9L6
В	7	HIS	-	expression tag	UNP B5M9L6
В	8	HIS	-	expression tag	UNP B5M9L6
В	9	HIS		expression tag	UNP B5M9L6



n previous	paae
	n previous

Chain	Residue	Modelled	Actual	Comment	Reference
В	10	HIS	-	expression tag	UNP B5M9L6
В	11	HIS	-	expression tag	UNP B5M9L6
В	12	ILE	-	expression tag	UNP B5M9L6
В	13	GLU	-	expression tag	UNP B5M9L6
В	14	GLY	-	expression tag	UNP B5M9L6
В	15	ARG	-	expression tag	UNP B5M9L6
В	16	HIS	-	expression tag	UNP B5M9L6


• Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	
9	۸	1	Total	С	N	О	Р	0	0	
	А	A	1	53	27	9	15	2	0	
9	D	1	Total	С	N	О	Р	0	0	
	Б	1	53	27	9	15	2	U	U	

• Molecule 3 is (4R,5S)-3-((2R,3S,4S)-2-ethyl-5-((2R,3R)-2-ethyl-3-(2-((2R,3R)-2-ethyl-3-methyloxiran-2-yl)ethyl)oxiran-2-yl)-3-hydroxy-4-methylpentanoyl)-4-methyl-5-phenyloxazolidi n-2-one (three-letter code: LSB) (formula:  $C_{29}H_{43}NO_6$ ) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	5	ZeroOcc	AltConf	
3	Λ	1	Total C	О	0	0	
9	Λ	1	17 15	2	0		
3	В	1	Total C	О	0	0	
3	Б	1	17 15	2	0		
2	D	1	Total C	О	0	0	
)	Ъ	1	17 15	2			

• Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Cl 1 1	0	0
4	В	1	Total Cl 1 1	0	0

• Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	170	Total O 170 170	0	0
5	В	176	Total O 176 176	0	0



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Putative epoxidase LasC





## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants	47.04Å 62.23Å 76.29Å	Donositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$74.79^{\circ}$ $81.97^{\circ}$ $76.81^{\circ}$	Depositor
Resolution (Å)	19.61 - 2.20	Depositor
rtesolution (A)	19.61 - 2.20	EDS
% Data completeness	98.1 (19.61-2.20)	Depositor
(in resolution range)	97.9 (19.61-2.20)	EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.43 (at 2.19Å)	Xtriage
Refinement program	PHENIX (1.16_3549: ???)	Depositor
D D.	0.227 , 0.271	Depositor
$R, R_{free}$	0.227 , $0.271$	DCC
$R_{free}$ test set	1977 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.3	Xtriage
Anisotropy	0.572	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.40 , 49.9	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.52, < L^2>=0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	7482	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 18.62% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: LSB, FAD, ELY, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles		
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.27	0/3541	0.46	0/4824	
1	В	0.26	0/3550	0.47	0/4836	
All	All	0.26	0/7091	0.46	0/9660	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintenain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	В	0	2
All	All	0	4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	298	ALA	Mainchain
1	A	299	ELY	Mainchain
1	В	298	ALA	Mainchain
1	В	299	ELY	Mainchain



### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3485	0	3466	28	0
1	В	3492	0	3478	42	0
2	A	53	0	31	1	0
2	В	53	0	31	1	0
3	A	17	0	25	4	0
3	В	34	0	50	8	0
4	A	1	0	0	0	0
4	В	1	0	0	0	0
5	A	170	0	0	5	0
5	В	176	0	0	6	0
All	All	7482	0	7081	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:256:ILE:HG23	5:A:820:HOH:O	1.66	0.94
1:B:199:LEU:HB3	1:B:204:LEU:HD12	1.60	0.82
1:B:252:VAL:HG21	3:B:703:LSB:H34A	1.66	0.78
1:B:416:GLN:O	1:B:416:GLN:HG3	1.88	0.74
1:A:361:LEU:HD22	1:A:370:LEU:HD11	1.72	0.70
1:B:473:ARG:NH1	5:B:803:HOH:O	2.29	0.66
1:A:342:VAL:HA	3:A:702:LSB:H34A	1.78	0.66
1:B:242:HIS:CD2	1:B:416:GLN:NE2	2.64	0.65
1:A:239:HIS:CE1	3:A:702:LSB:H31A	2.34	0.63
1:B:347:MET:HB2	5:B:821:HOH:O	1.98	0.62
1:B:400:ARG:CG	1:B:400:ARG:HH11	2.14	0.61
1:B:419:THR:HA	3:B:702:LSB:H31	1.85	0.58
1:B:246:ARG:HG2	1:B:247:PRO:HD2	1.85	0.58
1:A:364:ARG:NH2	5:A:812:HOH:O	2.35	0.57
1:B:60:LEU:HD13	1:B:68:ARG:HH11	1.70	0.57
1:A:252:VAL:HG21	3:A:702:LSB:H20	1.86	0.57
1:A:216:MET:O	1:A:312:THR:HG21	2.07	0.55



 $Continued\ from\ previous\ page...$ 

Continuea from prev		Interatomic	Clash	
Atom-1	Atom-2	${\rm distance} \ ({\rm \AA})$	overlap (Å)	
1:B:332:LEU:HB3	1:B:378:ILE:HD13	1.88	0.54	
1:B:400:ARG:NH1	1:B:400:ARG:HG3	2.23	0.53	
1:B:418:VAL:HG13	3:B:702:LSB:H32B	1.91	0.53	
1:A:416:GLN:NE2	5:A:817:HOH:O	2.39	0.53	
1:B:449:MET:HG3	3:B:702:LSB:O33	2.09	0.52	
2:A:701:FAD:N5	3:A:702:LSB:H24	2.25	0.52	
1:B:434:ARG:NH1	5:B:816:HOH:O	2.42	0.52	
1:B:400:ARG:HH11	1:B:400:ARG:HA	1.75	0.51	
1:B:217:VAL:HG12	1:B:310:ARG:HG3	1.91	0.51	
1:B:217:VAL:HG21	1:B:272:GLU:HB3	1.93	0.51	
1:B:376:ARG:NH2	5:B:817:HOH:O	2.43	0.51	
1:A:207:PRO:HG3	1:A:319:TYR:CE1	2.46	0.50	
1:A:163:ARG:NH2	5:A:827:HOH:O	2.44	0.50	
1:B:217:VAL:HG23	1:B:273:PRO:HD2	1.92	0.50	
1:B:276:ASP:OD2	5:B:801:HOH:O	2.20	0.50	
1:B:400:ARG:CG	1:B:400:ARG:NH1	2.73	0.49	
1:B:400:ARG:HH11	1:B:400:ARG:CB	2.26	0.49	
1:B:239:HIS:CD2	3:B:703:LSB:H31A	2.47	0.49	
1:A:233:PHE:CG	1:A:234:PRO:HD2	2.48	0.48	
1:B:195:SER:O	1:B:199:LEU:HG	2.13	0.48	
1:B:115:GLN:OE1	1:B:427:THR:HA	2.13	0.48	
1:A:239:HIS:O	1:A:289:ARG:NH1	2.38	0.48	
1:B:252:VAL:HG21	3:B:703:LSB:H20A	1.95	0.47	
1:B:213:ASP:HB3	1:B:400:ARG:HB3	1.97	0.47	
1:A:419:THR:HA	1:A:422:VAL:HG22	1.97	0.46	
1:B:415:ARG:NH1	5:B:815:HOH:O	2.41	0.46	
1:A:216:MET:HE1	1:A:342:VAL:HB	1.98	0.46	
1:B:233:PHE:CG	1:B:234:PRO:HD2	2.52	0.45	
1:B:242:HIS:CD2	1:B:416:GLN:HE21	2.34	0.44	
1:B:242:HIS:CD2	1:B:416:GLN:HE22	2.33	0.44	
1:B:107:VAL:HG21	1:B:472:LEU:HD23	1.99	0.44	
1:A:211:PHE:HA	1:A:314:ASN:O	2.18	0.43	
1:A:257:GLU:HG2	1:A:260:ARG:HH21	1.84	0.43	
1:B:110:THR:HA	1:B:239:HIS:ND1	2.32	0.43	
1:A:98:HIS:CD2	1:A:255:PRO:HB2	2.52	0.43	
1:B:199:LEU:HA	1:B:202:LEU:HB2	2.00	0.43	
1:A:209:GLU:O	1:A:402:GLN:NE2	2.46	0.43	
1:A:216:MET:HA	1:A:268:THR:HA	2.01	0.43	
1:B:44:GLY:O	1:B:145:ARG:HD2	2.19	0.43	
1:A:26:LEU:HD21	1:A:167:VAL:HG21	2.01	0.42	
1:A:206:ALA:O	5:A:801:HOH:O	2.22	0.42	



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${ m distance}({ m \AA})$	$\text{overlap } (\text{\AA})$
1:B:342:VAL:O	3:B:703:LSB:H21	2.18	0.42
1:B:346:GLY:HA3	2:B:701:FAD:H1'2	2.02	0.42
1:B:228:ALA:O	1:B:479:ARG:HD2	2.19	0.42
1:A:216:MET:HE3	1:A:218:TYR:HE1	1.85	0.42
1:B:252:VAL:CG2	3:B:703:LSB:H34A	2.45	0.42
1:A:74:TRP:CE3	1:A:441:THR:HG22	2.55	0.42
1:A:115:GLN:OE1	1:A:427:THR:HA	2.20	0.41
1:B:202:LEU:HD23	1:B:202:LEU:HA	1.91	0.41
1:A:299:ELY:HT2B	1:A:299:ELY:HH1A	1.89	0.41
1:A:400:ARG:HA	1:A:400:ARG:HD3	1.90	0.41
1:B:98:HIS:CD2	1:B:255:PRO:HB2	2.56	0.41
1:A:216:MET:CE	1:A:342:VAL:HB	2.51	0.41
1:A:205:PRO:HG2	1:A:322:LEU:HD22	2.04	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	465/488 (95%)	454 (98%)	11 (2%)	0	100	100
1	В	466/488 (96%)	460 (99%)	6 (1%)	0	100	100
All	All	931/976 (95%)	914 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	350/376 (93%)	346 (99%)	4 (1%)	70 82		
1	В	350/376~(93%)	344 (98%)	6 (2%)	56 71		
All	All	700/752~(93%)	690 (99%)	10 (1%)	62 77		

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	163	ARG
1	A	224	ARG
1	A	315	ARG
1	A	400	ARG
1	В	62	LYS
1	В	199	LEU
1	В	243	ARG
1	В	310	ARG
1	В	400	ARG
1	В	417	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	В	242	HIS
1	В	416	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the



expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Dog	Link	Во	ond leng	ths	В	ond ang	gles
WIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	ELY	В	299	1	11,12,13	0.57	0	8,13,15	0.70	0
1	ELY	A	299	1	11,12,13	0.56	0	8,13,15	0.76	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	ELY	В	299	1	-	3/12/13/15	-
1	ELY	A	299	1	-	3/12/13/15	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	299	ELY	CT1-CH1-NZ-CE
1	В	299	ELY	CT1-CH1-NZ-CE
1	A	299	ELY	CT1-CH1-NZ-CH2
1	В	299	ELY	CT1-CH1-NZ-CH2
1	A	299	ELY	CA-CB-CG-CD
1	В	299	ELY	CA-CB-CG-CD

There are no ring outliers.

1 monomer is involved in 1 short contact:

$\mathbf{Mol}$	Chain	Res	Type	Clashes	Symm-Clashes
1	A	299	ELY	1	0

### 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.



### 5.6 Ligand geometry (i)

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Truss	Chain	Dag	Link	Вс	ond leng	ths	Bond angles		
MIOI	Type	Chain	Res	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FAD	A	701	-	53,58,58	0.46	0	68,89,89	0.48	1 (1%)
3	LSB	A	702	-	16,18,39	1.73	3 (18%)	14,28,59	2.89	6 (42%)
2	FAD	В	701	-	53,58,58	0.44	0	68,89,89	0.52	2 (2%)
3	LSB	В	702	-	16,18,39	1.80	3 (18%)	14,28,59	2.37	6 (42%)
3	LSB	В	703	-	16,18,39	1.76	3 (18%)	14,28,59	2.49	6 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	LSB	A	702	-	1/1/5/12	11/17/33/71	0/2/2/4
2	FAD	A	701	-	-	2/30/50/50	0/6/6/6
2	FAD	В	701	-	-	2/30/50/50	0/6/6/6
3	LSB	В	702	-	1/1/5/12	11/17/33/71	0/2/2/4
3	LSB	В	703	-	1/1/5/12	9/17/33/71	0/2/2/4

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(A)
3	В	702	LSB	O33-C18	4.61	1.61	1.46
3	В	702	LSB	O36-C22	4.42	1.60	1.46
3	В	703	LSB	O33-C18	4.39	1.60	1.46
3	В	703	LSB	O36-C22	4.38	1.60	1.46
3	A	702	LSB	O36-C22	4.37	1.60	1.46
3	A	702	LSB	O33-C18	4.24	1.60	1.46
3	В	702	LSB	O33-C19	-2.64	1.40	1.45
3	A	702	LSB	O33-C19	-2.27	1.40	1.45



Continued from previous page...

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\text{\AA})$	$Ideal(\AA)$	
3	В	703	LSB	O33-C19	-2.21	1.41	1.45	l

#### All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$\operatorname{Ideal}({}^{o})$
3	A	702	LSB	O33-C19-C20	-6.94	105.86	117.74
3	A	702	LSB	O36-C23-C24	-4.86	107.53	117.33
3	В	703	LSB	C35-C34-C22	4.60	124.40	114.95
3	В	703	LSB	O33-C19-C20	-4.39	110.22	117.74
3	В	702	LSB	O36-C23-C24	-4.25	108.76	117.33
3	A	702	LSB	C32-C31-C18	4.07	123.32	114.95
3	В	702	LSB	C35-C34-C22	3.93	123.03	114.95
3	В	703	LSB	C32-C31-C18	3.92	123.02	114.95
3	В	703	LSB	O36-C23-C24	-3.82	109.64	117.33
3	A	702	LSB	C35-C34-C22	3.74	122.63	114.95
3	В	702	LSB	C20-C19-C18	-3.43	120.31	125.34
3	В	702	LSB	C32-C31-C18	3.38	121.90	114.95
3	В	702	LSB	C18-C17-C16	2.86	122.43	117.69
3	В	702	LSB	C31-C18-C19	2.29	123.60	118.75
2	В	701	FAD	C5A-C6A-N6A	2.29	123.83	120.35
2	A	701	FAD	C5A-C6A-N6A	2.27	123.80	120.35
3	В	703	LSB	C20-C21-C22	2.22	118.33	114.39
3	A	702	LSB	C18-C17-C16	2.20	121.35	117.69
3	В	703	LSB	C31-C18-C19	2.16	123.32	118.75
2	В	701	FAD	P-O3P-PA	-2.07	125.72	132.83
3	A	702	LSB	C20-C19-C18	-2.07	122.31	125.34

#### All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	702	LSB	C22
3	В	702	LSB	C22
3	В	703	LSB	C22

#### All (35) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	В	701	FAD	O4'-C4'-C5'-O5'
3	A	702	LSB	C15-C16-C17-C18
3	A	702	LSB	C16-C17-C18-C31
3	A	702	LSB	C20-C21-C22-C23
3	A	702	LSB	C20-C21-C22-C34



Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	702	LSB	C21-C22-C34-C35
3	A	702	LSB	O36-C22-C34-C35
3	В	702	LSB	C15-C16-C17-C18
3	В	702	LSB	C19-C18-C31-C32
3	В	702	LSB	O33-C18-C31-C32
3	В	702	LSB	C20-C21-C22-C34
3	В	703	LSB	C17-C18-C31-C32
3	В	703	LSB	C21-C22-C34-C35
3	В	703	LSB	O36-C22-C34-C35
3	В	702	LSB	C20-C21-C22-O36
2	A	701	FAD	O4'-C4'-C5'-O5'
3	В	703	LSB	C15-C16-C17-C18
3	A	702	LSB	O33-C19-C20-C21
3	В	703	LSB	C20-C21-C22-O36
3	В	703	LSB	C20-C21-C22-C23
3	В	702	LSB	C18-C19-C20-C21
3	В	703	LSB	C16-C17-C18-C31
3	A	702	LSB	C18-C19-C20-C21
3	A	702	LSB	C17-C18-C31-C32
3	В	702	LSB	C17-C18-C31-C32
2	В	701	FAD	O4B-C4B-C5B-O5B
3	В	702	LSB	O33-C19-C20-C21
2	A	701	FAD	O4B-C4B-C5B-O5B
3	В	702	LSB	C16-C17-C18-O33
3	В	703	LSB	C16-C17-C18-O33
3	A	702	LSB	C19-C20-C21-C22
3	В	702	LSB	C19-C20-C21-C22
3	A	702	LSB	C16-C17-C18-O33
3	В	703	LSB	C19-C20-C21-C22
3	В	702	LSB	C16-C17-C18-C31

There are no ring outliers.


5 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	701	FAD	1	0
3	A	702	LSB	4	0
2	В	701	FAD	1	0
3	В	702	LSB	3	0
3	В	703	LSB	5	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,




bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














## 5.7 Other polymers (i)

There are no such residues in this entry.

Torsions



Rings

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\# \mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q<0.9
1	A	467/488 (95%)	0.41	7 (1%) 71 68	20, 31, 45, 59	0
1	В	467/488 (95%)	0.36	12 (2%) 57 54	19, 31, 46, 77	1 (0%)
All	All	934/976 (95%)	0.39	19 (2%) 64 61	19, 31, 46, 77	1 (0%)

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	412	ALA	3.9
1	В	411	HIS	3.9
1	В	484	ALA	3.1
1	A	412	ALA	3.1
1	В	409	THR	2.7
1	A	301	GLN	2.7
1	В	416	GLN	2.7
1	В	400	ARG	2.5
1	В	54	PHE	2.5
1	A	90	ASP	2.5
1	A	461	GLY	2.5
1	В	193	SER	2.4
1	A	251	ALA	2.4
1	В	418	VAL	2.3
1	В	194	PRO	2.1
1	В	343	TYR	2.1
1	A	217	VAL	2.1
1	A	312	THR	2.1
1	В	394	ILE	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,



median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
1	ELY	В	299	13/14	0.83	0.11	27,33,38,38	0
1	ELY	A	299	13/14	0.89	0.09	23,30,36,39	0

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
3	LSB	В	702	17/36	0.57	0.22	41,57,60,63	0
3	LSB	В	703	17/36	0.58	0.20	36,45,55,55	0
3	LSB	A	702	17/36	0.60	0.18	32,41,50,53	0
2	FAD	В	701	53/53	0.91	0.09	19,28,33,36	0
2	FAD	A	701	53/53	0.92	0.09	16,26,30,32	0
4	CL	A	703	1/1	0.97	0.06	32,32,32,32	0
4	CL	В	704	1/1	0.98	0.03	39,39,39,39	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




#### Electron density around LSB B 702:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










#### Electron density around LSB B 703:

 $2 \text{mF}_o\text{-DF}_c$  (at 0.7 rmsd) in gray  $\text{mF}_o\text{-DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)












#### Electron density around LSB A 702:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










#### Electron density around FAD B 701:


 $2 {
m mF}_o {
m -DF}_c$  (at 0.7 rmsd) in gray  ${
m mF}_o {
m -DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)











## 6.5 Other polymers (i)

There are no such residues in this entry.

