



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 10:23 am BST

PDB ID : 1XYZ
Title : A COMMON PROTEIN FOLD AND SIMILAR ACTIVE SITE IN TWO
DISTINCT FAMILIES OF BETA-GLYCANASES
Authors : Alzari, P.M.; Spinelli, S.; Dominguez, R.
Deposited on : 1995-06-07
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1812 (1.40-1.40)
Ramachandran outliers	138981	1763 (1.40-1.40)
Sidechain outliers	138945	1762 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	347	
1	B	347	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5621 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 1,4-BETA-D-XYLAN-XYLANOHYDROLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	320	2582	1640	439	484	19	0	0	0
1	B	320	2582	1640	439	484	19	0	0	0

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	234	Total 234	O 234	0	0
2	B	223	Total 223	O 223	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.


- Molecule 1: 1,4-BETA-D-XYLAN-XYLANOHYDROLASE

Chain A: 



- Molecule 1: 1,4-BETA-D-XYLAN-XYLANOHYDROLASE

Chain B: 



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	47.10Å 51.10Å 70.74Å 100.54° 83.79° 101.64°	Depositor
Resolution (Å)	10.00 – 1.40	Depositor
% Data completeness (in resolution range)	(Not available) (10.00-1.40)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.183 , 0.221	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	5621	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.68	0/2650	0.76	1/3597 (0.0%)
1	B	0.67	0/2650	0.75	0/3597
All	All	0.68	0/5300	0.76	1/7194 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	565	ASP	CB-CG-OD1	5.36	123.12	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2582	0	2463	24	0
1	B	2582	0	2463	22	0
2	A	234	0	0	15	0
2	B	223	0	0	8	0
All	All	5621	0	4926	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:562:MET:SD	2:A:193:HOH:O	2.11	1.07
1:B:558:CYS:SG	2:B:308:HOH:O	2.34	0.86
1:A:708:MET:SD	2:A:173:HOH:O	2.40	0.79
1:A:558:CYS:SG	2:A:75:HOH:O	2.42	0.78
1:A:706:LYS:HE2	2:A:208:HOH:O	1.83	0.77
1:A:568:GLN:HE21	1:A:572:ASN:H	1.30	0.77
1:A:828:ASN:ND2	2:A:104:HOH:O	2.17	0.76
1:B:631:THR:HG22	2:B:420:HOH:O	1.89	0.70
1:A:561:GLU:HB2	2:A:75:HOH:O	1.91	0.69
1:B:567:LEU:HD21	2:B:278:HOH:O	1.94	0.68
1:B:558:CYS:CB	2:B:308:HOH:O	2.42	0.66
1:A:535:TYR:HE1	1:A:578:LYS:NZ	1.94	0.65
1:A:535:TYR:HE1	1:A:578:LYS:HZ3	1.45	0.64
1:A:583:LEU:HD21	1:A:636:LYS:HD3	1.82	0.59
1:A:832:GLU:HA	1:A:835:MET:HE2	1.84	0.59
1:A:535:TYR:CE1	1:A:578:LYS:NZ	2.71	0.58
1:A:523:GLU:HG3	2:A:66:HOH:O	2.02	0.58
1:A:627:THR:O	1:A:631:THR:HG23	2.05	0.56
1:B:690:ASN:HB2	2:B:413:HOH:O	2.07	0.55
1:B:561:GLU:HB2	2:B:308:HOH:O	2.10	0.51
1:A:562:MET:CE	2:A:193:HOH:O	2.50	0.51
1:A:561:GLU:HG3	2:A:167:HOH:O	2.11	0.51
1:A:558:CYS:CB	2:A:75:HOH:O	2.58	0.49
1:B:558:CYS:HB3	2:B:308:HOH:O	2.08	0.49
2:A:393:HOH:O	1:B:561:GLU:HG3	2.13	0.49
1:B:759:ILE:HG23	1:B:763:GLU:OE2	2.12	0.48
1:A:595:GLY:HA2	2:A:193:HOH:O	2.13	0.48
1:A:568:GLN:NE2	1:A:572:ASN:H	2.05	0.48
1:B:761:GLN:HG2	1:B:762:SER:N	2.28	0.48
1:A:774:ASN:ND2	2:A:82:HOH:O	2.47	0.47
1:A:551:ARG:HD3	2:A:144:HOH:O	2.15	0.47
1:B:644:ASN:HD22	1:B:687:ASN:HB3	1.81	0.45
1:B:764:ASN:HD21	1:B:766:ALA:HB3	1.81	0.45
1:B:804:ILE:HB	1:B:805:PRO:HD3	1.98	0.45
1:B:627:THR:O	1:B:631:THR:HG23	2.17	0.44
1:B:777:LYS:HG3	1:B:833:ALA:HB2	1.98	0.44
1:B:562:MET:HE1	1:B:595:GLY:HA2	1.99	0.44
1:A:673:ARG:NH1	2:A:189:HOH:O	2.42	0.43
1:A:707:SER:O	1:A:711:ARG:HG3	2.19	0.43
1:B:764:ASN:ND2	1:B:766:ALA:HB3	2.33	0.43
1:A:536:PRO:HG2	1:A:801:TYR:CZ	2.54	0.42
1:B:687:ASN:HD22	1:B:719:GLY:HA3	1.84	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:813:ASN:N	1:B:814:PRO:HD3	2.35	0.41
1:B:760:PRO:HG3	1:B:811:TYR:CE2	2.56	0.41
1:B:548:ILE:HD12	1:B:821:TYR:CD1	2.55	0.41
1:B:601:HIS:HD2	2:B:250:HOH:O	2.05	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	318/347 (92%)	314 (99%)	4 (1%)	0	100	100
1	B	318/347 (92%)	312 (98%)	5 (2%)	1 (0%)	41	18
All	All	636/694 (92%)	626 (98%)	9 (1%)	1 (0%)	47	21

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	762	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	278/301 (92%)	277 (100%)	1 (0%)	91	78

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	278/301 (92%)	275 (99%)	3 (1%)	73	50
All	All	556/602 (92%)	552 (99%)	4 (1%)	84	66

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	761	GLN
1	B	539	ASN
1	B	762	SER
1	B	795	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	534	ASN
1	A	550	GLN
1	A	568	GLN
1	A	571	GLN
1	A	572	ASN
1	A	581	GLN
1	A	592	GLN
1	A	610	ASN
1	A	624	ASN
1	A	761	GLN
1	A	774	ASN
1	A	828	ASN
1	B	534	ASN
1	B	539	ASN
1	B	601	HIS
1	B	644	ASN
1	B	652	ASN
1	B	665	GLN
1	B	687	ASN
1	B	764	ASN
1	B	790	ASN
1	B	813	ASN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.