



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 14, 2025 – 02:39 PM JST

PDB ID : 8XZC  
Title : Crystal structure of human cytosolic beta-alanyl lysine dipeptidase (PM20D2)  
Tyr314Phe mutant  
Authors : Chandravanshi, K.; Kumar, A.; Makde, R.D.  
Deposited on : 2024-01-21  
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.21  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.004 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.40


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1460 (2.36-2.36)
Clashscore	180529	1571 (2.36-2.36)
Ramachandran outliers	177936	1559 (2.36-2.36)
Sidechain outliers	177891	1559 (2.36-2.36)
RSRZ outliers	164620	1460 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	454	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3120 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Xaa-Arg dipeptidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	401	3024	1945	513	553	13	0	4	0

There are 38 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-17	MET	-	initiating methionine	UNP Q8IYS1
A	-16	ALA	-	expression tag	UNP Q8IYS1
A	-15	SER	-	expression tag	UNP Q8IYS1
A	-14	TRP	-	expression tag	UNP Q8IYS1
A	-13	SER	-	expression tag	UNP Q8IYS1
A	-12	HIS	-	expression tag	UNP Q8IYS1
A	-11	PRO	-	expression tag	UNP Q8IYS1
A	-10	GLN	-	expression tag	UNP Q8IYS1
A	-9	PHE	-	expression tag	UNP Q8IYS1
A	-8	GLU	-	expression tag	UNP Q8IYS1
A	-7	GLY	-	expression tag	UNP Q8IYS1
A	-6	GLY	-	expression tag	UNP Q8IYS1
A	-5	SER	-	expression tag	UNP Q8IYS1
A	-4	SER	-	expression tag	UNP Q8IYS1
A	-3	HIS	-	expression tag	UNP Q8IYS1
A	-2	HIS	-	expression tag	UNP Q8IYS1
A	-1	HIS	-	expression tag	UNP Q8IYS1
A	0	HIS	-	expression tag	UNP Q8IYS1
A	1	HIS	-	expression tag	UNP Q8IYS1
A	2	HIS	-	expression tag	UNP Q8IYS1
A	3	SER	-	expression tag	UNP Q8IYS1
A	4	SER	-	expression tag	UNP Q8IYS1
A	5	GLY	-	expression tag	UNP Q8IYS1
A	6	SER	-	expression tag	UNP Q8IYS1
A	7	GLY	-	expression tag	UNP Q8IYS1
A	8	GLY	-	expression tag	UNP Q8IYS1
A	9	GLY	-	expression tag	UNP Q8IYS1

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	10	GLY	-	expression tag	UNP Q8IYS1
A	11	GLY	-	expression tag	UNP Q8IYS1
A	12	GLU	-	expression tag	UNP Q8IYS1
A	13	ASN	-	expression tag	UNP Q8IYS1
A	14	LEU	-	expression tag	UNP Q8IYS1
A	15	TYR	-	expression tag	UNP Q8IYS1
A	16	PHE	-	expression tag	UNP Q8IYS1
A	17	GLN	-	expression tag	UNP Q8IYS1
A	18	GLY	-	expression tag	UNP Q8IYS1
A	19	SER	-	expression tag	UNP Q8IYS1
A	314	PHE	TYR	engineered mutation	UNP Q8IYS1

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O P 5 4 1	0	0
2	A	1	Total O P 5 4 1	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Zn	0	0
			2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	84	Total	O	0	0
			84	84		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Xaa-Arg dipeptidase



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	79.94Å 115.53Å 124.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.82 – 2.35 36.82 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.8 (36.82-2.35) 99.9 (36.82-2.35)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.89 (at 2.34Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.196 , 0.240 0.196 , 0.240	Depositor DCC
$R_{free}$ test set	1187 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	50.4	Xtrriage
Anisotropy	0.351	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 56.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3120	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	62.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.89% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/3108	0.46	0/4240

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3024	0	2833	16	0
2	A	10	0	0	0	0
3	A	2	0	0	0	0
4	A	84	0	0	1	0
All	All	3120	0	2833	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:242:ARG:HA	1:A:245:MET:HG3	1.72	0.71
1:A:120:ILE:HG22	1:A:379:GLU:HG3	1.88	0.56
1:A:198:LEU:HB3	1:A:348:PRO:HG3	1.91	0.53
1:A:187:MET:O	1:A:366:PRO:HD2	2.10	0.51
1:A:369:HIS:CE1	1:A:371:GLY:H	2.30	0.48
1:A:105:PRO:HA	1:A:153:PRO:HB2	1.96	0.48
1:A:365:HIS:O	1:A:365:HIS:ND1	2.46	0.47
1:A:124:CYS:SG	1:A:189:HIS:CE1	3.07	0.47
1:A:93:PRO:HA	1:A:94:PRO:HD3	1.83	0.45
1:A:325:LYS:HA	1:A:328[B]:MET:HE2	2.00	0.44
1:A:426:LEU:HD23	1:A:426:LEU:HA	1.87	0.44
1:A:95:GLU:HB3	4:A:674:HOH:O	2.19	0.43
1:A:246:LYS:HB2	1:A:249:TRP:CE2	2.54	0.42
1:A:360:VAL:HG23	1:A:361:VAL:HG23	2.00	0.42
1:A:96:ALA:O	1:A:98:ALA:N	2.49	0.41
1:A:14:LEU:HB2	1:A:16:PHE:CD1	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	397/454 (87%)	380 (96%)	15 (4%)	2 (0%)	25 28

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	148	PRO
1	A	98	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	288/360 (80%)	285 (99%)	3 (1%)	73 84

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	114	TYR
1	A	132	VAL
1	A	222	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	369	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PO4	A	501	-	4,4,4	0.88	0	6,6,6	0.40	0
2	PO4	A	502	-	4,4,4	0.88	0	6,6,6	0.43	0

There are no bond length outliers.

There are no bond angle outliers.


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight  $> 250$  and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	401/454 (88%)	0.45	36 (8%) <b>17</b> <b>19</b>	29, 59, 98, 133	4 (0%)

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	349	SER	6.6
1	A	99	PRO	6.4
1	A	220	TYR	5.5
1	A	348	PRO	4.7
1	A	14	LEU	4.5
1	A	98	ALA	4.4
1	A	222	TRP	4.4
1	A	97	ARG	4.3
1	A	350	GLY	4.2
1	A	214	ALA	4.0
1	A	103	PRO	3.8
1	A	193	GLU	3.7
1	A	223	GLU	3.6
1	A	351[A]	SER	3.5
1	A	146	GLY	3.3
1	A	339	ILE	3.3
1	A	96	ALA	3.2
1	A	94	PRO	3.2
1	A	149	ARG	3.1
1	A	17	GLN	3.0
1	A	199	PRO	3.0
1	A	151	PRO	2.9
1	A	197	TYR	2.9
1	A	274	TYR	2.9
1	A	124	CYS	2.8
1	A	95	GLU	2.8
1	A	116	ALA	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	314	PHE	2.7
1	A	430	GLN	2.6
1	A	15	TYR	2.5
1	A	150	PRO	2.4
1	A	16	PHE	2.3
1	A	148	PRO	2.2
1	A	224	GLY	2.2
1	A	377	HIS	2.1
1	A	213	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	PO4	A	502	5/5	0.72	0.11	118,118,121,132	5
2	PO4	A	501	5/5	0.83	0.11	83,85,90,111	0
3	ZN	A	503	1/1	0.87	0.09	41,41,41,41	1
3	ZN	A	504	1/1	0.91	0.08	37,37,37,37	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around PO4 A 502:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around PO4 A 501:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around ZN A 503:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.