



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2024 – 08:10 PM JST

PDB ID : 8Y0G  
Title : Human beta-catenin crystal structure  
Authors : Tim, F.  
Deposited on : 2024-01-22  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtrriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	549	

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 3820 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Catenin beta-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	510	3820	2403	686	706	25	0	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Catenin beta-1



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.84Å 103.43Å 186.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.50 39.75 – 2.50	Depositor EDS
% Data completeness (in resolution range)	92.4 (50.00-2.50) 92.5 (39.75-2.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.58 (at 2.51Å)	Xtrriage
Refinement program	REFMAC 5.5	Depositor
R, $R_{free}$	0.238 , 0.287 0.246 , 0.286	Depositor DCC
$R_{free}$ test set	1030 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	58.8	Xtrriage
Anisotropy	0.033	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 40.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3820	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	72.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.04% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/3873	0.49	1/5270 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	159	LEU	CB-CA-C	5.73	121.09	110.20

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3820	0	3905	79	0
All	All	3820	0	3905	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:515:ARG:HH21	1:A:575:GLY:HA3	1.47	0.80
1:A:241:VAL:HA	1:A:244:LEU:HD22	1.65	0.77

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:504:TRP:HD1	1:A:562:GLU:HG3	1.57	0.70
1:A:153:ILE:HG23	1:A:194:MET:HE1	1.74	0.69
1:A:189:MET:HG2	1:A:226:GLU:HG2	1.73	0.69
1:A:567:GLU:N	1:A:567:GLU:OE1	2.27	0.68
1:A:176:HIS:HB2	1:A:213:CYS:HB2	1.81	0.63
1:A:475:HIS:CE1	1:A:477:GLU:HG2	2.37	0.59
1:A:299:ASP:O	1:A:303:ILE:HG13	2.03	0.59
1:A:618:LEU:HD22	1:A:631:ILE:HG12	1.86	0.58
1:A:188:ILE:HG23	1:A:194:MET:HG3	1.84	0.58
1:A:157:THR:HG22	1:A:194:MET:HE2	1.86	0.58
1:A:584:VAL:HG12	1:A:587:ARG:HH22	1.68	0.58
1:A:607:ILE:HG22	1:A:610:ILE:H	1.69	0.58
1:A:274:ARG:CZ	1:A:308:ASN:HD22	2.19	0.56
1:A:203:GLN:HA	1:A:242:LYS:HE3	1.88	0.56
1:A:202:MET:HG3	1:A:214:THR:HB	1.89	0.54
1:A:256:ILE:HG13	1:A:296:ILE:HG22	1.90	0.53
1:A:162:ASP:OD1	1:A:167:VAL:HB	2.08	0.53
1:A:453:ARG:HH21	1:A:457:ARG:NH2	2.07	0.52
1:A:213:CYS:O	1:A:217:THR:OG1	2.26	0.52
1:A:619:CYS:HA	1:A:658:VAL:HG22	1.92	0.52
1:A:202:MET:HE3	1:A:243:MET:HG3	1.93	0.51
1:A:244:LEU:HB3	1:A:281:LYS:HB3	1.93	0.51
1:A:169:ASN:C	1:A:169:ASN:HD22	2.13	0.50
1:A:413:ASP:HB3	1:A:416:VAL:HB	1.93	0.50
1:A:195:VAL:HA	1:A:198:ILE:HD12	1.91	0.50
1:A:408:LEU:O	1:A:411:SER:HB3	2.11	0.50
1:A:627:ALA:O	1:A:631:ILE:HD12	2.13	0.49
1:A:588:ILE:HG12	1:A:589:VAL:N	2.27	0.49
1:A:160:LEU:HD23	1:A:194:MET:HB2	1.95	0.48
1:A:202:MET:HB2	1:A:214:THR:CG2	2.44	0.48
1:A:604:TYR:CZ	1:A:639:PRO:HG3	2.48	0.48
1:A:179:SER:HB3	1:A:217:THR:HG23	1.96	0.48
1:A:384:THR:O	1:A:388:LEU:HG	2.12	0.48
1:A:182:GLU:HA	1:A:185:ARG:HG2	1.95	0.48
1:A:212:ARG:HG3	1:A:254:TYR:HE2	1.79	0.47
1:A:242:LYS:HG2	1:A:243:MET:HG2	1.96	0.47
1:A:162:ASP:HB3	1:A:168:VAL:HB	1.97	0.47
1:A:511:VAL:O	1:A:515:ARG:HG3	2.16	0.46
1:A:164:ASP:OD2	1:A:167:VAL:HG23	2.15	0.46
1:A:242:LYS:HE2	1:A:243:MET:HG2	1.98	0.46
1:A:374:SER:O	1:A:378:VAL:HG23	2.16	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:278:GLY:O	1:A:282:MET:HG3	2.17	0.45
1:A:202:MET:HB2	1:A:214:THR:HG21	1.99	0.45
1:A:308:ASN:OD1	1:A:310:GLU:HG2	2.17	0.45
1:A:338:TRP:O	1:A:342:ARG:HG3	2.17	0.45
1:A:600:VAL:HG22	1:A:636:ALA:HB2	1.99	0.44
1:A:497:LEU:HD22	1:A:506:LEU:HD21	1.98	0.44
1:A:437:MET:O	1:A:441:VAL:HG22	2.18	0.44
1:A:474:ARG:H	1:A:474:ARG:HH11	1.65	0.44
1:A:541:VAL:O	1:A:545:GLN:HG3	2.18	0.44
1:A:202:MET:CE	1:A:243:MET:HG3	2.48	0.44
1:A:458:GLU:OE2	1:A:503:HIS:ND1	2.44	0.43
1:A:212:ARG:NE	1:A:250:SER:OG	2.46	0.43
1:A:253:PHE:N	1:A:253:PHE:CD2	2.86	0.43
1:A:654:TYR:O	1:A:658:VAL:HG23	2.19	0.43
1:A:658:VAL:O	1:A:662:MET:HG3	2.18	0.43
1:A:435:LYS:HD3	1:A:475:HIS:HB2	1.99	0.43
1:A:181:LYS:HD3	1:A:182:GLU:N	2.33	0.42
1:A:515:ARG:NH2	1:A:575:GLY:HA3	2.26	0.42
1:A:181:LYS:HG3	1:A:184:SER:H	1.85	0.42
1:A:153:ILE:O	1:A:157:THR:HG23	2.20	0.42
1:A:171:ALA:O	1:A:175:VAL:HG23	2.20	0.42
1:A:537:VAL:O	1:A:541:VAL:HG23	2.20	0.42
1:A:188:ILE:HA	1:A:194:MET:SD	2.61	0.41
1:A:203:GLN:HG2	1:A:242:LYS:HZ2	1.84	0.41
1:A:513:LEU:O	1:A:517:LEU:HG	2.21	0.41
1:A:292:LYS:HB2	1:A:292:LYS:HE3	1.69	0.41
1:A:226:GLU:CD	1:A:229:LEU:HD22	2.41	0.41
1:A:198:ILE:HG23	1:A:214:THR:HG23	2.01	0.41
1:A:227:GLY:O	1:A:231:ILE:HG13	2.20	0.41
1:A:370:LEU:HA	1:A:378:VAL:HG13	2.02	0.41
1:A:436:MET:HG3	1:A:475:HIS:CE1	2.56	0.41
1:A:473:SER:HB2	1:A:474:ARG:NH1	2.36	0.41
1:A:165:GLN:HA	1:A:168:VAL:HG12	2.04	0.40
1:A:649:GLU:O	1:A:652:ALA:N	2.52	0.40
1:A:622:ALA:HB3	1:A:661:ARG:HD3	2.03	0.40
1:A:155:GLU:O	1:A:159:LEU:HG	2.22	0.40

There are no symmetry-related clashes.



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	506/549 (92%)	497 (98%)	8 (2%)	1 (0%)	47 68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	247	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	407/459 (89%)	384 (94%)	23 (6%)	20 39

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	159	LEU
1	A	160	LEU
1	A	162	ASP
1	A	169	ASN
1	A	186	HIS
1	A	189	MET
1	A	190	ARG
1	A	193	GLN
1	A	194	MET
1	A	219	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	228	LEU
1	A	232	PHE
1	A	242	LYS
1	A	244	LEU
1	A	288	LYS
1	A	390	ASP
1	A	411	SER
1	A	457	ARG
1	A	475	HIS
1	A	508	LYS
1	A	588	ILE
1	A	637	THR
1	A	643	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	169	ASN
1	A	440	GLN
1	A	578	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	510/549 (92%)	0.68	59 (11%) <b>4</b> <b>4</b>	42, 60, 134, 167	0

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	149	ALA	16.8
1	A	146	ALA	14.9
1	A	153	ILE	11.6
1	A	156	LEU	10.4
1	A	195	VAL	9.2
1	A	144	ASP	9.0
1	A	154	PRO	8.7
1	A	145	ASP	7.7
1	A	157	THR	7.6
1	A	150	THR	6.9
1	A	160	LEU	6.7
1	A	147	GLU	6.1
1	A	560	PHE	5.2
1	A	172	ALA	4.7
1	A	248	VAL	4.7
1	A	561	VAL	4.7
1	A	151	ARG	4.6
1	A	159	LEU	4.4
1	A	234	SER	4.2
1	A	148	LEU	4.2
1	A	152	ALA	4.2
1	A	663	SER	4.1
1	A	185	ARG	4.0
1	A	190	ARG	3.6
1	A	253	PHE	3.5
1	A	252	LEU	3.5
1	A	208	VAL	3.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	251	VAL	3.2
1	A	643	LEU	3.2
1	A	242	LYS	3.0
1	A	178	LEU	2.9
1	A	192	PRO	2.9
1	A	211	ALA	2.9
1	A	175	VAL	2.8
1	A	224	HIS	2.8
1	A	549	ARG	2.8
1	A	199	VAL	2.7
1	A	204	ASN	2.7
1	A	186	HIS	2.7
1	A	155	GLU	2.6
1	A	189	MET	2.5
1	A	515	ARG	2.5
1	A	194	MET	2.3
1	A	606	PRO	2.3
1	A	547	THR	2.3
1	A	477	GLU	2.2
1	A	170	LYS	2.2
1	A	213	CYS	2.2
1	A	233	LYS	2.2
1	A	167	VAL	2.1
1	A	174	MET	2.1
1	A	644	LEU	2.1
1	A	200	ARG	2.1
1	A	660	PHE	2.1
1	A	181	LYS	2.1
1	A	212	ARG	2.0
1	A	562	GLU	2.0
1	A	168	VAL	2.0
1	A	250	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.