



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2018 – 10:20 pm GMT

PDB ID : 1Y1T  
Title : Crystal Structure of the Uridine Phosphorylase from Salmonella Typhimurium at 1.77Å Resolution  
Authors : Gabdoulkhakov, A.G.; Dontsova, M.V.; Kachalova, G.S.; Betzel, C.; Ealick, S.E.; Mikhailov, A.M.  
Deposited on : 2004-11-19  
Resolution : 1.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	7886 (1.80-1.76)
Clashscore	122126	8858 (1.80-1.76)
Ramachandran outliers	120053	8764 (1.80-1.76)
Sidechain outliers	120020	8763 (1.80-1.76)
RSRZ outliers	108989	7741 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	253	
1	F	253	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	5001	-	X	-	-

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 3773 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uridine phosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	250	Total 1876	C 1174	N 330	O 360	S 12	0	0	0
1	F	218	Total 1607	C 1009	N 278	O 309	S 11	0	0	0

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 6	C 3	O 3	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	F	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	153	Total	O	0	0
			153	153		
4	F	126	Total	O	0	0
			126	126		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Uridine phosphorylase



- Molecule 1: Uridine phosphorylase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	150.82Å 150.82Å 47.84Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	25.40 – 1.77 25.39 – 1.77	Depositor EDS
% Data completeness (in resolution range)	97.3 (25.40-1.77) 97.4 (25.39-1.77)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.45 (at 1.77Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.184 , 0.205 0.183 , 0.203	Depositor DCC
$R_{free}$ test set	1925 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.9	Xtrriage
Anisotropy	0.478	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 64.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.027 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3773	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.16% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.36	0/1906	0.64	1/2584 (0.0%)
1	F	0.40	1/1629 (0.1%)	0.68	3/2208 (0.1%)
All	All	0.38	1/3535 (0.0%)	0.66	4/4792 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	2184	MET	C-N	6.84	1.49	1.34

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	2184	MET	O-C-N	6.34	132.85	122.70
1	F	2253	LEU	CA-C-O	-5.53	108.50	120.10
1	F	2184	MET	CA-C-N	-5.24	105.68	117.20
1	A	1253	LEU	CA-C-O	-5.14	109.30	120.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1876	0	1887	43	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	1607	0	1627	29	0
2	A	6	0	4	0	0
3	F	5	0	0	0	0
4	A	153	0	0	4	0
4	F	126	0	0	2	0
All	All	3773	0	3518	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1092:ILE:HD11	1:A:1241:ALA:HB1	1.53	0.90
1:A:1094:THR:HG21	1:A:1238:GLU:HG2	1.69	0.74
1:F:2069:ILE:HG23	1:F:2197:MET:HE1	1.74	0.69
1:F:2158:SER:HB3	1:F:2200:ALA:HB2	1.75	0.68
1:F:2038:MET:HG2	1:F:2057:LEU:HD21	1.77	0.66
1:A:1079:GLU:O	1:A:1083:GLN:HG3	1.97	0.65
1:A:1182:GLY:O	1:A:1185:GLU:HG3	1.97	0.64
1:A:1060:LYS:HD2	1:A:1253:LEU:HB3	1.80	0.62
1:A:1048:ARG:HH11	1:A:1048:ARG:HG2	1.65	0.62
1:A:1094:THR:CG2	1:A:1238:GLU:HG2	2.29	0.61
1:A:1194:ASN:HD22	1:A:1194:ASN:N	1.98	0.61
1:A:1032:GLU:HG3	1:A:1053:TRP:CH2	2.38	0.59
1:A:1048:ARG:HG3	4:A:3090:HOH:O	2.04	0.57
1:A:1049:GLU:HG2	1:F:2049:GLU:HG3	1.87	0.57
1:A:1030:ARG:HH22	1:A:1093:GLY:HA2	1.70	0.57
1:A:1220:ILE:HB	1:A:1233:THR:O	2.04	0.56
1:F:2069:ILE:CG2	1:F:2197:MET:HE1	2.35	0.56
1:A:1158:SER:HB3	1:A:1200:ALA:HB2	1.87	0.56
1:A:1030:ARG:NH2	1:A:1093:GLY:HA2	2.22	0.55
1:F:2235:LYS:HB2	1:F:2235:LYS:HZ3	1.70	0.55
1:A:1092:ILE:HD11	1:A:1241:ALA:CB	2.33	0.55
1:A:1049:GLU:HG2	1:F:2049:GLU:CD	2.29	0.53
1:A:1054:ARG:HG3	1:A:1063:ILE:HD13	1.91	0.52
1:A:1030:ARG:HA	1:A:1033:LYS:HD2	1.90	0.52
1:F:2147:ILE:HD11	1:F:2244:ILE:HD11	1.93	0.51
1:F:2186:GLU:HG3	1:F:2190:MET:HE3	1.93	0.51
1:F:2033:LYS:HE3	4:F:3145:HOH:O	2.09	0.51
1:F:2044:LEU:HD11	1:F:2054:ARG:HB2	1.93	0.51

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1005:ASP:OD2	1:A:1012:THR:HA	2.10	0.51
1:A:1044:LEU:HD11	1:A:1054:ARG:HB2	1.93	0.50
1:A:1034:ILE:HG12	1:A:1242:VAL:HG13	1.93	0.50
1:A:1076:ILE:HD11	1:F:2162:PHE:HD1	1.77	0.49
1:A:1058:ASP:OD2	1:A:1250:ARG:HG3	2.13	0.49
1:A:1137:THR:O	1:A:1141:VAL:HG23	2.13	0.49
1:F:2092:ILE:HD13	1:F:2093:GLY:N	2.27	0.49
1:F:2013:LYS:HG2	4:F:3173:HOH:O	2.13	0.49
1:F:2092:ILE:C	1:F:2092:ILE:HD13	2.34	0.48
1:F:2235:LYS:NZ	1:F:2235:LYS:HB2	2.29	0.47
1:A:1049:GLU:HG2	1:F:2049:GLU:CG	2.45	0.47
1:F:2105:GLY:HA2	1:F:2237:THR:HG22	1.96	0.47
1:F:2005:ASP:OD2	1:F:2013:LYS:HD2	2.15	0.47
1:A:1230:ASN:O	1:A:1233:THR:N	2.48	0.46
1:F:2242:VAL:O	1:F:2245:VAL:HG12	2.15	0.46
1:F:2111:THR:HG23	1:F:2153:VAL:HG12	1.98	0.45
1:A:1009:LEU:HG	1:A:1080:GLU:OE1	2.17	0.45
1:A:1185:GLU:HG2	4:A:3042:HOH:O	2.17	0.45
1:A:1238:GLU:O	1:A:1242:VAL:HG23	2.18	0.44
1:F:2096:GLY:HA2	1:F:2221:VAL:O	2.16	0.44
1:A:1194:ASN:ND2	1:A:1194:ASN:N	2.65	0.44
1:F:2024:VAL:O	1:F:2091:ARG:HG3	2.18	0.44
1:A:1031:VAL:HG13	1:A:1064:VAL:HG12	2.00	0.44
1:A:1025:PRO:O	1:A:1066:SER:HA	2.18	0.44
1:A:1054:ARG:HG3	1:A:1063:ILE:CD1	2.48	0.43
1:A:1016:LEU:HG	1:A:1063:ILE:HG13	2.00	0.43
1:F:2238:GLU:O	1:F:2242:VAL:HG23	2.19	0.43
1:A:1178:ARG:HD2	4:A:3209:HOH:O	2.18	0.42
1:F:2022:ALA:HA	1:F:2063:ILE:O	2.19	0.42
1:F:2034:ILE:HG12	1:F:2242:VAL:HG13	2.00	0.42
1:F:2162:PHE:O	1:F:2162:PHE:HD2	2.02	0.42
1:A:1247:GLU:HB3	4:A:3213:HOH:O	2.19	0.42
1:A:1163:TYR:HB2	1:A:1164:PRO:CD	2.50	0.42
1:A:1091:ARG:HH11	1:A:1091:ARG:HG3	1.86	0.41
1:A:1147:ILE:HG23	1:A:1147:ILE:O	2.20	0.41
1:F:2016:LEU:HD22	1:F:2063:ILE:CD1	2.50	0.41
1:F:2236:GLN:HB2	1:F:2236:GLN:HE21	1.65	0.41
1:A:1093:GLY:O	1:A:1217:ALA:HA	2.21	0.41
1:A:1037:LEU:HA	1:A:1037:LEU:HD22	1.92	0.40
1:A:1028:PRO:HG3	1:A:1049:GLU:OE2	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/253 (98%)	234 (94%)	10 (4%)	4 (2%)	11	2
1	F	212/253 (84%)	210 (99%)	1 (0%)	1 (0%)	31	15
All	All	460/506 (91%)	444 (96%)	11 (2%)	5 (1%)	16	4

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1235	LYS
1	F	2184	MET
1	A	1163	TYR
1	A	1231	ALA
1	A	1233	THR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	199/202 (98%)	190 (96%)	9 (4%)	30	13
1	F	171/202 (85%)	162 (95%)	9 (5%)	25	9
All	All	370/404 (92%)	352 (95%)	18 (5%)	27	10

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1005	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	1029	GLU
1	A	1037	LEU
1	A	1048	ARG
1	A	1049	GLU
1	A	1147	ILE
1	A	1185	GLU
1	A	1194	ASN
1	A	1196	GLU
1	F	2092	ILE
1	F	2161	THR
1	F	2162	PHE
1	F	2184	MET
1	F	2186	GLU
1	F	2195	TYR
1	F	2196	GLU
1	F	2235	LYS
1	F	2236	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1083	GLN
1	A	1122	HIS
1	A	1194	ASN
1	A	1209	GLN
1	A	1226	GLN
1	F	2236	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GOL	A	5001	-	5,5,5	4.72	5 (100%)	5,5,5	5.56	3 (60%)
3	SO4	F	4001	-	4,4,4	0.57	0	6,6,6	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	5001	-	-	0/4/4/4	0/0/0/0
3	SO4	F	4001	-	-	0/0/0/0	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	5001	GOL	C3-C2	-7.81	1.22	1.52
2	A	5001	GOL	C1-C2	-2.74	1.41	1.52
2	A	5001	GOL	O2-C2	-2.43	1.36	1.43
2	A	5001	GOL	O3-C3	3.48	1.57	1.42
2	A	5001	GOL	O1-C1	4.97	1.63	1.42

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	5001	GOL	O1-C1-C2	3.08	125.04	110.11
2	A	5001	GOL	O2-C2-C3	6.50	138.68	109.00
2	A	5001	GOL	O3-C3-C2	10.15	159.34	110.11

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	250/253 (98%)	0.48	15 (6%) 22 21	11, 21, 47, 78	0
1	F	218/253 (86%)	0.37	12 (5%) 25 24	10, 19, 41, 92	0
All	All	468/506 (92%)	0.43	27 (5%) 23 22	10, 20, 46, 92	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1007	PHE	8.3
1	F	2183	SER	7.7
1	A	1231	ALA	7.7
1	F	2162	PHE	7.2
1	A	1233	THR	6.8
1	A	1234	MET	6.2
1	F	2235	LYS	5.4
1	F	2237	THR	4.6
1	A	1048	ARG	4.5
1	F	2184	MET	4.2
1	A	1235	LYS	4.1
1	F	2185	GLU	3.9
1	A	1253	LEU	3.4
1	A	1170	ASP	3.3
1	F	2236	GLN	3.3
1	A	1232	GLU	3.2
1	F	2195	TYR	3.0
1	A	1040	LYS	3.0
1	A	1059	GLY	2.8
1	A	1004	SER	2.7
1	F	2238	GLU	2.6
1	F	2221	VAL	2.3
1	A	1005	ASP	2.3
1	A	1228	ILE	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	F	2186	GLU	2.1
1	A	1024	VAL	2.0
1	F	2101	HIS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GOL	A	5001	6/6	0.68	0.32	20,20,20,20	0
3	SO4	F	4001	5/5	0.89	0.38	20,20,20,20	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.