



Full wwPDB X-ray Structure Validation Report ⓘ

May 17, 2020 – 08:41 am BST

PDB ID : 4Y2A
Title : Crystal Structure of Coxsackie Virus B3 3D polymerase in complex with GPC-N114 inhibitor
Authors : Vives-Adrian, L.; Ferrer-Orta, C.; Verdaguer, N.
Deposited on : 2015-02-09
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	466	 83% 14%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	1FS	A	501	-	-	X	-

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 3844 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 3D polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	466	3739	2397	628	692	22	5	3	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	463	HIS	-	expression tag	UNP P03313
A	464	HIS	-	expression tag	UNP P03313
A	465	HIS	-	expression tag	UNP P03313
A	466	HIS	-	expression tag	UNP P03313

- Molecule 2 is 2,2'-[(4-chlorobenzene-1,2-diyl)bis(oxy)]bis(5-nitrobenzonitrile) (three-letter code: 1FS) (formula: C₂₀H₉ClN₄O₆).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Cl	N	O		
2	A	1	31	20	1	4	6	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	C O	0	0
			6	3 3		
4	A	1	Total	C O	0	0
			6	3 3		
4	A	1	Total	C O	0	0
			6	3 3		
4	A	1	Total	C O	0	0
			6	3 3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	25	Total	O	0	0
			25	25		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3D polymerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	74.74Å 74.74Å 289.05Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.36 – 2.90 74.74 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.7 (72.36-2.90) 99.7 (74.74-2.90)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.85 (at 2.91Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.198 , 0.250 0.201 , 0.201	Depositor DCC
R_{free} test set	1730 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	60.1	Xtriage
Anisotropy	0.050	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 35.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3844	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4, 1FS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.70	1/3852 (0.0%)	0.82	1/5214 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	428	HIS	CB-CG	-14.44	1.24	1.50

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	360	LYS	N-CA-C	-5.05	97.36	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	464	HIS	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3739	0	3689	39	0
2	A	31	0	9	12	0
3	A	25	0	0	1	0
4	A	24	0	32	0	0
5	A	25	0	0	1	0
All	All	3844	0	3730	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:501:1FS:H4	2:A:501:1FS:C2	1.94	0.95
1:A:174[A]:ARG:NH1	3:A:502:SO4:O1	2.24	0.70
2:A:501:1FS:H3	2:A:501:1FS:C20	2.23	0.69
1:A:188:ARG:HH11	2:A:501:1FS:H8	1.60	0.67
2:A:501:1FS:C2	2:A:501:1FS:C9	2.67	0.66
1:A:293:GLY:HA2	2:A:501:1FS:N1	2.14	0.62
1:A:188:ARG:HH11	2:A:501:1FS:C15	2.12	0.62
1:A:188:ARG:HD2	2:A:501:1FS:H8	1.83	0.60
1:A:194:LEU:HD23	1:A:194:LEU:C	2.23	0.58
2:A:501:1FS:C6	2:A:501:1FS:C20	2.83	0.56
1:A:265:TYR:O	1:A:268:TYR:HB3	2.06	0.56
2:A:501:1FS:H3	2:A:501:1FS:N4	2.23	0.52
1:A:188:ARG:HD2	2:A:501:1FS:C15	2.39	0.52
1:A:423:TRP:CZ2	1:A:424:HIS:CE1	2.98	0.52
1:A:161:GLU:OE2	1:A:174[A]:ARG:HG3	2.10	0.51
1:A:161:GLU:OE2	1:A:174[B]:ARG:HG3	2.10	0.50
1:A:314:TYR:O	1:A:317:ILE:HB	2.13	0.49
1:A:107:LEU:HD12	2:A:501:1FS:CL	2.50	0.49
1:A:210:VAL:HA	1:A:327:TYR:CE2	2.48	0.48
1:A:273:HIS:HE2	1:A:280:HIS:CD2	2.32	0.48
1:A:162:LEU:O	1:A:163:ARG:HD2	2.13	0.48
1:A:6:ILE:HG23	1:A:281:TYR:HB3	1.94	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:194:LEU:C	1:A:194:LEU:CD2	2.82	0.48
1:A:75:LEU:HD21	1:A:251:MET:HE3	1.96	0.48
1:A:410:ASN:O	1:A:411:THR:C	2.50	0.47
1:A:331:VAL:HG12	1:A:332:ILE:N	2.30	0.46
1:A:103:GLY:O	1:A:104:THR:HB	2.15	0.46
1:A:287:MET:HG2	1:A:298:ASN:HD21	1.81	0.45
1:A:306:ILE:HG21	1:A:322:PHE:HZ	1.82	0.45
1:A:146:ASP:OD1	1:A:146:ASP:N	2.50	0.45
1:A:183:ASP:O	1:A:184:SER:C	2.55	0.44
1:A:310:MET:HE3	1:A:344:LEU:HD11	1.99	0.44
1:A:359:ASP:C	1:A:360:LYS:O	2.51	0.44
1:A:412:GLN:O	1:A:416:ARG:HG3	2.17	0.44
1:A:142:LYS:O	1:A:146:ASP:OD1	2.35	0.44
1:A:311:LEU:HD21	1:A:319:LEU:HD11	2.00	0.44
1:A:147:LYS:NZ	5:A:601:HOH:O	2.51	0.43
1:A:194:LEU:HD23	1:A:194:LEU:O	2.17	0.43
2:A:501:1FS:C6	2:A:501:1FS:N4	2.81	0.43
1:A:115:SER:O	1:A:127:LYS:HE3	2.19	0.43
1:A:273:HIS:HE2	1:A:280:HIS:HE2	1.66	0.42
1:A:287:MET:HG2	1:A:298:ASN:ND2	2.34	0.42
1:A:368:THR:HB	1:A:370:THR:H	1.85	0.42
1:A:466:HIS:C	1:A:466:HIS:ND1	2.73	0.42
1:A:348:GLY:O	1:A:351:TYR:N	2.45	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	467/466 (100%)	433 (93%)	31 (7%)	3 (1%)	25 58

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	136	LYS
1	A	359	ASP
1	A	104	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	410/407 (101%)	376 (92%)	34 (8%)	11 32

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	ILE
1	A	22	LYS
1	A	72	GLU
1	A	96	LYS
1	A	105	GLU
1	A	108	GLU
1	A	110	LEU
1	A	128	ARG
1	A	146	ASP
1	A	151	ASN
1	A	174[A]	ARG
1	A	174[B]	ARG
1	A	179	SER
1	A	204	VAL
1	A	208	SER
1	A	218	TRP
1	A	220	LYS
1	A	242	SER
1	A	265	TYR
1	A	280	HIS
1	A	289	SER
1	A	317	ILE
1	A	329	ASP
1	A	359	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	362	GLU
1	A	366	GLU
1	A	368	THR
1	A	383	GLU
1	A	432	GLU
1	A	452	SER
1	A	460[A]	ASP
1	A	460[B]	ASP
1	A	464	HIS
1	A	466	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	189	GLN
1	A	193	ASN
1	A	199	HIS
1	A	228	HIS
1	A	298	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	1FS	A	501	-	31,33,33	3.71	12 (38%)	42,46,46	8.35	15 (35%)
4	GOL	A	507	-	5,5,5	0.49	0	5,5,5	0.25	0
3	SO4	A	505	-	4,4,4	0.60	0	6,6,6	0.43	0
3	SO4	A	502	-	4,4,4	0.53	0	6,6,6	0.52	0
4	GOL	A	510	-	5,5,5	0.67	0	5,5,5	1.12	0
4	GOL	A	509	-	5,5,5	0.70	0	5,5,5	1.08	0
3	SO4	A	504	-	4,4,4	0.57	0	6,6,6	0.46	0
3	SO4	A	503	-	4,4,4	0.50	0	6,6,6	0.82	0
4	GOL	A	508	-	5,5,5	0.27	0	5,5,5	0.60	0
3	SO4	A	506	-	4,4,4	0.17	0	6,6,6	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1FS	A	501	-	-	5/16/20/20	0/3/3/3
4	GOL	A	507	-	-	2/4/4/4	-
4	GOL	A	509	-	-	2/4/4/4	-
4	GOL	A	510	-	-	4/4/4/4	-
4	GOL	A	508	-	-	4/4/4/4	-

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	1FS	O5-N3	9.29	1.38	1.22
2	A	501	1FS	C1-N1	8.35	1.34	1.14
2	A	501	1FS	C20-N4	8.12	1.33	1.14
2	A	501	1FS	C2-C1	7.27	1.55	1.44
2	A	501	1FS	O2-N2	5.88	1.32	1.22
2	A	501	1FS	C19-C20	4.96	1.51	1.44
2	A	501	1FS	C12-C11	4.13	1.45	1.38
2	A	501	1FS	C16-C17	-3.96	1.31	1.38
2	A	501	1FS	C4-N2	-3.86	1.35	1.45
2	A	501	1FS	C17-N3	-3.48	1.36	1.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	1FS	C5-C4	3.34	1.45	1.38
2	A	501	1FS	C3-C4	-2.34	1.34	1.39

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	1FS	C2-C1-N1	-37.41	116.27	177.88
2	A	501	1FS	C19-C20-N4	-33.70	122.39	177.88
2	A	501	1FS	C8-O3-C7	-9.51	95.39	118.04
2	A	501	1FS	C12-C11-CL	8.70	130.02	119.15
2	A	501	1FS	C3-C4-N2	-5.93	113.52	118.75
2	A	501	1FS	C14-C19-C20	5.90	125.32	119.57
2	A	501	1FS	C10-C11-CL	-5.41	110.90	119.35
2	A	501	1FS	C13-O4-C14	-5.24	105.57	118.04
2	A	501	1FS	C5-C4-C3	3.68	124.76	120.09
2	A	501	1FS	O4-C14-C15	-3.13	112.37	120.73
2	A	501	1FS	C5-C4-N2	3.09	121.70	119.38
2	A	501	1FS	O4-C14-C19	2.83	127.41	116.89
2	A	501	1FS	C6-C5-C4	-2.75	116.25	120.08
2	A	501	1FS	C16-C17-C18	2.64	123.44	120.09
2	A	501	1FS	C3-C2-C1	2.41	123.36	119.74

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	1FS	N1-C1-C2-C7
2	A	501	1FS	C3-C4-N2-O2
2	A	501	1FS	C5-C4-N2-O2
2	A	501	1FS	C16-C17-N3-O5
2	A	501	1FS	C18-C17-N3-O5
4	A	510	GOL	O1-C1-C2-C3
4	A	510	GOL	C1-C2-C3-O3
4	A	509	GOL	C1-C2-C3-O3
4	A	508	GOL	C1-C2-C3-O3
4	A	507	GOL	O1-C1-C2-C3
4	A	508	GOL	O1-C1-C2-C3
4	A	510	GOL	O1-C1-C2-O2
4	A	510	GOL	O2-C2-C3-O3
4	A	509	GOL	O2-C2-C3-O3
4	A	508	GOL	O2-C2-C3-O3
4	A	507	GOL	O1-C1-C2-O2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	A	508	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	1FS	12	0
3	A	502	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	466/466 (100%)	0.07	0 100 100	29, 53, 78, 111	1 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	A	506	5/5	0.73	0.25	85,94,112,126	0
2	1FS	A	501	31/31	0.74	0.37	61,99,123,127	31
4	GOL	A	509	6/6	0.76	0.21	54,67,73,87	0
3	SO4	A	505	5/5	0.88	0.27	72,80,91,99	0
4	GOL	A	510	6/6	0.91	0.19	48,60,64,65	0
3	SO4	A	502	5/5	0.93	0.13	63,64,80,96	0
4	GOL	A	508	6/6	0.94	0.35	54,63,71,71	0
3	SO4	A	504	5/5	0.94	0.12	73,78,84,91	0
4	GOL	A	507	6/6	0.95	0.20	57,62,63,65	0
3	SO4	A	503	5/5	0.97	0.15	58,66,75,78	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.