



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 3, 2024 – 05:56 AM EST

PDB ID : 1Y4J
Title : Crystal structure of the paralogue of the human formylglycine generating enzyme
Authors : Dickmanns, A.; Rudolph, M.G.; Ficner, R.
Deposited on : 2004-12-01
Resolution : 1.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3097 (1.86-1.86)
Clashscore	180529	3359 (1.86-1.86)
Ramachandran outliers	177936	3335 (1.86-1.86)
Sidechain outliers	177891	3335 (1.86-1.86)
RSRZ outliers	164620	3097 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	284	 86% 7% 6%
1	B	284	 88% 6% 6%
2	C	3	 67% 33%
3	D	2	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	D	1	X	-	-	-

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 4982 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Sulfatase modifying factor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	266	2173	1374	400	389	10	0	7	0
1	B	268	2169	1372	394	395	8	0	5	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	302	ARG	-	expression tag	UNP Q8NBJ7
A	303	GLY	-	expression tag	UNP Q8NBJ7
A	304	SER	-	expression tag	UNP Q8NBJ7
A	305	HIS	-	expression tag	UNP Q8NBJ7
A	306	HIS	-	expression tag	UNP Q8NBJ7
A	307	HIS	-	expression tag	UNP Q8NBJ7
A	308	HIS	-	expression tag	UNP Q8NBJ7
A	309	HIS	-	expression tag	UNP Q8NBJ7
A	310	HIS	-	expression tag	UNP Q8NBJ7
B	302	ARG	-	expression tag	UNP Q8NBJ7
B	303	GLY	-	expression tag	UNP Q8NBJ7
B	304	SER	-	expression tag	UNP Q8NBJ7
B	305	HIS	-	expression tag	UNP Q8NBJ7
B	306	HIS	-	expression tag	UNP Q8NBJ7
B	307	HIS	-	expression tag	UNP Q8NBJ7
B	308	HIS	-	expression tag	UNP Q8NBJ7
B	309	HIS	-	expression tag	UNP Q8NBJ7
B	310	HIS	-	expression tag	UNP Q8NBJ7

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	3	38	22	2	14	0	0	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	D	2	28	16	2	10	0	0	0

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Ca	0	0
			2	2		
4	B	2	Total	Ca	0	0
			2	2		

- Molecule 5 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O 8 6 2	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	286	Total O 286 286	0	0
6	B	276	Total O 276 276	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	47.08Å 100.75Å 62.70Å 90.00° 101.68° 90.00°	Depositor
Resolution (Å)	32.00 – 1.86 32.00 – 1.86	Depositor EDS
% Data completeness (in resolution range)	99.3 (32.00-1.86) 99.3 (32.00-1.86)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.87 (at 1.87Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.170 , 0.205 0.177 , 0.209	Depositor DCC
R_{free} test set	2414 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	28.4	Xtrriage
Anisotropy	0.197	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 43.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4982	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, NAG, MPD, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.60	0/2274	0.70	0/3080
1	B	0.59	0/2255	0.70	1/3061 (0.0%)
All	All	0.59	0/4529	0.70	1/6141 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	253	ARG	NE-CZ-NH2	-5.79	117.41	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2173	0	2053	14	0
1	B	2169	0	2035	10	0
2	C	38	0	34	1	0
3	D	28	0	25	0	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
5	B	8	0	14	1	0
6	A	286	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	276	0	0	5	0
All	All	4982	0	4161	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:84[B]:ARG:NH1	6:B:2146:HOH:O	1.89	1.04
1:B:32[A]:GLN:HE21	1:B:294:ALA:HB1	1.42	0.81
1:B:32[A]:GLN:NE2	1:B:294:ALA:HB1	1.99	0.77
1:B:127:ARG:HG2	6:B:2187:HOH:O	1.93	0.68
1:A:28:THR:HG23	1:A:30:MET:H	1.58	0.67
6:A:1226:HOH:O	2:C:3:FUC:O2	2.13	0.66
1:B:29:SER:OG	6:B:2152:HOH:O	2.15	0.62
1:A:55:ARG:NH2	6:A:1105:HOH:O	2.34	0.61
1:B:46:ASP:HB3	1:B:270:ARG:HH22	1.68	0.59
1:A:125:PHE:CE1	1:A:128:GLN:HG3	2.39	0.57
1:A:49:ASP:OD1	1:A:242:TYR:OH	2.16	0.56
1:A:86:GLU:HA	1:A:89[A]:MET:CE	2.38	0.54
5:B:2001:MPD:H11	6:B:2212:HOH:O	2.10	0.51
1:A:173:ALA:HA	1:A:230:LEU:HD13	1.95	0.49
1:A:72:LYS:HG2	6:A:1280:HOH:O	2.14	0.48
1:B:244:ALA:O	1:B:247:GLN:HG2	2.14	0.47
1:A:86:GLU:HA	1:A:89[A]:MET:HE3	1.97	0.46
1:B:37:ARG:HG2	1:B:58:THR:OG1	2.16	0.45
1:A:175:GLY:HA2	1:A:225:TYR:CD2	2.53	0.44
1:A:105:ARG:HG2	1:A:105:ARG:HH11	1.84	0.43
1:A:237:TRP:CZ2	1:A:286:LEU:HD21	2.54	0.43
1:A:86:GLU:HA	1:A:89[A]:MET:HE2	2.00	0.42
1:B:187:TRP:CD1	6:B:2195:HOH:O	2.57	0.41
1:A:277:ASN:ND2	1:A:281:SER:HB2	2.36	0.41
1:A:86:GLU:HA	1:A:89[B]:MET:CE	2.50	0.41
1:B:44:SER:OG	1:B:46:ASP:HB2	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	269/284 (95%)	260 (97%)	8 (3%)	1 (0%)	30	18
1	B	270/284 (95%)	259 (96%)	10 (4%)	1 (0%)	30	18
All	All	539/568 (95%)	519 (96%)	18 (3%)	2 (0%)	30	18

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	233	ASN
1	B	233	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/231 (96%)	218 (98%)	5 (2%)	47	32
1	B	221/231 (96%)	219 (99%)	2 (1%)	75	70
All	All	444/462 (96%)	437 (98%)	7 (2%)	62	46

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	48	ARG
1	A	55	ARG
1	A	79[A]	ARG
1	A	79[B]	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	105	ARG
1	B	137	ARG
1	B	277	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	142	HIS
1	A	243	GLN
1	A	247	GLN
1	B	32[A]	GLN
1	B	243	GLN
1	B	277	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	1,2	14,14,15	0.52	0	17,19,21	1.84	5 (29%)
2	NAG	C	2	2	14,14,15	0.32	0	17,19,21	0.97	1 (5%)
2	FUC	C	3	2	10,10,11	0.40	0	14,14,16	0.92	1 (7%)
3	NAG	D	1	3,1	14,14,15	0.52	0	17,19,21	1.91	4 (23%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	D	2	3	14,14,15	0.63	0	17,19,21	1.07	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
2	FUC	C	3	2	-	-	0/1/1/1
3	NAG	D	1	3,1	1/1/5/7	2/6/23/26	0/1/1/1
3	NAG	D	2	3	-	5/6/23/26	0/1/1/1

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	1	NAG	C1-O5-C5	5.67	119.79	112.19
2	C	1	NAG	C1-C2-N2	-4.68	103.06	110.43
2	C	1	NAG	O5-C1-C2	-3.08	106.52	111.29
3	D	2	NAG	C4-C3-C2	2.94	115.32	111.02
3	D	1	NAG	O5-C1-C2	2.89	115.77	111.29
2	C	3	FUC	O5-C1-C2	2.71	117.26	110.79
3	D	1	NAG	C4-C3-C2	-2.55	107.29	111.02
2	C	1	NAG	O5-C5-C6	-2.47	102.86	107.66
2	C	2	NAG	C1-C2-N2	-2.46	106.56	110.43
3	D	1	NAG	O4-C4-C5	2.45	115.37	109.32
2	C	1	NAG	C2-N2-C7	-2.40	119.68	122.90
2	C	1	NAG	C1-O5-C5	2.11	115.02	112.19

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	D	1	NAG	C1

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	2	NAG	O5-C5-C6-O6

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	C	1	NAG	O5-C5-C6-O6
3	D	2	NAG	C4-C5-C6-O6
2	C	2	NAG	C4-C5-C6-O6
3	D	2	NAG	C8-C7-N2-C2
3	D	2	NAG	O7-C7-N2-C2
2	C	2	NAG	O5-C5-C6-O6
2	C	1	NAG	C4-C5-C6-O6
3	D	1	NAG	C8-C7-N2-C2
3	D	2	NAG	C3-C2-N2-C7
3	D	1	NAG	O7-C7-N2-C2


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	3	FUC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	MPD	B	2001	-	7,7,7	0.36	0	9,10,10	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MPD	B	2001	-	-	0/5/5/5	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	2001	MPD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	266/284 (93%)	-0.18	4 (1%) 71 75	17, 29, 45, 63	7 (2%)
1	B	268/284 (94%)	-0.19	4 (1%) 71 75	14, 30, 47, 53	4 (1%)
All	All	534/568 (94%)	-0.19	8 (1%) 71 75	14, 29, 47, 63	11 (2%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	27	ALA	4.1
1	A	108	ALA	3.2
1	A	187	TRP	2.8
1	A	106	ASN	2.6
1	B	187	TRP	2.5
1	B	109	THR	2.5
1	A	28	THR	2.4
1	B	28	THR	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	D	2	14/15	0.67	0.15	78,80,80,80	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NAG	D	1	14/15	0.71	0.14	63,67,71,75	0
2	NAG	C	1	14/15	0.76	0.14	45,55,59,60	0
2	NAG	C	2	14/15	0.82	0.11	58,59,61,62	0
2	FUC	C	3	10/11	0.94	0.08	36,37,41,42	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MPD	B	2001	8/8	0.74	0.23	61,61,62,62	0
4	CA	B	1003	1/1	0.98	0.08	31,31,31,31	0
4	CA	B	1002	1/1	0.99	0.03	27,27,27,27	0
4	CA	A	1001	1/1	0.99	0.02	23,23,23,23	0
4	CA	A	1004	1/1	0.99	0.06	30,30,30,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.