



# Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 12:04 am BST

PDB ID : 1Y6V  
Title : Structure of E. coli Alkaline Phosphatase in presence of cobalt at 1.60 Å resolution  
Authors : Wang, J.; Stieglitz, K.; Kantrowitz, E.R.  
Deposited on : 2004-12-07  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	449	
1	B	449	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SO4	A	858	-	-	X	-

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7346 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Alkaline phosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	449	3304	2042	581	669	12	0	0	0
1	B	449	3304	2042	581	669	12	0	0	0

- Molecule 2 is COBALT (II) ION (three-letter code: CO) (formula: Co).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	3	Total	Co	0	0
			3	3		
2	A	3	Total	Co	0	0
			3	3		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O P 5 4 1	0	0
3	B	1	Total O P 5 4 1	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	408	Total O 408 408	0	0
5	B	304	Total O 304 304	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alkaline phosphatase



- Molecule 1: Alkaline phosphatase



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	76.47Å 164.26Å 192.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.60 49.20 – 1.59	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.00-1.60) 96.1 (49.20-1.59)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.86 (at 1.59Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.191 , 0.212 0.192 , 0.212	Depositor DCC
$R_{free}$ test set	15852 reflections (9.89%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.4	Xtrriage
Anisotropy	0.351	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 54.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7346	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.30% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, CO, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.93	1/3359 (0.0%)	0.96	5/4560 (0.1%)
1	B	0.86	2/3359 (0.1%)	0.93	6/4560 (0.1%)
All	All	0.90	3/6718 (0.0%)	0.94	11/9120 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	322	GLU	CD-OE1	11.47	1.38	1.25
1	B	322	GLU	CB-CG	7.66	1.66	1.52
1	A	103	ALA	CA-CB	5.03	1.63	1.52

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	267	ARG	NE-CZ-NH1	-7.25	116.67	120.30
1	B	293	ASN	N-CA-C	6.44	128.39	111.00
1	A	209	LYS	CD-CE-NZ	6.32	126.23	111.70
1	A	418	ARG	NE-CZ-NH1	-6.11	117.25	120.30
1	B	323	GLY	N-CA-C	-5.96	98.20	113.10
1	A	323	GLY	N-CA-C	-5.96	98.21	113.10
1	A	330	ASP	CB-CG-OD2	5.84	123.55	118.30
1	B	97	ASP	N-CA-C	-5.76	95.45	111.00
1	A	24	ARG	NE-CZ-NH1	-5.74	117.43	120.30
1	B	322	GLU	CG-CD-OE1	5.39	129.08	118.30
1	B	312	LYS	CD-CE-NZ	5.16	123.58	111.70

There are no chirality outliers.

There are no planarity outliers.



## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3304	0	3248	37	0
1	B	3304	0	3248	55	0
2	A	3	0	0	0	0
2	B	3	0	0	0	0
3	A	5	0	0	1	0
3	B	5	0	0	1	0
4	A	5	0	0	2	0
4	B	5	0	0	0	0
5	A	408	0	0	12	0
5	B	304	0	0	2	0
All	All	7346	0	6496	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:293:ASN:ND2	1:B:295:SER:H	1.64	0.93
1:A:279:ILE:HD13	1:A:382:LYS:HE3	1.47	0.93
1:B:293:ASN:HD22	1:B:295:SER:H	0.90	0.90
1:B:293:ASN:HD22	1:B:295:SER:N	1.75	0.80
1:A:292:ARG:O	1:A:293:ASN:HB2	1.83	0.78
1:A:312:LYS:HE2	5:A:1093:HOH:O	1.86	0.74
1:A:267:ARG:HA	1:A:292:ARG:HD2	1.70	0.74
1:A:176:GLU:HB2	5:A:1202:HOH:O	1.87	0.73
1:B:90:ASN:OD1	1:B:92:LYS:HG2	1.88	0.72
1:B:236:LEU:HD23	1:B:256:LEU:HB3	1.69	0.72
1:A:279:ILE:CD1	1:A:382:LYS:HE3	2.19	0.72
1:B:449:LYS:HB3	1:B:449:LYS:NZ	2.09	0.68
1:A:227:GLU:HA	1:A:230:GLN:HE21	1.59	0.66
1:B:176:GLU:HG3	1:B:177:LYS:HG3	1.78	0.65
4:A:858:SO4:O1	5:A:859:HOH:O	2.14	0.64
1:A:230:GLN:HG2	5:A:1098:HOH:O	1.96	0.64
1:B:8:GLU:CD	1:B:8:GLU:H	2.01	0.63

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:248:GLU:HA	1:B:312:LYS:NZ	2.14	0.63
1:B:247:THR:O	1:B:248:GLU:HG3	1.99	0.63
1:A:172:SER:O	1:A:176:GLU:HG2	1.99	0.62
1:B:214:THR:HG22	1:B:224:THR:HA	1.82	0.61
1:A:292:ARG:NH1	5:A:902:HOH:O	2.36	0.58
1:A:40:LYS:N	1:A:40:LYS:HD3	2.18	0.58
1:B:235:GLN:NE2	1:B:253:LYS:HG2	2.19	0.58
1:A:102:SER:OG	3:A:856:PO4:P	2.61	0.57
1:B:2:PRO:HG2	1:B:357:LYS:HG2	1.85	0.57
1:B:293:ASN:ND2	1:B:295:SER:OG	2.38	0.57
1:B:247:THR:C	1:B:248:GLU:HG3	2.26	0.56
1:A:357:LYS:HE3	5:A:1088:HOH:O	2.05	0.55
1:A:188:LYS:H	1:A:188:LYS:CE	2.19	0.55
1:A:188:LYS:HE2	1:A:188:LYS:H	1.72	0.54
1:B:293:ASN:ND2	1:B:294:ASP:H	2.04	0.54
1:B:90:ASN:HD21	1:B:92:LYS:HE3	1.72	0.54
1:A:1:THR:N	1:A:358:LYS:O	2.41	0.54
1:B:248:GLU:HA	1:B:312:LYS:HZ2	1.73	0.54
1:B:195:LEU:HD23	1:B:195:LEU:C	2.29	0.53
1:A:379:PRO:HA	1:A:399:VAL:HG21	1.91	0.53
1:B:201:ASP:OD2	1:B:251:GLN:NE2	2.42	0.53
1:A:314:GLU:HA	1:A:314:GLU:OE2	2.09	0.53
1:A:279:ILE:HD13	1:A:382:LYS:CE	2.30	0.52
1:A:292:ARG:O	1:A:293:ASN:CB	2.56	0.52
1:B:293:ASN:ND2	1:B:295:SER:N	2.47	0.52
1:B:214:THR:HG22	1:B:224:THR:CA	2.39	0.51
1:B:449:LYS:HB3	1:B:449:LYS:HZ2	1.75	0.51
1:B:214:THR:HG22	1:B:224:THR:N	2.26	0.51
1:B:449:LYS:CB	1:B:449:LYS:NZ	2.74	0.50
1:B:195:LEU:HD23	1:B:195:LEU:O	2.11	0.50
1:B:234:TYR:CD1	1:B:254:PRO:HG2	2.47	0.50
1:B:309:LEU:HA	1:B:312:LYS:HE2	1.93	0.49
1:B:358:LYS:HD3	1:B:359:GLU:N	2.27	0.49
1:B:3:GLU:HA	1:B:3:GLU:OE2	2.13	0.49
1:B:229:ALA:O	1:B:234:TYR:HB2	2.13	0.49
1:B:102:SER:OG	3:B:956:PO4:P	2.71	0.49
1:B:449:LYS:HB3	1:B:449:LYS:HZ3	1.78	0.48
1:A:188:LYS:H	1:A:188:LYS:NZ	2.11	0.48
1:A:253:LYS:HE2	5:A:1183:HOH:O	2.14	0.47
1:B:15:ASP:O	1:B:21:GLY:HA3	2.15	0.47
1:A:1:THR:O	1:A:1:THR:HG23	2.15	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:137:LYS:HE2	1:A:198:ALA:O	2.15	0.46
1:B:304:ASP:OD2	1:B:351:ARG:NH1	2.48	0.46
1:A:188:LYS:HE2	1:A:188:LYS:HB2	1.55	0.45
4:A:858:SO4:O2	5:A:902:HOH:O	2.21	0.44
1:B:287:THR:CG2	5:B:1142:HOH:O	2.65	0.44
1:B:314:GLU:HA	1:B:314:GLU:OE1	2.18	0.44
1:A:267:ARG:HG2	1:A:268:TRP:CD1	2.53	0.44
1:B:116:TYR:CZ	1:B:119:ALA:HB2	2.53	0.44
1:B:137:LYS:HE3	1:B:251:GLN:OE1	2.17	0.44
1:B:214:THR:HG22	1:B:223:LYS:C	2.39	0.43
1:A:248:GLU:HB2	5:A:1183:HOH:O	2.18	0.43
1:A:247:THR:O	1:A:312:LYS:NZ	2.51	0.43
1:B:293:ASN:ND2	1:B:294:ASP:N	2.66	0.43
1:A:33:LEU:HD12	1:A:33:LEU:HA	1.90	0.42
1:A:365:ILE:HD13	1:A:438:LEU:HD11	2.00	0.42
1:B:225:LEU:HD12	1:B:225:LEU:HA	1.82	0.42
1:B:424:PRO:O	1:B:425:HIS:HB2	2.19	0.42
1:B:208:ALA:HB2	1:B:258:LEU:HB3	2.00	0.42
1:B:298:THR:OG1	1:B:301:GLN:HG3	2.19	0.42
1:B:33:LEU:HA	1:B:33:LEU:HD12	1.83	0.42
1:A:73:LYS:HD3	5:A:1237:HOH:O	2.20	0.42
1:B:199:ARG:HA	1:B:234:TYR:OH	2.19	0.42
1:A:288:PRO:HB3	1:A:292:ARG:NH1	2.35	0.42
1:B:312:LYS:HE3	1:B:312:LYS:HB2	1.76	0.42
1:B:407:GLU:H	1:B:407:GLU:HG2	1.70	0.41
1:B:155:THR:HB	1:B:322:GLU:OE1	2.20	0.41
1:B:40:LYS:HE3	5:B:1099:HOH:O	2.21	0.41
1:A:292:ARG:CZ	5:A:902:HOH:O	2.67	0.41
1:A:196:LEU:HD23	1:A:196:LEU:HA	1.90	0.41
1:A:293:ASN:HB3	1:A:295:SER:H	1.85	0.41
1:B:213:GLU:O	1:B:224:THR:HA	2.20	0.41
1:B:351:ARG:HA	1:B:351:ARG:HD2	1.68	0.41
1:A:181:ASN:HD22	1:A:188:LYS:NZ	2.19	0.41
1:A:6:VAL:HG12	5:A:967:HOH:O	2.19	0.41
1:B:155:THR:HB	1:B:322:GLU:CD	2.41	0.41
1:B:1:THR:OG1	1:B:2:PRO:HD2	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	447/449 (100%)	437 (98%)	9 (2%)	1 (0%)	47	26
1	B	447/449 (100%)	434 (97%)	13 (3%)	0	100	100
All	All	894/898 (100%)	871 (97%)	22 (2%)	1 (0%)	51	29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	292	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	340/340 (100%)	332 (98%)	8 (2%)	49	24
1	B	340/340 (100%)	321 (94%)	19 (6%)	21	5
All	All	680/680 (100%)	653 (96%)	27 (4%)	31	10

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	GLU
1	A	33	LEU
1	A	40	LYS
1	A	84	TYR
1	A	188	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	226	ARG
1	A	236	LEU
1	A	382	LYS
1	B	4	MET
1	B	8	GLU
1	B	33	LEU
1	B	84	TYR
1	B	92	LYS
1	B	171	PRO
1	B	172	SER
1	B	185	LYS
1	B	225	LEU
1	B	239	ASP
1	B	247	THR
1	B	261	ASP
1	B	287	THR
1	B	293	ASN
1	B	304	ASP
1	B	314	GLU
1	B	351	ARG
1	B	358	LYS
1	B	407	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	230	GLN
1	A	252	GLN
1	B	83	GLN
1	B	235	GLN
1	B	293	ASN
1	B	391	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 6 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	PO4	B	956	2	4,4,4	0.74	0	6,6,6	0.55	0
4	SO4	A	858	1	4,4,4	0.59	0	6,6,6	0.22	0
4	SO4	B	958	-	4,4,4	0.61	0	6,6,6	0.18	0
3	PO4	A	856	2	4,4,4	1.02	0	6,6,6	1.00	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	956	PO4	1	0
4	A	858	SO4	2	0
3	A	856	PO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	449/449 (100%)	-0.24	11 (2%) 59 56	11, 18, 35, 119	0
1	B	449/449 (100%)	0.35	37 (8%) 11 10	13, 25, 51, 144	0
All	All	898/898 (100%)	0.06	48 (5%) 26 24	11, 21, 48, 144	0

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	THR	17.9
1	A	1	THR	14.3
1	B	2	PRO	11.8
1	A	2	PRO	11.7
1	B	187	GLY	4.4
1	B	222	GLY	4.4
1	B	231	ALA	4.2
1	B	216	THR	3.7
1	A	408	ASP	3.5
1	B	234	TYR	3.4
1	B	449	LYS	3.4
1	B	224	THR	3.4
1	B	185	LYS	3.2
1	B	244	ASN	3.1
1	A	244	ASN	3.0
1	A	3	GLU	2.9
1	A	314	GLU	2.9
1	B	225	LEU	2.9
1	A	292	ARG	2.9
1	B	214	THR	2.8
1	B	182	ALA	2.8
1	B	245	SER	2.7
1	B	230	GLN	2.7
1	B	236	LEU	2.7

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	179	PRO	2.6
1	B	223	LYS	2.6
1	B	189	GLY	2.5
1	B	212	ALA	2.5
1	A	407	GLU	2.4
1	B	186	GLY	2.4
1	B	218	GLY	2.4
1	A	449	LYS	2.4
1	B	408	ASP	2.3
1	B	221	GLN	2.3
1	B	220	TRP	2.3
1	B	314	GLU	2.3
1	B	48	LEU	2.2
1	B	176	GLU	2.2
1	B	4	MET	2.2
1	B	312	LYS	2.2
1	A	248	GLU	2.2
1	B	241	ALA	2.2
1	B	3	GLU	2.1
1	A	312	LYS	2.1
1	B	227	GLU	2.0
1	B	170	GLY	2.0
1	B	233	GLY	2.0
1	B	253	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	PO4	B	956	5/5	0.90	0.12	27,28,31,31	0
3	PO4	A	856	5/5	0.92	0.10	17,18,20,24	0
4	SO4	B	958	5/5	0.93	0.12	26,27,27,28	5
4	SO4	A	858	5/5	0.94	0.13	26,27,29,30	5
2	CO	B	952	1/1	0.98	0.04	19,19,19,19	0
2	CO	B	950	1/1	0.99	0.04	19,19,19,19	0
2	CO	A	852	1/1	1.00	0.04	14,14,14,14	0
2	CO	A	851	1/1	1.00	0.02	12,12,12,12	0
2	CO	A	850	1/1	1.00	0.04	11,11,11,11	0
2	CO	B	951	1/1	1.00	0.04	19,19,19,19	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.