



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 26, 2021 – 06:02 pm BST

PDB ID : 6Y6R
Title : Crystal structure of MINDY1 T335D mutant
Authors : Abdul Rehman, S.A.; Kulathu, Y.
Deposited on : 2020-02-27
Resolution : 3.32 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1089 (3.36-3.28)
Clashscore	141614	1137 (3.36-3.28)
Ramachandran outliers	138981	1115 (3.36-3.28)
Sidechain outliers	138945	1114 (3.36-3.28)
RSRZ outliers	127900	1059 (3.36-3.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	289	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1993 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquitin carboxyl-terminal hydrolase MINDY-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	257	1993	1274	322	385	12	0	1	0


There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	96	GLY	-	expression tag	UNP Q8N5J2
A	97	PRO	-	expression tag	UNP Q8N5J2
A	98	LEU	-	expression tag	UNP Q8N5J2
A	99	GLY	-	expression tag	UNP Q8N5J2
A	100	SER	-	expression tag	UNP Q8N5J2
A	101	PRO	-	expression tag	UNP Q8N5J2
A	102	GLU	-	expression tag	UNP Q8N5J2
A	103	PHE	-	expression tag	UNP Q8N5J2
A	104	PRO	-	expression tag	UNP Q8N5J2
A	105	GLY	-	expression tag	UNP Q8N5J2
A	106	ARG	-	expression tag	UNP Q8N5J2
A	107	LEU	-	expression tag	UNP Q8N5J2
A	108	GLU	-	expression tag	UNP Q8N5J2
A	109	MET	-	expression tag	UNP Q8N5J2
A	335	ASP	THR	engineered mutation	UNP Q8N5J2

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ubiquitin carboxyl-terminal hydrolase MINDY-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	99.81Å 99.81Å 164.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	85.21 – 3.32 85.28 – 3.32	Depositor EDS
% Data completeness (in resolution range)	99.1 (85.21-3.32) 99.2 (85.28-3.32)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.39 (at 3.33Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.200 , 0.243 0.219 , 0.256	Depositor DCC
R_{free} test set	599 reflections (4.68%)	wwPDB-VP
Wilson B-factor (Å ²)	142.8	Xtriage
Anisotropy	0.504	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	1993	wwPDB-VP
Average B, all atoms (Å ²)	168.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.77	0/2046	0.94	0/2798

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1993	0	1878	38	0
All	All	1993	0	1878	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:186:LEU:HB3	1:A:189:ASN:HB2	1.37	1.02
1:A:167:MET:HE2	1:A:205:ALA:HB2	1.55	0.88
1:A:186:LEU:HB3	1:A:189:ASN:CB	2.13	0.79
1:A:138:PRO:HB3	1:A:320:PHE:CG	2.23	0.72
1:A:138:PRO:HB3	1:A:320:PHE:CD2	2.23	0.72

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:334:VAL:HG23	1:A:346:VAL:O	1.90	0.71
1:A:186:LEU:HD22	1:A:189:ASN:HD22	1.62	0.64
1:A:179:PRO:HG3	1:A:190:PHE:CE2	2.35	0.61
1:A:122:LYS:CB	1:A:150:TRP:CH2	2.84	0.60
1:A:200:VAL:HG11	1:A:224:PRO:HG2	1.82	0.60
1:A:334:VAL:O	1:A:334:VAL:HG12	2.02	0.60
1:A:201:LEU:HB3	1:A:202:PRO:HD3	1.86	0.57
1:A:137:CYS:HB2	1:A:204:LEU:O	2.04	0.57
1:A:344:GLN:O	1:A:361:ASP:HA	2.05	0.56
1:A:122:LYS:CB	1:A:150:TRP:HH2	2.17	0.56
1:A:240:TRP:CZ3	1:A:293:GLN:HG2	2.40	0.56
1:A:167:MET:HE2	1:A:205:ALA:CB	2.34	0.53
1:A:292:ALA:HB1	1:A:316:ARG:HH21	1.74	0.53
1:A:130:THR:HA	1:A:335:ASP:O	2.09	0.52
1:A:331:TYR:HB3	1:A:347:TRP:HB3	1.91	0.51
1:A:259:ASN:O	1:A:262:VAL:HG22	2.10	0.51
1:A:342:GLU:HB3	1:A:345:VAL:HG23	1.92	0.51
1:A:278:THR:O	1:A:282:ILE:HG12	2.12	0.50
1:A:143:MET:HE3	1:A:170:LEU:HB2	1.94	0.50
1:A:240:TRP:CH2	1:A:293:GLN:HG2	2.48	0.49
1:A:123:GLY:O	1:A:124:GLU:HG2	2.13	0.48
1:A:279:GLU:HA	1:A:282:ILE:HB	1.96	0.47
1:A:167:MET:O	1:A:170:LEU:HB3	2.16	0.46
1:A:143:MET:CE	1:A:146:LEU:HD12	2.45	0.46
1:A:360:CYS:HA	1:A:365:HIS:O	2.16	0.46
1:A:237:TYR:HE2	1:A:312:SER:CB	2.30	0.45
1:A:342:GLU:HG2	1:A:344:GLN:HE22	1.80	0.45
1:A:240:TRP:CE3	1:A:293:GLN:HG2	2.52	0.44
1:A:217:VAL:O	1:A:237:TYR:HA	2.19	0.43
1:A:115:CYS:HB3	1:A:340:LEU:HD21	2.01	0.42
1:A:142:ILE:O	1:A:145:ILE:HB	2.19	0.42
1:A:325:LYS:HA	1:A:329:HIS:O	2.20	0.42
1:A:286:PHE:O	1:A:290:THR:HG23	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	254/289 (88%)	227 (89%)	27 (11%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	220/253 (87%)	198 (90%)	22 (10%)	7 28

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	112	ASP
1	A	128	ILE
1	A	131	GLN
1	A	157	GLN
1	A	173	CYS
1	A	176	SER
1	A	187	GLN
1	A	204	LEU
1	A	219	ASP
1	A	223	THR
1	A	246	SER
1	A	248	GLU
1	A	256	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	264	ARG
1	A	265	ILE
1	A	269	LYS
1	A	272	SER
1	A	312	SER
1	A	316	ARG
1	A	340	LEU
1	A	362	SER
1	A	366	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	157	GLN
1	A	189	ASN
1	A	245	GLN
1	A	259	ASN
1	A	275	ASN
1	A	297	HIS
1	A	344	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	257/289 (88%)	1.04	55 (21%) 0 0	120, 165, 223, 291	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	185	GLY	4.2
1	A	347	TRP	3.5
1	A	322	THR	3.5
1	A	240	TRP	3.5
1	A	302	LEU	3.5
1	A	139	LEU	3.3
1	A	204	LEU	3.3
1	A	287	LEU	3.1
1	A	166	LEU	3.1
1	A	241	LEU	3.1
1	A	229	PHE	3.0
1	A	333	LEU	2.9
1	A	242	VAL	2.8
1	A	320	PHE	2.8
1	A	142	ILE	2.8
1	A	350	LEU	2.7
1	A	345	VAL	2.7
1	A	359	PHE	2.7
1	A	214	PHE	2.7
1	A	236	LEU	2.7
1	A	315	PHE	2.7
1	A	143	MET	2.7
1	A	294	LEU	2.6
1	A	231	LEU	2.6
1	A	261	LEU	2.6
1	A	119	ILE	2.6
1	A	311	LEU	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	283	ALA	2.5
1	A	217	VAL	2.5
1	A	160	VAL	2.4
1	A	174	LEU	2.4
1	A	346	VAL	2.4
1	A	368	HIS	2.4
1	A	146	LEU	2.4
1	A	147	PHE	2.4
1	A	306	ALA	2.4
1	A	331	TYR	2.3
1	A	234	ILE	2.3
1	A	128	ILE	2.3
1	A	364	PHE	2.3
1	A	253	VAL	2.3
1	A	141	ALA	2.2
1	A	232	LEU	2.2
1	A	314	PHE	2.2
1	A	228	VAL	2.2
1	A	205	ALA	2.2
1	A	129	ILE	2.1
1	A	332	LEU	2.1
1	A	313	VAL	2.1
1	A	265	ILE	2.1
1	A	171	GLY	2.1
1	A	203	LYS	2.1
1	A	282	ILE	2.1
1	A	208	LEU	2.0
1	A	138	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.