



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 7, 2024 – 04:17 PM JST

PDB ID : 8Y6B  
Title : Structure of human LGI1-ADAM22 complex in space group P212121  
Authors : Liu, H.; Xu, F.  
Deposited on : 2024-02-02  
Resolution : 3.49 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1659 (3.60-3.40)
Clashscore	141614	1036 (3.58-3.42)
Ramachandran outliers	138981	1005 (3.58-3.42)
Sidechain outliers	138945	1006 (3.58-3.42)
RSRZ outliers	127900	1559 (3.60-3.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	486	 13% 83% 17%
1	B	486	 84% 15%
1	C	486	 2% 84% 16%
2	D	526	 3% 72% 24%
2	E	526	 2% 76% 21%
2	F	526	 74% 23%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

<b>Mol</b>	<b>Type</b>	<b>Chain</b>	<b>Res</b>	<b>Chirality</b>	<b>Geometry</b>	<b>Clashes</b>	<b>Electron density</b>
3	NAG	A	801	-	-	-	X
3	NAG	A	803	-	-	-	X
3	NAG	D	602	-	-	-	X
3	NAG	E	601	-	-	-	X

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 24226 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Disintegrin and metalloproteinase domain-containing protein 22.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	486	3728	2303	643	731	51	0	0	0
1	B	486	3728	2303	643	731	51	0	0	0
1	C	486	3728	2303	643	731	51	0	0	0

- Molecule 2 is a protein called Leucine-rich glioma-inactivated protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	517	4194	2700	698	781	15	0	0	0
2	F	517	4194	2700	698	781	15	0	0	0
2	E	517	4194	2700	698	781	15	0	0	0

There are 21 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
D	32	HIS	-	expression tag	UNP O95970
D	33	HIS	-	expression tag	UNP O95970
D	34	HIS	-	expression tag	UNP O95970
D	35	HIS	-	expression tag	UNP O95970
D	36	HIS	-	expression tag	UNP O95970
D	37	HIS	-	expression tag	UNP O95970
D	38	HIS	-	expression tag	UNP O95970
F	32	HIS	-	expression tag	UNP O95970
F	33	HIS	-	expression tag	UNP O95970
F	34	HIS	-	expression tag	UNP O95970
F	35	HIS	-	expression tag	UNP O95970
F	36	HIS	-	expression tag	UNP O95970

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	37	HIS	-	expression tag	UNP O95970
F	38	HIS	-	expression tag	UNP O95970
E	32	HIS	-	expression tag	UNP O95970
E	33	HIS	-	expression tag	UNP O95970
E	34	HIS	-	expression tag	UNP O95970
E	35	HIS	-	expression tag	UNP O95970
E	36	HIS	-	expression tag	UNP O95970
E	37	HIS	-	expression tag	UNP O95970
E	38	HIS	-	expression tag	UNP O95970

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	D	1	Total	C	N	O	0	0
			14	8	1	5		
3	D	1	Total	C	N	O	0	0
			14	8	1	5		
3	D	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total C N O 14 8 1 5	0	0
3	B	1	Total C N O 14 8 1 5	0	0
3	B	1	Total C N O 14 8 1 5	0	0
3	C	1	Total C N O 14 8 1 5	0	0
3	C	1	Total C N O 14 8 1 5	0	0
3	C	1	Total C N O 14 8 1 5	0	0
3	F	1	Total C N O 14 8 1 5	0	0
3	F	1	Total C N O 14 8 1 5	0	0
3	F	1	Total C N O 14 8 1 5	0	0
3	E	1	Total C N O 14 8 1 5	0	0
3	E	1	Total C N O 14 8 1 5	0	0
3	E	1	Total C N O 14 8 1 5	0	0

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	3	Total Ca 3 3	0	0
4	D	1	Total Ca 1 1	0	0
4	B	3	Total Ca 3 3	0	0
4	C	3	Total Ca 3 3	0	0
4	F	1	Total Ca 1 1	0	0
4	E	1	Total Ca 1 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	2	Total O 2 2	0	0
5	D	13	Total O 13 13	0	0
5	B	88	Total O 88 88	0	0
5	C	54	Total O 54 54	0	0
5	F	13	Total O 13 13	0	0
5	E	26	Total O 26 26	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Disintegrin and metalloproteinase domain-containing protein 22




- Molecule 1: Disintegrin and metalloproteinase domain-containing protein 22




- Molecule 1: Disintegrin and metalloproteinase domain-containing protein 22









• Molecule 2: Leucine-rich glioma-inactivated protein 1




• Molecule 2: Leucine-rich glioma-inactivated protein 1



• Molecule 2: Leucine-rich glioma-inactivated protein 1

Chain E:  2% 76% 21% ..



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.45Å 196.08Å 420.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.46 – 3.49 42.46 – 3.49	Depositor EDS
% Data completeness (in resolution range)	96.8 (42.46-3.49) 96.8 (42.46-3.49)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.29 (at 3.48Å)	Xtrriage
Refinement program	PHENIX (1.10.1_2155)	Depositor
R, $R_{free}$	0.236 , 0.277 0.236 , 0.277	Depositor DCC
$R_{free}$ test set	2894 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	81.9	Xtrriage
Anisotropy	0.493	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.23 , 41.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	24226	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	115.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.97% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/3792	0.46	0/5105
1	B	0.31	0/3792	0.51	0/5105
1	C	0.29	0/3792	0.50	0/5105
2	D	0.28	0/4301	0.48	0/5836
2	E	0.27	0/4301	0.46	0/5836
2	F	0.25	0/4301	0.46	0/5836
All	All	0.28	0/24279	0.48	0/32823

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3728	0	3579	65	0
1	B	3728	0	3580	55	0
1	C	3728	0	3579	47	0
2	D	4194	0	4110	96	0
2	E	4194	0	4111	75	0
2	F	4194	0	4110	73	0
3	A	42	0	39	3	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	42	0	39	7	0
3	C	42	0	39	0	0
3	D	42	0	39	4	0
3	E	42	0	39	7	0
3	F	42	0	39	2	0
4	A	3	0	0	0	0
4	B	3	0	0	0	0
4	C	3	0	0	0	0
4	D	1	0	0	0	0
4	E	1	0	0	0	0
4	F	1	0	0	0	0
5	A	2	0	0	2	0
5	B	88	0	0	3	0
5	C	54	0	0	2	0
5	D	13	0	0	0	0
5	E	26	0	0	1	0
5	F	13	0	0	0	0
All	All	24226	0	23303	421	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 421 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:715:PHE:CD2	1:A:716:PRO:HD2	1.74	1.21
1:A:664:LEU:HD23	1:A:665:GLU:CG	1.84	1.07
1:A:664:LEU:CD2	1:A:665:GLU:HG2	1.90	1.02
1:A:664:LEU:HD23	1:A:665:GLU:HG2	1.05	1.01
2:D:477:MET:HG3	2:D:523:ARG:NH1	1.78	0.97

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	484/486 (100%)	469 (97%)	15 (3%)	0	100	100
1	B	484/486 (100%)	469 (97%)	15 (3%)	0	100	100
1	C	484/486 (100%)	475 (98%)	9 (2%)	0	100	100
2	D	515/526 (98%)	488 (95%)	26 (5%)	1 (0%)	47	81
2	E	515/526 (98%)	491 (95%)	24 (5%)	0	100	100
2	F	515/526 (98%)	497 (96%)	18 (4%)	0	100	100
All	All	2997/3036 (99%)	2889 (96%)	107 (4%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	522	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	420/420 (100%)	415 (99%)	5 (1%)	71	87
1	B	420/420 (100%)	407 (97%)	13 (3%)	40	70
1	C	420/420 (100%)	408 (97%)	12 (3%)	42	71
2	D	474/483 (98%)	456 (96%)	18 (4%)	33	65
2	E	474/483 (98%)	456 (96%)	18 (4%)	33	65
2	F	474/483 (98%)	452 (95%)	22 (5%)	27	61
All	All	2682/2709 (99%)	2594 (97%)	88 (3%)	38	68

5 of 88 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	245	LEU
2	E	51	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	F	274	ASN
2	F	421	THR
2	E	149	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
2	E	52	ASN
2	F	480	GLN
1	C	280	GLN
2	F	333	ASN
2	D	485	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 30 ligands modelled in this entry, 12 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	D	601	2	14,14,15	0.23	0	17,19,21	0.41	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	A	802	1	14,14,15	0.78	1 (7%)	17,19,21	1.27	2 (11%)
3	NAG	E	603	2	14,14,15	0.36	0	17,19,21	0.50	0
3	NAG	D	602	2	14,14,15	0.19	0	17,19,21	0.39	0
3	NAG	B	802	1	14,14,15	0.25	0	17,19,21	0.50	0
3	NAG	C	801	1	14,14,15	0.44	0	17,19,21	0.67	1 (5%)
3	NAG	A	801	1	14,14,15	0.21	0	17,19,21	0.41	0
3	NAG	B	801	1	14,14,15	0.47	0	17,19,21	1.30	1 (5%)
3	NAG	C	803	1	14,14,15	0.22	0	17,19,21	0.40	0
3	NAG	F	603	2	14,14,15	0.25	0	17,19,21	0.47	0
3	NAG	F	602	2	14,14,15	0.38	0	17,19,21	0.45	0
3	NAG	E	601	2	14,14,15	0.43	0	17,19,21	1.26	2 (11%)
3	NAG	A	803	1	14,14,15	0.81	1 (7%)	17,19,21	1.29	2 (11%)
3	NAG	B	803	1	14,14,15	0.45	0	17,19,21	0.45	0
3	NAG	F	601	2	14,14,15	0.68	1 (7%)	17,19,21	0.78	1 (5%)
3	NAG	D	603	2	14,14,15	0.21	0	17,19,21	0.43	0
3	NAG	E	602	2	14,14,15	0.51	0	17,19,21	1.29	2 (11%)
3	NAG	C	802	1	14,14,15	0.20	0	17,19,21	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	D	601	2	-	4/6/23/26	0/1/1/1
3	NAG	A	802	1	-	4/6/23/26	0/1/1/1
3	NAG	E	603	2	-	4/6/23/26	0/1/1/1
3	NAG	D	602	2	-	3/6/23/26	0/1/1/1
3	NAG	B	802	1	-	4/6/23/26	0/1/1/1
3	NAG	C	801	1	-	4/6/23/26	0/1/1/1
3	NAG	A	801	1	-	4/6/23/26	0/1/1/1
3	NAG	B	801	1	-	5/6/23/26	0/1/1/1
3	NAG	C	803	1	-	2/6/23/26	0/1/1/1
3	NAG	F	603	2	-	4/6/23/26	0/1/1/1
3	NAG	F	602	2	-	2/6/23/26	0/1/1/1
3	NAG	E	601	2	-	5/6/23/26	0/1/1/1
3	NAG	A	803	1	-	5/6/23/26	0/1/1/1
3	NAG	B	803	1	-	4/6/23/26	0/1/1/1

Continued on next page...



*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	F	601	2	-	4/6/23/26	0/1/1/1
3	NAG	D	603	2	-	3/6/23/26	0/1/1/1
3	NAG	E	602	2	-	3/6/23/26	0/1/1/1
3	NAG	C	802	1	-	3/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	803	NAG	C1-C2	2.76	1.56	1.52
3	A	802	NAG	C1-C2	2.59	1.56	1.52
3	F	601	NAG	C1-C2	2.18	1.55	1.52

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	802	NAG	C2-N2-C7	4.36	129.11	122.90
3	E	601	NAG	C2-N2-C7	4.36	129.11	122.90
3	B	801	NAG	C2-N2-C7	4.28	129.00	122.90
3	A	803	NAG	C2-N2-C7	4.28	129.00	122.90
3	E	602	NAG	C2-N2-C7	4.21	128.89	122.90

There are no chirality outliers.

5 of 67 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	801	NAG	O5-C5-C6-O6
3	A	803	NAG	C4-C5-C6-O6
3	C	801	NAG	C4-C5-C6-O6
3	E	601	NAG	C4-C5-C6-O6
3	A	801	NAG	C8-C7-N2-C2

There are no ring outliers.

9 monomers are involved in 23 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	601	NAG	4	0
3	A	802	NAG	2	0
3	E	603	NAG	4	0
3	B	801	NAG	1	0
3	E	601	NAG	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	803	NAG	1	0
3	B	803	NAG	6	0
3	F	601	NAG	2	0
3	E	602	NAG	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	486/486 (100%)	0.66	62 (12%) <b>3</b> <b>4</b>	102, 169, 251, 286	0
1	B	486/486 (100%)	-0.24	2 (0%) <b>92</b> <b>90</b>	35, 61, 118, 156	0
1	C	486/486 (100%)	-0.06	11 (2%) <b>60</b> <b>54</b>	48, 95, 174, 215	0
2	D	517/526 (98%)	0.26	14 (2%) <b>54</b> <b>48</b>	80, 130, 178, 202	0
2	E	517/526 (98%)	0.05	8 (1%) <b>73</b> <b>68</b>	60, 98, 165, 212	0
2	F	517/526 (98%)	-0.02	5 (0%) <b>82</b> <b>77</b>	42, 106, 177, 217	0
All	All	3009/3036 (99%)	0.11	102 (3%) <b>45</b> <b>40</b>	35, 112, 196, 286	0

The worst 5 of 102 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	576	GLU	7.5
1	A	455	GLY	5.7
1	A	637	GLY	5.6
1	A	716	PRO	5.1
1	A	718	ASN	4.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	A	801	14/15	0.23	0.59	224,252,264,266	0
3	NAG	A	803	14/15	0.49	0.67	222,240,246,247	0
3	NAG	E	601	14/15	0.56	0.62	200,211,220,221	0
3	NAG	D	602	14/15	0.57	0.44	187,209,214,217	0
3	NAG	A	802	14/15	0.58	0.29	178,183,189,189	0
3	NAG	C	802	14/15	0.72	0.27	168,184,191,192	0
3	NAG	F	601	14/15	0.76	0.20	167,193,208,210	0
4	CA	A	806	1/1	0.77	0.12	281,281,281,281	0
3	NAG	D	603	14/15	0.81	0.22	139,151,164,167	0
3	NAG	D	601	14/15	0.81	0.27	118,164,181,183	0
3	NAG	C	803	14/15	0.81	0.46	121,157,163,167	0
3	NAG	E	603	14/15	0.83	0.27	131,145,158,162	0
3	NAG	B	801	14/15	0.83	0.32	112,133,141,150	0
3	NAG	B	803	14/15	0.84	0.30	78,101,123,124	0
4	CA	A	805	1/1	0.85	0.06	181,181,181,181	0
4	CA	B	804	1/1	0.85	0.08	82,82,82,82	0
4	CA	C	806	1/1	0.86	0.11	129,129,129,129	0
3	NAG	F	603	14/15	0.87	0.25	122,130,135,140	0
3	NAG	C	801	14/15	0.88	0.31	101,118,137,142	0
3	NAG	E	602	14/15	0.89	0.27	104,135,157,161	0
3	NAG	B	802	14/15	0.90	0.15	60,84,112,134	0
3	NAG	F	602	14/15	0.90	0.37	96,107,123,132	0
4	CA	D	604	1/1	0.91	0.25	82,82,82,82	0
4	CA	B	806	1/1	0.94	0.12	34,34,34,34	0
4	CA	C	804	1/1	0.95	0.18	72,72,72,72	0
4	CA	A	804	1/1	0.96	0.16	169,169,169,169	0
4	CA	E	604	1/1	0.97	0.35	71,71,71,71	0
4	CA	F	604	1/1	0.98	0.44	54,54,54,54	0
4	CA	B	805	1/1	0.98	0.11	36,36,36,36	0
4	CA	C	805	1/1	0.99	0.18	44,44,44,44	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.