



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 12, 2024 – 07:08 AM EST

PDB ID : 1Y7H
Title : Structural and biochemical studies identify tobacco SABP2 as a methylsalicylate esterase and further implicate it in plant innate immunity, Northeast Structural Genomics Target AR2241
Authors : Forouhar, F.; Yang, Y.; Kumar, D.; Chen, Y.; Fridman, E.; Park, S.W.; Chiang, Y.; Acton, T.B.; Montelione, G.T.; Pichersky, E.; Klessig, D.F.; Tong, L.; Northeast Structural Genomics Consortium (NESG)
Deposited on : 2004-12-08
Resolution : 2.52 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	6935 (2.54-2.50)
Clashscore	180529	7778 (2.54-2.50)
Ramachandran outliers	177936	7674 (2.54-2.50)
Sidechain outliers	177891	7676 (2.54-2.50)
RSRZ outliers	164620	6935 (2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	268	 64% 31% . .
1	B	268	 57% 38% . .
1	C	268	 61% 35% .
1	D	268	 62% 33% . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	268	
1	F	268	
1	G	268	
1	H	268	

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 16579 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called salicylic acid-binding protein 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	257	2031	1310	330	377	5	9	0	0	0
1	B	259	2051	1323	333	381	5	9	0	0	0
1	C	257	2031	1310	330	377	5	9	0	0	0
1	D	259	2051	1323	333	381	5	9	0	0	0
1	E	258	2043	1319	331	379	5	9	0	0	0
1	F	258	2042	1317	331	380	5	9	0	0	0
1	G	257	2031	1310	330	377	5	9	0	0	0
1	H	259	2051	1323	333	381	5	9	0	0	0

There are 136 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	63	MSE	MET	modified residue	UNP Q6RYA0
A	66	MSE	MET	modified residue	UNP Q6RYA0
A	85	MSE	MET	modified residue	UNP Q6RYA0
A	91	MSE	MET	modified residue	UNP Q6RYA0
A	108	MSE	MET	modified residue	UNP Q6RYA0
A	149	MSE	MET	modified residue	UNP Q6RYA0
A	183	MSE	MET	modified residue	UNP Q6RYA0
A	239	MSE	MET	modified residue	UNP Q6RYA0
A	241	MSE	MET	modified residue	UNP Q6RYA0
A	261	LEU	-	cloning artifact	UNP Q6RYA0
A	262	GLU	-	cloning artifact	UNP Q6RYA0
A	263	HIS	-	cloning artifact	UNP Q6RYA0
A	264	HIS	-	cloning artifact	UNP Q6RYA0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	265	HIS	-	cloning artifact	UNP Q6RYA0
A	266	HIS	-	cloning artifact	UNP Q6RYA0
A	267	HIS	-	cloning artifact	UNP Q6RYA0
A	268	HIS	-	cloning artifact	UNP Q6RYA0
B	63	MSE	MET	modified residue	UNP Q6RYA0
B	66	MSE	MET	modified residue	UNP Q6RYA0
B	85	MSE	MET	modified residue	UNP Q6RYA0
B	91	MSE	MET	modified residue	UNP Q6RYA0
B	108	MSE	MET	modified residue	UNP Q6RYA0
B	149	MSE	MET	modified residue	UNP Q6RYA0
B	183	MSE	MET	modified residue	UNP Q6RYA0
B	239	MSE	MET	modified residue	UNP Q6RYA0
B	241	MSE	MET	modified residue	UNP Q6RYA0
B	261	LEU	-	cloning artifact	UNP Q6RYA0
B	262	GLU	-	cloning artifact	UNP Q6RYA0
B	263	HIS	-	cloning artifact	UNP Q6RYA0
B	264	HIS	-	cloning artifact	UNP Q6RYA0
B	265	HIS	-	cloning artifact	UNP Q6RYA0
B	266	HIS	-	cloning artifact	UNP Q6RYA0
B	267	HIS	-	cloning artifact	UNP Q6RYA0
B	268	HIS	-	cloning artifact	UNP Q6RYA0
C	63	MSE	MET	modified residue	UNP Q6RYA0
C	66	MSE	MET	modified residue	UNP Q6RYA0
C	85	MSE	MET	modified residue	UNP Q6RYA0
C	91	MSE	MET	modified residue	UNP Q6RYA0
C	108	MSE	MET	modified residue	UNP Q6RYA0
C	149	MSE	MET	modified residue	UNP Q6RYA0
C	183	MSE	MET	modified residue	UNP Q6RYA0
C	239	MSE	MET	modified residue	UNP Q6RYA0
C	241	MSE	MET	modified residue	UNP Q6RYA0
C	261	LEU	-	cloning artifact	UNP Q6RYA0
C	262	GLU	-	cloning artifact	UNP Q6RYA0
C	263	HIS	-	cloning artifact	UNP Q6RYA0
C	264	HIS	-	cloning artifact	UNP Q6RYA0
C	265	HIS	-	cloning artifact	UNP Q6RYA0
C	266	HIS	-	cloning artifact	UNP Q6RYA0
C	267	HIS	-	cloning artifact	UNP Q6RYA0
C	268	HIS	-	cloning artifact	UNP Q6RYA0
D	63	MSE	MET	modified residue	UNP Q6RYA0
D	66	MSE	MET	modified residue	UNP Q6RYA0
D	85	MSE	MET	modified residue	UNP Q6RYA0
D	91	MSE	MET	modified residue	UNP Q6RYA0

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
D	108	MSE	MET	modified residue	UNP Q6RYA0
D	149	MSE	MET	modified residue	UNP Q6RYA0
D	183	MSE	MET	modified residue	UNP Q6RYA0
D	239	MSE	MET	modified residue	UNP Q6RYA0
D	241	MSE	MET	modified residue	UNP Q6RYA0
D	261	LEU	-	cloning artifact	UNP Q6RYA0
D	262	GLU	-	cloning artifact	UNP Q6RYA0
D	263	HIS	-	cloning artifact	UNP Q6RYA0
D	264	HIS	-	cloning artifact	UNP Q6RYA0
D	265	HIS	-	cloning artifact	UNP Q6RYA0
D	266	HIS	-	cloning artifact	UNP Q6RYA0
D	267	HIS	-	cloning artifact	UNP Q6RYA0
D	268	HIS	-	cloning artifact	UNP Q6RYA0
E	63	MSE	MET	modified residue	UNP Q6RYA0
E	66	MSE	MET	modified residue	UNP Q6RYA0
E	85	MSE	MET	modified residue	UNP Q6RYA0
E	91	MSE	MET	modified residue	UNP Q6RYA0
E	108	MSE	MET	modified residue	UNP Q6RYA0
E	149	MSE	MET	modified residue	UNP Q6RYA0
E	183	MSE	MET	modified residue	UNP Q6RYA0
E	239	MSE	MET	modified residue	UNP Q6RYA0
E	241	MSE	MET	modified residue	UNP Q6RYA0
E	261	LEU	-	cloning artifact	UNP Q6RYA0
E	262	GLU	-	cloning artifact	UNP Q6RYA0
E	263	HIS	-	cloning artifact	UNP Q6RYA0
E	264	HIS	-	cloning artifact	UNP Q6RYA0
E	265	HIS	-	cloning artifact	UNP Q6RYA0
E	266	HIS	-	cloning artifact	UNP Q6RYA0
E	267	HIS	-	cloning artifact	UNP Q6RYA0
E	268	HIS	-	cloning artifact	UNP Q6RYA0
F	63	MSE	MET	modified residue	UNP Q6RYA0
F	66	MSE	MET	modified residue	UNP Q6RYA0
F	85	MSE	MET	modified residue	UNP Q6RYA0
F	91	MSE	MET	modified residue	UNP Q6RYA0
F	108	MSE	MET	modified residue	UNP Q6RYA0
F	149	MSE	MET	modified residue	UNP Q6RYA0
F	183	MSE	MET	modified residue	UNP Q6RYA0
F	239	MSE	MET	modified residue	UNP Q6RYA0
F	241	MSE	MET	modified residue	UNP Q6RYA0
F	261	LEU	-	cloning artifact	UNP Q6RYA0
F	262	GLU	-	cloning artifact	UNP Q6RYA0
F	263	HIS	-	cloning artifact	UNP Q6RYA0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	264	HIS	-	cloning artifact	UNP Q6RYA0
F	265	HIS	-	cloning artifact	UNP Q6RYA0
F	266	HIS	-	cloning artifact	UNP Q6RYA0
F	267	HIS	-	cloning artifact	UNP Q6RYA0
F	268	HIS	-	cloning artifact	UNP Q6RYA0
G	63	MSE	MET	modified residue	UNP Q6RYA0
G	66	MSE	MET	modified residue	UNP Q6RYA0
G	85	MSE	MET	modified residue	UNP Q6RYA0
G	91	MSE	MET	modified residue	UNP Q6RYA0
G	108	MSE	MET	modified residue	UNP Q6RYA0
G	149	MSE	MET	modified residue	UNP Q6RYA0
G	183	MSE	MET	modified residue	UNP Q6RYA0
G	239	MSE	MET	modified residue	UNP Q6RYA0
G	241	MSE	MET	modified residue	UNP Q6RYA0
G	261	LEU	-	cloning artifact	UNP Q6RYA0
G	262	GLU	-	cloning artifact	UNP Q6RYA0
G	263	HIS	-	cloning artifact	UNP Q6RYA0
G	264	HIS	-	cloning artifact	UNP Q6RYA0
G	265	HIS	-	cloning artifact	UNP Q6RYA0
G	266	HIS	-	cloning artifact	UNP Q6RYA0
G	267	HIS	-	cloning artifact	UNP Q6RYA0
G	268	HIS	-	cloning artifact	UNP Q6RYA0
H	63	MSE	MET	modified residue	UNP Q6RYA0
H	66	MSE	MET	modified residue	UNP Q6RYA0
H	85	MSE	MET	modified residue	UNP Q6RYA0
H	91	MSE	MET	modified residue	UNP Q6RYA0
H	108	MSE	MET	modified residue	UNP Q6RYA0
H	149	MSE	MET	modified residue	UNP Q6RYA0
H	183	MSE	MET	modified residue	UNP Q6RYA0
H	239	MSE	MET	modified residue	UNP Q6RYA0
H	241	MSE	MET	modified residue	UNP Q6RYA0
H	261	LEU	-	cloning artifact	UNP Q6RYA0
H	262	GLU	-	cloning artifact	UNP Q6RYA0
H	263	HIS	-	cloning artifact	UNP Q6RYA0
H	264	HIS	-	cloning artifact	UNP Q6RYA0
H	265	HIS	-	cloning artifact	UNP Q6RYA0
H	266	HIS	-	cloning artifact	UNP Q6RYA0
H	267	HIS	-	cloning artifact	UNP Q6RYA0
H	268	HIS	-	cloning artifact	UNP Q6RYA0

- Molecule 2 is THIOCYANATE ION (three-letter code: SCN) (formula: CNS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	S	0	0
			3	1	1	1		
2	B	1	Total	C	N	S	0	0
			3	1	1	1		
2	C	1	Total	C	N	S	0	0
			3	1	1	1		
2	D	1	Total	C	N	S	0	0
			3	1	1	1		
2	E	1	Total	C	N	S	0	0
			3	1	1	1		
2	F	1	Total	C	N	S	0	0
			3	1	1	1		
2	G	1	Total	C	N	S	0	0
			3	1	1	1		
2	H	1	Total	C	N	S	0	0
			3	1	1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	19	Total	O	0	0
			19	19		
3	B	27	Total	O	0	0
			27	27		
3	C	30	Total	O	0	0
			30	30		
3	D	25	Total	O	0	0
			25	25		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	E	28	Total 28	O 28	0	0
3	F	30	Total 30	O 30	0	0
3	G	28	Total 28	O 28	0	0
3	H	37	Total 37	O 37	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: salicylic acid-binding protein 2

Chain A: 




- Molecule 1: salicylic acid-binding protein 2


Chain B: 



- Molecule 1: salicylic acid-binding protein 2


Chain C: 






- Molecule 1: salicylic acid-binding protein 2

Chain D: 62% 33%




- Molecule 1: salicylic acid-binding protein 2

Chain E: 60% 34%




- Molecule 1: salicylic acid-binding protein 2


Chain F: 66% 27%



- Molecule 1: salicylic acid-binding protein 2


Chain G: 60% 34%





- Molecule 1: salicylic acid-binding protein 2

Chain H: 60% 35%



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	45.63Å 137.74Å 171.21Å 90.00° 90.10° 90.00°	Depositor
Resolution (Å)	28.17 – 2.52 28.17 – 2.52	Depositor EDS
% Data completeness (in resolution range)	92.3 (28.17-2.52) 97.2 (28.17-2.52)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.67 (at 2.51Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.227 , 0.297 0.228 , 0.299	Depositor DCC
R_{free} test set	6879 reflections (9.72%)	wwPDB-VP
Wilson B-factor (Å ²)	18.6	Xtrriage
Anisotropy	0.147	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 0.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.23$	Xtrriage
Estimated twinning fraction	0.388 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	16579	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SCN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/2073	0.59	0/2790
1	B	0.43	0/2094	0.58	0/2819
1	C	0.43	0/2073	0.59	0/2790
1	D	0.44	0/2094	0.61	0/2819
1	E	0.42	0/2086	0.59	0/2808
1	F	0.43	0/2085	0.58	0/2808
1	G	0.42	0/2073	0.57	0/2790
1	H	0.42	0/2094	0.59	0/2819
All	All	0.43	0/16672	0.59	0/22443

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2031	0	2013	69	0
1	B	2051	0	2028	76	0
1	C	2031	0	2013	62	0
1	D	2051	0	2028	76	0
1	E	2043	0	2022	75	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	2042	0	2015	65	0
1	G	2031	0	2013	72	0
1	H	2051	0	2028	70	0
2	A	3	0	0	1	0
2	B	3	0	0	1	0
2	C	3	0	0	0	0
2	D	3	0	0	0	0
2	E	3	0	0	0	0
2	F	3	0	0	0	0
2	G	3	0	0	0	0
2	H	3	0	0	0	0
3	A	19	0	0	1	0
3	B	27	0	0	1	0
3	C	30	0	0	2	0
3	D	25	0	0	0	0
3	E	28	0	0	4	0
3	F	30	0	0	0	0
3	G	28	0	0	1	0
3	H	37	0	0	3	0
All	All	16579	0	16160	558	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 558 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:162:GLN:HE21	1:B:237:ASP:HB2	1.28	0.96
1:C:156:LEU:HD12	1:C:160:LEU:HD12	1.49	0.95
1:E:23:LEU:HD22	1:E:248:LEU:HD13	1.50	0.93
1:H:132:LEU:HD12	1:H:155:PHE:HA	1.53	0.91
1:F:132:LEU:HD12	1:F:155:PHE:HA	1.53	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	255/268 (95%)	233 (91%)	21 (8%)	1 (0%)	30	48
1	B	257/268 (96%)	229 (89%)	26 (10%)	2 (1%)	16	29
1	C	255/268 (95%)	225 (88%)	28 (11%)	2 (1%)	16	29
1	D	257/268 (96%)	231 (90%)	22 (9%)	4 (2%)	8	14
1	E	256/268 (96%)	239 (93%)	15 (6%)	2 (1%)	16	29
1	F	256/268 (96%)	232 (91%)	23 (9%)	1 (0%)	30	48
1	G	255/268 (95%)	233 (91%)	20 (8%)	2 (1%)	16	29
1	H	257/268 (96%)	234 (91%)	22 (9%)	1 (0%)	30	48
All	All	2048/2144 (96%)	1856 (91%)	177 (9%)	15 (1%)	19	33

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	117	PHE
1	D	140	GLY
1	E	3	GLU
1	E	257	HIS
1	C	117	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	219/221 (99%)	213 (97%)	6 (3%)	40	65

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	221/221 (100%)	212 (96%)	9 (4%)	26	47
1	C	219/221 (99%)	211 (96%)	8 (4%)	29	52
1	D	221/221 (100%)	212 (96%)	9 (4%)	26	47
1	E	220/221 (100%)	210 (96%)	10 (4%)	23	43
1	F	220/221 (100%)	210 (96%)	10 (4%)	23	43
1	G	219/221 (99%)	212 (97%)	7 (3%)	34	58
1	H	221/221 (100%)	213 (96%)	8 (4%)	30	53
All	All	1760/1768 (100%)	1693 (96%)	67 (4%)	28	51

5 of 67 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	188	LYS
1	H	3	GLU
1	H	194	ASP
1	D	52	LEU
1	D	26	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 33 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	257	HIS
1	H	96	GLN
1	H	221	GLN
1	C	224	ASN
1	C	221	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SCN	G	297	-	1,2,2	0.11	0	0,1,1	-	-
2	SCN	H	298	-	1,2,2	0.68	0	0,1,1	-	-
2	SCN	B	292	-	1,2,2	0.37	0	0,1,1	-	-
2	SCN	E	295	-	1,2,2	0.41	0	0,1,1	-	-
2	SCN	F	296	-	1,2,2	0.23	0	0,1,1	-	-
2	SCN	A	291	-	1,2,2	0.16	0	0,1,1	-	-
2	SCN	D	294	-	1,2,2	0.33	0	0,1,1	-	-
2	SCN	C	293	-	1,2,2	0.07	0	0,1,1	-	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	292	SCN	1	0
2	A	291	SCN	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	248/268 (92%)	-1.68	0 100 100	2, 15, 34, 45	0
1	B	250/268 (93%)	-1.65	0 100 100	1, 17, 38, 46	0
1	C	248/268 (92%)	-1.67	0 100 100	1, 15, 34, 43	0
1	D	250/268 (93%)	-1.72	0 100 100	1, 13, 32, 49	0
1	E	249/268 (92%)	-1.72	0 100 100	1, 12, 31, 42	0
1	F	249/268 (92%)	-1.67	0 100 100	3, 17, 35, 47	0
1	G	248/268 (92%)	-1.65	0 100 100	2, 16, 35, 44	0
1	H	250/268 (93%)	-1.68	0 100 100	3, 16, 33, 49	0
All	All	1992/2144 (92%)	-1.68	0 100 100	1, 15, 34, 49	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	SCN	B	292	3/3	0.99	0.04	26,26,27,28	0
2	SCN	D	294	3/3	0.99	0.06	40,40,42,43	0
2	SCN	F	296	3/3	0.99	0.05	13,13,16,17	0
2	SCN	G	297	3/3	0.99	0.04	28,28,29,30	0
2	SCN	H	298	3/3	0.99	0.04	28,28,28,30	0
2	SCN	C	293	3/3	1.00	0.03	18,18,19,20	0
2	SCN	A	291	3/3	1.00	0.03	32,32,32,34	0
2	SCN	E	295	3/3	1.00	0.03	26,26,26,29	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.