



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2018 – 12:23 am GMT

PDB ID : 4Y8D  
Title : Crystal structure of Cyclin-G associated kinase (GAK) complexed with selective 12i inhibitor  
Authors : Chaikuad, A.; Heroven, C.; Nowak, R.; De Jonghe, S.; von Delft, F.; Arrow-smith, C.H.; Edwards, A.M.; Bountra, C.; Knapp, S.; Structural Genomics Consortium (SGC)  
Deposited on : 2015-02-16  
Resolution : 2.10 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk31020  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk31020


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	4608 (2.10-2.10)
Clashscore	122126	5109 (2.10-2.10)
Ramachandran outliers	120053	5059 (2.10-2.10)
Sidechain outliers	120020	5060 (2.10-2.10)
RSRZ outliers	108989	4497 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	340	 11% 71% 6% 23%
1	B	340	 10% 72% 5% 23%
2	C	140	 2% 81% 14%
2	D	140	 3% 84% 14%

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6441 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclin-G-associated kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	262	2083	1324	370	377	12	0	4	0
1	B	262	2095	1331	371	381	12	0	6	0


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	12	SER	-	expression tag	UNP O14976
A	13	MET	-	expression tag	UNP O14976
B	12	SER	-	expression tag	UNP O14976
B	13	MET	-	expression tag	UNP O14976

- Molecule 2 is a protein called nanobody.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	120	894	546	160	182	6	0	0	0
2	D	121	903	552	163	182	6	0	1	0

- Molecule 3 is 2-methoxy-4-[3-(morpholin-4-yl)[1,2]thiazolo[4,3-b]pyridin-6-yl]aniline (three-letter code: 49J) (formula: C<sub>17</sub>H<sub>18</sub>N<sub>4</sub>O<sub>2</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			24	17	4	2	1		
3	B	1	Total	C	N	O	S	0	0
			24	17	4	2	1		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	1
			8	4	4		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	110	Total O 110 110	0	0
5	C	96	Total O 96 96	0	0
5	B	98	Total O 98 98	0	0
5	D	90	Total O 90 90	0	0


### 3 Residue-property plots i

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cyclin-G-associated kinase




- Molecule 1: Cyclin-G-associated kinase




- Molecule 2: nanobody



- Molecule 2: nanobody





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	37.71Å 68.70Å 89.87Å 109.70° 95.08° 99.04°	Depositor
Resolution (Å)	43.58 – 2.10 42.58 – 2.10	Depositor EDS
% Data completeness (in resolution range)	94.7 (43.58-2.10) 94.7 (42.58-2.10)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.04 (at 2.10Å)	Xtrriage
Refinement program	REFMAC 5.8.0049	Depositor
R, $R_{free}$	0.192 , 0.232 0.217 , 0.250	Depositor DCC
$R_{free}$ test set	2310 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.0	Xtrriage
Anisotropy	0.253	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 43.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6441	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	70.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.56% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, 49J

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.58	0/2133	0.70	3/2881 (0.1%)
1	B	0.58	0/2151	0.70	3/2905 (0.1%)
2	C	0.80	0/907	0.83	1/1226 (0.1%)
2	D	0.78	0/919	0.85	1/1242 (0.1%)
All	All	0.65	0/6110	0.74	8/8254 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	50	ARG	NE-CZ-NH1	6.79	123.70	120.30
1	B	293	ARG	NE-CZ-NH1	-6.47	117.06	120.30
1	B	293	ARG	NE-CZ-NH2	6.36	123.48	120.30
1	A	293	ARG	NE-CZ-NH1	-5.60	117.50	120.30
1	A	293	ARG	NE-CZ-NH2	5.29	122.95	120.30
1	A	157	ARG	NE-CZ-NH2	-5.15	117.72	120.30
1	B	157	ARG	NE-CZ-NH2	-5.12	117.74	120.30
2	C	50	ARG	NE-CZ-NH1	5.01	122.81	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	236	SER	Peptide
1	B	236	SER	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2083	0	2097	15	0
1	B	2095	0	2112	11	0
2	C	894	0	858	5	0
2	D	903	0	867	6	0
3	A	24	0	18	1	0
3	B	24	0	18	1	0
4	B	4	0	6	0	0
4	C	12	0	18	0	0
4	D	8	0	12	0	0
5	A	110	0	0	4	0
5	B	98	0	0	1	0
5	C	96	0	0	3	0
5	D	90	0	0	4	0
All	All	6441	0	6006	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:105:ARG:NH2	5:D:341:HOH:O	2.24	0.69
2:D:62[A]:ARG:NH1	5:D:301:HOH:O	2.27	0.67
1:A:37:GLU:N	5:A:559:HOH:O	2.29	0.64
3:A:401:49J:NAL	3:A:401:49J:H7	2.12	0.64
1:A:297:GLN:HG3	5:A:501:HOH:O	1.98	0.63
1:B:261:HIS:CD2	1:B:262:PRO:HD2	2.39	0.57
2:C:105:ARG:NH2	5:C:350:HOH:O	2.36	0.56

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:261:HIS:CD2	1:A:262:PRO:HD2	2.40	0.56
1:A:236:SER:HA	5:A:555:HOH:O	2.06	0.55
1:B:40:LEU:HD13	1:B:57:ALA:HB1	1.91	0.52
1:A:40:LEU:HD13	1:A:57:ALA:HB1	1.92	0.52
2:D:28:THR:O	2:D:50:ARG:HD2	2.12	0.50
1:B:297:GLN:HG3	5:B:501:HOH:O	2.12	0.49
1:A:241:GLY:HA2	1:B:76:GLU:CD	2.34	0.48
1:A:62:SER:OG	1:A:64:ARG:NH2	2.47	0.48
2:C:28:THR:O	2:C:50:ARG:HD2	2.14	0.47
2:D:62[A]:ARG:CZ	5:D:301:HOH:O	2.63	0.45
1:B:261:HIS:CG	1:B:262:PRO:HD2	2.52	0.45
1:A:261:HIS:CG	1:A:262:PRO:HD2	2.52	0.45
1:A:100[B]:GLN:NE2	5:A:554:HOH:O	2.34	0.44
1:A:241:GLY:HA2	1:B:76:GLU:HG3	1.99	0.44
1:A:40:LEU:HB3	1:A:59:ASP:HA	2.00	0.43
1:A:89:MET:SD	1:A:99:VAL:HG23	2.59	0.43
1:A:76:GLU:CD	1:B:241:GLY:HA2	2.40	0.43
2:C:42:ARG:NH2	5:C:393:HOH:O	2.43	0.42
2:C:42:ARG:NH1	5:C:387:HOH:O	2.53	0.42
3:B:401:49J:CAK	3:B:401:49J:NAL	2.83	0.42
1:B:40:LEU:HB3	1:B:59:ASP:HA	2.01	0.42
1:B:89:MET:SD	1:B:99:VAL:HG23	2.60	0.42
1:A:72:LEU:CD2	1:A:118:GLU:HG2	2.50	0.42
2:D:62[A]:ARG:NH2	5:D:301:HOH:O	2.53	0.42
1:A:110:GLU:H	1:A:110:GLU:CD	2.23	0.42
2:D:62[B]:ARG:HG3	2:D:62[B]:ARG:HH21	1.84	0.42
1:B:110:GLU:H	1:B:110:GLU:CD	2.23	0.41
2:C:84:LYS:HB2	2:C:85:PRO:HD2	2.01	0.41
1:B:72:LEU:CD2	1:B:118:GLU:HG2	2.50	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/340 (76%)	251 (96%)	9 (4%)	0	100	100
1	B	262/340 (77%)	249 (95%)	13 (5%)	0	100	100
2	C	118/140 (84%)	117 (99%)	1 (1%)	0	100	100
2	D	120/140 (86%)	120 (100%)	0	0	100	100
All	All	760/960 (79%)	737 (97%)	23 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	227/294 (77%)	227 (100%)	0	100	100
1	B	230/294 (78%)	230 (100%)	0	100	100
2	C	94/110 (86%)	94 (100%)	0	100	100
2	D	94/110 (86%)	94 (100%)	0	100	100
All	All	645/808 (80%)	645 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	49J	A	401	-	23,27,27	2.12	8 (34%)	29,38,38	2.61	11 (37%)
3	49J	B	401	-	23,27,27	2.22	6 (26%)	29,38,38	2.47	12 (41%)
4	EDO	B	402	-	3,3,3	0.43	0	2,2,2	0.26	0
4	EDO	C	201	-	3,3,3	0.42	0	2,2,2	0.48	0
4	EDO	C	202[A]	-	3,3,3	0.75	0	2,2,2	0.20	0
4	EDO	C	202[B]	-	3,3,3	0.64	0	2,2,2	0.11	0
4	EDO	D	201	-	3,3,3	0.51	0	2,2,2	0.78	0
4	EDO	D	202	-	3,3,3	0.64	0	2,2,2	0.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	49J	A	401	-	-	0/6/18/18	0/4/4/4
3	49J	B	401	-	-	0/6/18/18	0/4/4/4
4	EDO	B	402	-	-	0/1/1/1	0/0/0/0
4	EDO	C	201	-	-	0/1/1/1	0/0/0/0
4	EDO	C	202[A]	-	-	0/1/1/1	0/0/0/0
4	EDO	C	202[B]	-	-	0/1/1/1	0/0/0/0
4	EDO	D	201	-	-	0/1/1/1	0/0/0/0
4	EDO	D	202	-	-	0/1/1/1	0/0/0/0

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	401	49J	CAG-CAV	-5.18	1.33	1.41
3	A	401	49J	CAG-CAV	-4.88	1.34	1.41
3	A	401	49J	CAU-CAW	-4.14	1.33	1.42
3	B	401	49J	CAU-CAW	-3.95	1.33	1.42
3	A	401	49J	CAR-CAS	-3.52	1.40	1.49
3	B	401	49J	CAR-CAS	-2.64	1.42	1.49
3	A	401	49J	CAJ-CAH	-2.50	1.40	1.50
3	B	401	49J	CAQ-CAT	-2.49	1.37	1.40
3	A	401	49J	CAW-CAV	-2.14	1.35	1.42
3	A	401	49J	CAW-NAL	-2.13	1.34	1.37
3	A	401	49J	CAK-NAX	2.23	1.50	1.46
3	B	401	49J	CAJ-NAX	2.42	1.50	1.46
3	A	401	49J	CAE-NAL	4.10	1.38	1.31
3	B	401	49J	CAE-NAL	5.90	1.41	1.31

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	401	49J	CAS-CAE-NAL	-6.70	118.65	125.51
3	B	401	49J	CAS-CAE-NAL	-5.81	119.56	125.51
3	B	401	49J	CAT-CAQ-NAB	-5.55	114.89	119.45
3	A	401	49J	CAV-CAW-NAL	-4.52	119.63	125.00
3	B	401	49J	CAV-CAW-NAL	-4.15	120.07	125.00
3	A	401	49J	OAO-CAH-CAJ	-3.90	103.25	111.81
3	B	401	49J	OAO-CAI-CAK	-3.74	103.60	111.81
3	A	401	49J	CAJ-NAX-CAU	-3.49	116.90	121.29
3	A	401	49J	CAH-CAJ-NAX	-3.10	104.30	109.92
3	B	401	49J	CAF-CAT-CAQ	-2.70	119.16	120.92
3	A	401	49J	CAF-CAT-CAQ	-2.58	119.23	120.92
3	B	401	49J	CAH-CAJ-NAX	-2.54	105.31	109.92
3	A	401	49J	CAT-CAQ-NAB	-2.40	117.48	119.45
3	B	401	49J	CAG-CAV-NAM	-2.07	125.42	130.22
3	B	401	49J	CAI-OAO-CAH	2.03	116.71	109.89
3	B	401	49J	OAN-CAT-CAF	2.31	128.06	124.15
3	A	401	49J	OAN-CAT-CAQ	2.36	115.64	114.05
3	B	401	49J	CAK-NAX-CAJ	2.89	117.75	111.54
3	B	401	49J	CAE-NAL-CAW	3.25	120.05	116.76
3	B	401	49J	CAC-CAQ-CAT	3.68	121.37	118.27
3	A	401	49J	CAK-NAX-CAJ	3.88	119.89	111.54
3	A	401	49J	CAC-CAQ-CAT	3.93	121.58	118.27
3	A	401	49J	CAE-NAL-CAW	5.33	122.14	116.76

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	401	49J	1	0
3	B	401	49J	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	262/340 (77%)	0.83	39 (14%) <b>2</b> <b>3</b>	32, 74, 155, 170	2 (0%)
1	B	262/340 (77%)	0.74	34 (12%) <b>3</b> <b>4</b>	31, 70, 149, 186	1 (0%)
2	C	120/140 (85%)	0.35	3 (2%) 57 63	31, 50, 78, 105	0
2	D	121/140 (86%)	0.38	4 (3%) 46 54	32, 51, 76, 90	1 (0%)
All	All	765/960 (79%)	0.65	80 (10%) <b>6</b> <b>8</b>	31, 61, 148, 186	4 (0%)

All (80) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	61	GLY	8.3
1	A	50	GLY	7.0
1	A	51	PHE	6.6
1	A	112	SER	5.7
1	B	36	GLY	5.4
1	B	49	GLY	5.2
1	A	107	ILE	5.2
1	B	62	SER	5.1
1	B	64	ARG	5.0
1	A	49	GLY	4.8
1	B	35	LEU	4.7
1	B	60	VAL	4.5
2	D	121	SER	4.4
1	A	141	GLY	4.4
1	B	51	PHE	4.3
1	A	113	ASP	4.0
2	C	1	GLN	3.9
1	A	114	THR	3.8
1	A	108	GLY	3.7
1	B	112	SER	3.7
1	A	61	GLY	3.7

*Continued on next page...*



*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	52	ALA	3.6
1	A	263	PHE	3.6
1	A	139	SER	3.5
1	A	236	SER	3.5
1	A	60	VAL	3.5
1	A	135	LYS	3.5
1	A	333	GLU	3.3
1	B	235	TYR	3.3
1	B	107	ILE	3.3
1	B	38	LEU	3.2
1	B	50	GLY	3.2
1	A	47	ALA	3.2
1	A	48	GLU	3.1
1	B	331	LEU	3.0
1	B	263	PHE	2.8
1	B	57	ALA	2.8
1	B	63	GLY	2.7
1	A	37	GLU	2.7
1	B	37	GLU	2.7
1	A	57	ALA	2.7
1	A	235	TYR	2.7
1	A	331	LEU	2.7
1	B	113	ASP	2.6
1	A	38	LEU	2.6
1	A	332	LEU	2.6
1	B	47	ALA	2.6
1	A	201	TYR	2.5
1	A	43	ARG	2.5
1	A	55	TYR	2.4
1	B	43	ARG	2.4
1	A	202	PRO	2.4
1	B	30	GLY	2.4
1	B	201	TYR	2.4
1	B	202	PRO	2.3
1	A	74	ASN	2.3
2	D	1	GLN	2.3
1	B	46	LEU	2.3
2	C	2	VAL	2.3
1	B	137	MET	2.3
1	B	74	ASN	2.3
1	A	118	GLU	2.2
1	A	29	VAL	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	277	SER	2.2
1	A	136	LYS	2.2
1	A	75	GLU	2.1
2	D	39	GLY	2.1
1	B	261	HIS	2.1
1	A	200	HIS	2.1
1	A	191	ASP	2.1
2	D	12	VAL	2.1
1	A	277	SER	2.0
1	B	139	SER	2.0
1	B	332	LEU	2.0
1	A	28	PHE	2.0
1	B	140	ARG	2.0
1	A	111	GLU	2.0
1	B	77	GLU	2.0
1	B	333	GLU	2.0
2	C	113	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	EDO	C	201	4/4	0.80	0.25	58,64,64,73	0
4	EDO	B	402	4/4	0.85	0.19	63,64,65,66	0
4	EDO	C	202[A]	4/4	0.86	0.21	39,39,40,41	4
4	EDO	C	202[B]	4/4	0.86	0.21	35,37,39,40	4
3	49J	A	401	24/24	0.92	0.26	38,49,59,60	0
4	EDO	D	201	4/4	0.94	0.15	34,37,40,46	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	EDO	D	202	4/4	0.94	0.13	47,50,53,53	0
3	49J	B	401	24/24	0.94	0.22	35,45,49,49	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.