



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 14, 2024 – 12:10 am GMT

PDB ID : 6YAX  
Title : Crystal structure of CD32b (Fc Gamma Receptor IIb) in complex with Human IgG1 Fab fragment (5C05)  
Authors : Fisher, H.; Tews, I.; Orr, C.  
Deposited on : 2020-03-14  
Resolution : 2.80 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	HHH	218	
1	III	218	
2	LLL	218	
2	MMM	218	
3	AAA	181	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	BBB	181	<p>2% 79% 11% 9%</p>
4	A	2	<p>100%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	A	2	X	-	-	-

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 16762 atoms, of which 8097 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 5C05 F(ab) heavy chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	HHH	204	3053	981	1509	257	300	6	87	0	0
1	III	167	2441	790	1190	211	246	4	79	0	0

- Molecule 2 is a protein called 5C05 F(ab) light chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	MMM	182	2530	811	1225	218	272	4	94	0	0
2	LLL	214	3096	983	1524	264	321	4	100	0	0

- Molecule 3 is a protein called Low affinity immunoglobulin gamma Fc region receptor II-c.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
3	AAA	165	2610	841	1285	232	248	4	98	1	0
3	BBB	164	2595	836	1278	227	250	4	97	2	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AAA	218	HIS	-	expression tag	UNP P31995
AAA	219	HIS	-	expression tag	UNP P31995
AAA	220	HIS	-	expression tag	UNP P31995
AAA	221	HIS	-	expression tag	UNP P31995
AAA	222	HIS	-	expression tag	UNP P31995
AAA	223	HIS	-	expression tag	UNP P31995
BBB	218	HIS	-	expression tag	UNP P31995
BBB	219	HIS	-	expression tag	UNP P31995

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
BBB	220	HIS	-	expression tag	UNP P31995
BBB	221	HIS	-	expression tag	UNP P31995
BBB	222	HIS	-	expression tag	UNP P31995
BBB	223	HIS	-	expression tag	UNP P31995

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
			Total	C	H	N				O
4	A	2	56	16	28	2	10	5	0	0

- Molecule 5 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: C<sub>8</sub>H<sub>18</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
5	HHH	1	31	8	18	5	1	0
5	III	1	31	8	18	5	1	0

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
6	AAA	1	28	8	14	1	5	3	0

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
7	LLL	1	14	3	8	3	2	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	HHH	53	Total 53	O 53	0	0
8	MMM	34	Total 34	O 34	0	0
8	AAA	34	Total 34	O 34	0	0
8	LLL	40	Total 40	O 40	0	0
8	III	42	Total 42	O 42	0	0
8	BBB	74	Total 74	O 74	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 5C05 F(ab) heavy chain




- Molecule 1: 5C05 F(ab) heavy chain




- Molecule 2: 5C05 F(ab) light chain





- Molecule 2: 5C05 F(ab) light chain









- Molecule 3: Low affinity immunoglobulin gamma Fc region receptor II-c



- Molecule 3: Low affinity immunoglobulin gamma Fc region receptor II-c



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	95.73Å 95.73Å 186.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.86 – 2.80 47.86 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.6 (47.86-2.80) 99.6 (47.86-2.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.53 (at 2.81Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, $R_{free}$	0.230 , 0.264 0.232 , 0.267	Depositor DCC
$R_{free}$ test set	1971 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	65.5	Xtrriage
Anisotropy	0.027	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 80.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.037 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	16762	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	77.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.59% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, PG4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	HHH	0.64	0/1581	0.78	0/2152
1	III	0.80	4/1279 (0.3%)	0.80	1/1738 (0.1%)
2	LLL	0.66	0/1611	0.78	0/2202
2	MMM	0.67	0/1330	0.81	0/1814
3	AAA	0.61	0/1368	0.78	0/1865
3	BBB	0.61	0/1361	0.76	0/1855
All	All	0.67	4/8530 (0.0%)	0.78	1/11626 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	III	67	ARG	CZ-NH2	10.91	1.47	1.33
1	III	38	ARG	CZ-NH2	-7.55	1.23	1.33
1	III	67	ARG	CZ-NH1	7.33	1.42	1.33
1	III	67	ARG	CD-NE	5.51	1.55	1.46

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	III	67	ARG	NE-CZ-NH1	-5.33	117.64	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	HHH	1544	1509	1504	23	1
1	III	1251	1190	1170	18	0
2	LLL	1572	1524	1514	27	0
2	MMM	1305	1225	1196	30	0
3	AAA	1325	1285	1273	14	1
3	BBB	1317	1278	1268	16	0
4	A	28	28	25	0	0
5	HHH	13	18	18	0	0
5	III	13	18	18	1	0
6	AAA	14	14	13	0	0
7	LLL	6	8	8	0	0
8	AAA	34	0	0	1	1
8	BBB	74	0	0	1	1
8	HHH	53	0	0	0	0
8	III	42	0	0	1	0
8	LLL	40	0	0	2	0
8	MMM	34	0	0	2	0
All	All	8665	8097	8007	126	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 126 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:HHH:188:VAL:HB	1:HHH:189:PRO:HD2	1.34	1.06
2:MMM:82:SER:HA	2:MMM:111:VAL:HG21	1.61	0.80
1:HHH:188:VAL:HB	1:HHH:189:PRO:CD	2.12	0.80
2:LLL:172:LYS:HD3	2:LLL:178:TYR:CZ	2.17	0.79
2:LLL:87:ASP:OD1	2:LLL:108:LYS:HG2	1.88	0.74

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:HHH:208:ASN:HD22	3:AAA:195:HIS:HD2[4_555]	1.19	0.41
8:AAA:433:HOH:O	8:BBB:466:HOH:O[1_655]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	HHH	198/218 (91%)	188 (95%)	9 (4%)	1 (0%)	29	61
1	III	159/218 (73%)	147 (92%)	12 (8%)	0	100	100
2	LLL	212/218 (97%)	202 (95%)	10 (5%)	0	100	100
2	MMM	168/218 (77%)	155 (92%)	13 (8%)	0	100	100
3	AAA	162/181 (90%)	157 (97%)	5 (3%)	0	100	100
3	BBB	162/181 (90%)	155 (96%)	7 (4%)	0	100	100
All	All	1061/1234 (86%)	1004 (95%)	56 (5%)	1 (0%)	51	81

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	HHH	190	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	HHH	172/182 (94%)	170 (99%)	2 (1%)	71	92
1	III	132/182 (72%)	132 (100%)	0	100	100
2	LLL	174/181 (96%)	174 (100%)	0	100	100
2	MMM	139/181 (77%)	135 (97%)	4 (3%)	42	76
3	AAA	153/166 (92%)	152 (99%)	1 (1%)	84	95
3	BBB	153/166 (92%)	152 (99%)	1 (1%)	84	95

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	923/1058 (87%)	915 (99%)	8 (1%)	78 94

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	BBB	98	SER
3	AAA	181	PHE
2	MMM	173	GLN
2	MMM	128	SER
2	MMM	184	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	NAG	A	1	3,4	14,14,15	0.51	0	17,19,21	1.06	2 (11%)
4	NAG	A	2	4	14,14,15	0.60	0	17,19,21	2.32	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1	3,4	-	2/6/23/26	0/1/1/1
4	NAG	A	2	4	1/1/7/7	1/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2	NAG	C1-O5-C5	6.60	121.13	112.19
4	A	2	NAG	C1-C2-N2	4.31	117.86	110.49
4	A	2	NAG	C2-N2-C7	3.73	128.22	122.90
4	A	1	NAG	C1-O5-C5	2.94	116.17	112.19
4	A	2	NAG	O5-C5-C6	2.05	110.42	107.20

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	A	2	NAG	C1


All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	2	NAG	C1-C2-N2-C7
4	A	1	NAG	C4-C5-C6-O6
4	A	1	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
7	GOL	LLL	301	-	5,5,5	0.11	0	5,5,5	0.33	0
6	NAG	AAA	301	3	14,14,15	0.54	0	17,19,21	1.31	2 (11%)
5	PG4	HHH	301	-	12,12,12	0.16	0	11,11,11	0.18	0
5	PG4	III	301	-	12,12,12	0.15	0	11,11,11	0.20	0



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	LLL	301	-	-	1/4/4/4	-
6	NAG	AAA	301	3	-	0/6/23/26	0/1/1/1
5	PG4	HHH	301	-	-	6/10/10/10	-
5	PG4	III	301	-	-	5/10/10/10	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	AAA	301	NAG	C1-C2-N2	-2.50	106.21	110.49
6	AAA	301	NAG	C1-O5-C5	2.14	115.09	112.19

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	HHH	301	PG4	O3-C5-C6-O4
5	III	301	PG4	O3-C5-C6-O4
5	HHH	301	PG4	O1-C1-C2-O2
5	III	301	PG4	O2-C3-C4-O3
5	HHH	301	PG4	O4-C7-C8-O5

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	III	301	PG4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	HHH	204/218 (93%)	0.53	9 (4%) 34 24	53, 72, 108, 134	0
1	III	167/218 (76%)	0.64	12 (7%) 15 8	47, 75, 120, 126	0
2	LLL	214/218 (98%)	0.49	11 (5%) 28 19	52, 79, 104, 121	0
2	MMM	182/218 (83%)	0.68	13 (7%) 16 9	58, 89, 115, 141	0
3	AAA	165/181 (91%)	0.68	16 (9%) 7 4	55, 74, 111, 135	0
3	BBB	164/181 (90%)	0.36	4 (2%) 59 49	43, 62, 93, 109	0
All	All	1096/1234 (88%)	0.56	65 (5%) 22 14	43, 76, 113, 141	0

The worst 5 of 65 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	AAA	117	GLN	5.9
3	BBB	82	ILE	4.3
2	MMM	199	CYS	4.1
3	AAA	95	THR	3.9
3	AAA	82	ILE	3.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	NAG	A	2	14/15	0.85	0.25	104,106,116,119	3
4	NAG	A	1	14/15	0.89	0.15	86,100,112,112	2

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	NAG	AAA	301	14/15	0.68	0.38	105,108,112,113	3
7	GOL	LLL	301	6/6	0.86	0.21	69,75,81,81	2
5	PG4	III	301	13/13	0.87	0.34	63,71,79,79	1
5	PG4	HHH	301	13/13	0.91	0.29	69,73,83,83	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.