



# wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 14, 2023 – 05:28 pm GMT

PDB ID : 2YBF  
Title : Complex of Rad18 (Rad6 binding domain) with Rad6b  
Authors : Hibbert, R.G.; Sixma, T.K.  
Deposited on : 2011-03-08  
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtrriage (Phenix) : 1.13  
EDS : **FAILED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36

## 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

There are no overall percentile quality scores available for this entry.

ENTRY-COMPOSITION INFOmissingINFO

SEQUENCE-PLOTS INFOmissingINFO

## 2 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	58.21Å 58.21Å 167.07Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	24.92 – 2.00	Depositor
% Data completeness (in resolution range)	99.3 (24.92-2.00)	Depositor
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.62 (at 1.99Å)	Xtrriage
Refinement program	REFMAC 5.5.0099	Depositor
R, $R_{free}$	0.199 , 0.242	Depositor
Wilson B-factor (Å <sup>2</sup> )	24.6	Xtrriage
Anisotropy	0.198	Xtrriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	1532	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.56% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

### 3 Model quality [i](#)

#### 3.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, SO4, BME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	1/1291 (0.1%)	0.50	0/1757
2	B	0.32	0/197	0.41	0/258
All	All	0.37	1/1488 (0.1%)	0.49	0/2015

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	133	ASN	C-N	-5.22	1.22	1.34

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

#### 3.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

#### 3.3 Torsion angles [i](#)

##### 3.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

##### 3.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

### 3.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 3.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 3.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 3.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 could not be matched to an existing wwPDB Chemical Component Dictionary definition at this stage - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 3.7 Other polymers [i](#)

There are no such residues in this entry.

### 3.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 4 Fit of model and data

### 4.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

### 4.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

### 4.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

### 4.4 Ligands

EDS failed to run properly - this section is therefore empty.

### 4.5 Other polymers

EDS failed to run properly - this section is therefore empty.