



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 14, 2023 – 04:02 AM JST

PDB ID : 7YCM

Title : TR-SFX MmCPDII-DNA complex: 100 ps snapshot. Includes 100ps, dark, and extrapolated structure factors

Authors : Maestre-Reyna, M.; Wang, P.-H.; Nango, E.; Hosokawa, Y.; Saft, M.; Furrer, A.; Yang, C.-H.; Ngura Putu, E.P.G.; Wu, W.-J.; Emmerich, H.-J.; Engilberge, S.; Caramello, N.; Wranik, M.; Glover, H.L.; Franz-Badur, S.; Wu, H.-Y.; Lee, C.-C.; Huang, W.-C.; Huang, K.-F.; Chang, Y.-K.; Liao, J.-H.; Weng, J.-H.; Gad, W.; Chang, C.-W.; Pang, A.H.; Gashi, D.; Beale, E.; Ozerov, D.; Milne, C.; Cirelli, C.; Bacellar, C.; Sugahara, M.; Owada, S.; Joti, Y.; Yamashita, A.; Tanaka, R.; Tanaka, T.; Luo, F.J.; Tono, K.; Kiontke, S.; Spadaccini, R.; Royant, A.; Yamamoto, J.; Iwata, S.; Standfuss, J.; Essen, L.-O.; Bessho, Y.; Tsai, M.-D.

Deposited on : 2022-07-01

Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)






The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	482	 82% 13% 5%

Continued on next page...

EDS : 2.36
 buster-report : 1.1.7 (2018)
 Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
 Refmac : 5.8.0158
 CCP4 : 7.0.044 (Gargrove)
 Ideal geometry (proteins) : Engh & Huber (2001)
 Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
 Validation Pipeline (wwPDB-VP) : 2.36

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	B	482	 80% 9% 10%
2	C	14	 43% 29% 7% 21%
2	E	14	 29% 43% 29%
3	D	14	 93% 7%
3	F	14	 64% 29% 7%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 8700 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Deoxyribodipyrimidine photolyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	456	3683	2376	606	686	15	0	11	0
1	B	434	3512	2263	588	646	15	0	10	0

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-17	MET	-	initiating methionine	UNP A0A0F8I5V2
A	-16	GLY	-	expression tag	UNP A0A0F8I5V2
A	-15	SER	-	expression tag	UNP A0A0F8I5V2
A	-14	SER	-	expression tag	UNP A0A0F8I5V2
A	-13	HIS	-	expression tag	UNP A0A0F8I5V2
A	-12	HIS	-	expression tag	UNP A0A0F8I5V2
A	-11	HIS	-	expression tag	UNP A0A0F8I5V2
A	-10	HIS	-	expression tag	UNP A0A0F8I5V2
A	-9	HIS	-	expression tag	UNP A0A0F8I5V2
A	-8	HIS	-	expression tag	UNP A0A0F8I5V2
A	-7	SER	-	expression tag	UNP A0A0F8I5V2
A	-6	SER	-	expression tag	UNP A0A0F8I5V2
A	-5	GLY	-	expression tag	UNP A0A0F8I5V2
A	-4	LEU	-	expression tag	UNP A0A0F8I5V2
A	-3	VAL	-	expression tag	UNP A0A0F8I5V2
A	-2	PRO	-	expression tag	UNP A0A0F8I5V2
A	-1	ARG	-	expression tag	UNP A0A0F8I5V2
A	0	GLY	-	expression tag	UNP A0A0F8I5V2
A	1	SER	-	expression tag	UNP A0A0F8I5V2
A	2	HIS	-	expression tag	UNP A0A0F8I5V2
A	377	THR	MET	engineered mutation	UNP A0A0F8I5V2
B	-17	MET	-	initiating methionine	UNP A0A0F8I5V2
B	-16	GLY	-	expression tag	UNP A0A0F8I5V2
B	-15	SER	-	expression tag	UNP A0A0F8I5V2
B	-14	SER	-	expression tag	UNP A0A0F8I5V2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-13	HIS	-	expression tag	UNP A0A0F8I5V2
B	-12	HIS	-	expression tag	UNP A0A0F8I5V2
B	-11	HIS	-	expression tag	UNP A0A0F8I5V2
B	-10	HIS	-	expression tag	UNP A0A0F8I5V2
B	-9	HIS	-	expression tag	UNP A0A0F8I5V2
B	-8	HIS	-	expression tag	UNP A0A0F8I5V2
B	-7	SER	-	expression tag	UNP A0A0F8I5V2
B	-6	SER	-	expression tag	UNP A0A0F8I5V2
B	-5	GLY	-	expression tag	UNP A0A0F8I5V2
B	-4	LEU	-	expression tag	UNP A0A0F8I5V2
B	-3	VAL	-	expression tag	UNP A0A0F8I5V2
B	-2	PRO	-	expression tag	UNP A0A0F8I5V2
B	-1	ARG	-	expression tag	UNP A0A0F8I5V2
B	0	GLY	-	expression tag	UNP A0A0F8I5V2
B	1	SER	-	expression tag	UNP A0A0F8I5V2
B	2	HIS	-	expression tag	UNP A0A0F8I5V2
B	377	THR	MET	engineered mutation	UNP A0A0F8I5V2


- Molecule 2 is a DNA chain called CPD photolesion containing DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	11	Total	C	N	O	P	0	0	0
			244	115	44	73	12			
2	E	10	Total	C	N	O	P	0	0	0
			225	106	41	67	11			

- Molecule 3 is a DNA chain called complementary oligonucleotide to the CPD containing DNA.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	14	Total	C	N	O	P	0	0	0
			286	136	56	81	13			
3	F	13	Total	C	N	O	P	0	0	0
			266	126	54	74	12			

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
5	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		


- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	196	Total 196	O 196	0	0
6	B	132	Total 132	O 132	0	0
6	C	19	Total 19	O 19	0	0
6	D	8	Total 8	O 8	0	0
6	E	5	Total 5	O 5	0	0
6	F	13	Total 13	O 13	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Deoxyribodipyrimidine photolyase

Chain A: 



- Molecule 1: Deoxyribodipyrimidine photolyase

Chain B: 




- Molecule 2: CPD photolesion containing DNA

Chain C: 



- Molecule 2: CPD photolesion containing DNA

Chain E: 



- Molecule 3: complementary oligonucleotide to the CPD containing DNA

Chain D:
93% 7%

A horizontal progress bar for Chain D. It is 93% green and 7% yellow.



- Molecule 3: complementary oligonucleotide to the CPD containing DNA

Chain F:
64% 29% 7%

A horizontal progress bar for Chain F. It is 64% green, 29% yellow, and 7% grey.



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	72.47Å 115.93Å 169.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	16.76 – 2.00 16.76 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (16.76-2.00) 99.9 (16.76-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.36 (at 2.00Å)	Xtrriage
Refinement program	PHENIX (1.19_4092: ???)	Depositor
R, R_{free}	0.193 , 0.212 0.194 , 0.212	Depositor DCC
R_{free} test set	4771 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	23.0	Xtrriage
Anisotropy	0.036	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8700	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TTD, FAD, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.27	0/3817	0.46	0/5174
1	B	0.24	0/3639	0.44	0/4935
2	C	0.45	0/227	0.70	0/345
2	E	0.53	0/206	0.75	0/313
3	D	0.45	0/321	0.79	0/494
3	F	0.48	0/299	0.74	0/460
All	All	0.29	0/8509	0.50	0/11721

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3683	0	3528	32	1
1	B	3512	0	3345	27	0
2	C	244	0	135	8	0
2	E	225	0	124	4	0
3	D	286	0	158	1	0
3	F	266	0	146	2	0
4	A	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	53	0	31	0	0
5	B	53	0	31	1	0
6	A	196	0	0	3	0
6	B	132	0	0	3	1
6	C	19	0	0	1	0
6	D	8	0	0	2	0
6	E	5	0	0	1	0
6	F	13	0	0	0	0
All	All	8700	0	7498	69	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:188:PRO:HG2	6:A:787:HOH:O	1.71	0.89
1:A:89:PRO:HG3	1:A:188:PRO:HG3	1.67	0.75
1:B:316:ASN:ND2	6:B:603:HOH:O	2.22	0.71
1:B:185:SER:O	6:B:601:HOH:O	2.07	0.71
2:C:10:DC:N4	6:D:101:HOH:O	2.28	0.66
3:F:4:DG:H2''	3:F:5:DC:H5''	1.78	0.65
1:B:460:LYS:NZ	1:B:460:LYS:HB2	2.11	0.64
3:F:1:DT:H2''	3:F:2:DG:OP1	2.00	0.60
1:A:451:LYS:NZ	2:C:9:DG:OP2	2.31	0.59
1:B:357:ASP:OD2	1:B:380:TYR:OH	2.14	0.58
1:B:307:GLU:OE1	6:B:602:HOH:O	2.17	0.58
1:A:21:VAL:HG13	1:A:112:THR:HB	1.86	0.58
1:B:105:VAL:HG13	1:B:110:ALA:HB3	1.86	0.57
1:B:460:LYS:HB2	1:B:460:LYS:HZ3	1.71	0.56
1:A:359:LEU:HD13	1:A:454:VAL:HG13	1.86	0.56
3:D:11:DC:N4	6:D:103:HOH:O	2.38	0.56
1:B:26[B]:SER:OG	1:B:116:ASP:OD1	2.24	0.55
1:A:161:HIS:HB3	2:C:7:TTD:OP2	2.08	0.53
1:A:291:ASN:HB3	1:A:294:SER:HB2	1.89	0.53
1:A:121[A]:ARG:NH2	1:A:320:ASP:OD1	2.39	0.53
1:B:121:ARG:NH2	1:B:320:ASP:OD1	2.39	0.52
1:B:450[A]:ARG:NH2	2:E:11:DG:O6	2.42	0.52
1:B:291:ASN:HB3	1:B:294:SER:HB2	1.92	0.52
1:B:302:ILE:HG23	1:B:303:LEU:HD22	1.91	0.52
1:A:376:TYR:OH	6:A:601:HOH:O	2.19	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:351:GLU:OE2	1:B:406:TYR:OH	2.21	0.51
2:C:4:DG:N3	6:C:101:HOH:O	2.34	0.51
1:A:383:LYS:HD2	1:A:443:MET:HG3	1.93	0.50
2:E:4:DG:H4'	2:E:5:DG:OP1	2.11	0.50
1:B:435:GLU:O	1:B:437:THR:N	2.45	0.49
1:A:357:ASP:OD2	1:A:380:TYR:OH	2.24	0.49
1:A:61:PHE:HD2	1:A:211:LEU:HD11	1.78	0.49
1:B:19:GLY:HA3	1:B:109:ASN:O	2.12	0.49
1:A:397[B]:GLU:OE1	6:A:602:HOH:O	2.20	0.47
1:B:244:PHE:CD2	1:B:265:SER:HA	2.50	0.47
1:A:336:HIS:NE2	1:A:433:GLU:OE2	2.48	0.47
1:A:379:MET:SD	2:C:7:TTD:H71	2.55	0.46
1:B:358:PRO:HB2	1:B:458:ILE:HD13	1.98	0.46
1:B:372:LYS:NZ	5:B:501:FAD:O2P	2.38	0.46
1:A:252:TYR:O	1:A:256[A]:ARG:HB3	2.16	0.46
1:A:41:ARG:NH2	1:A:183:PRO:O	2.45	0.46
1:B:454:VAL:O	1:B:458:ILE:HG13	2.16	0.46
1:A:108:TYR:OH	1:A:188:PRO:HB3	2.15	0.46
1:B:129:LYS:HA	1:B:129:LYS:HD3	1.75	0.45
1:A:345:TYR:CE2	1:A:355:THR:HA	2.51	0.45
2:E:10:DC:H2''	2:E:11:DG:C8	2.50	0.45
1:A:171:LEU:HD23	1:A:171:LEU:HA	1.81	0.44
1:A:266:ASN:HA	1:A:372:LYS:HE3	1.98	0.44
1:B:384:LYS:HA	1:B:384:LYS:HD2	1.71	0.44
1:A:89:PRO:HG3	1:A:188:PRO:CG	2.44	0.44
1:A:218:LEU:HD12	1:A:220:ASN:H	1.83	0.43
1:B:161:HIS:CD2	1:B:162:THR:HG23	2.54	0.43
1:A:259:PRO:HA	1:A:374:HIS:CD2	2.53	0.42
1:A:299:LEU:O	1:A:303:LEU:HB2	2.19	0.42
2:E:9:DG:N2	6:E:101:HOH:O	2.50	0.42
1:A:24:TRP:HA	1:A:54:VAL:HG13	2.00	0.42
1:B:44:ALA:HB1	1:B:49:VAL:O	2.20	0.41
1:A:384:LYS:HA	1:A:384:LYS:HD3	1.74	0.41
1:A:268:SER:OG	1:A:269:PRO:HD3	2.20	0.41
1:A:76:LEU:HD22	1:A:92:PHE:HE1	1.86	0.41
2:C:8:DC:H2'	2:C:9:DG:C8	2.55	0.41
1:A:379:MET:HE1	2:C:7:TTD:C2T	2.50	0.41
1:B:456:LEU:HD12	1:B:456:LEU:HA	1.95	0.41
2:C:7:TTD:H2''	2:C:7:TTD:H6	1.70	0.41
1:A:343:HIS:HB3	1:A:345:TYR:CE1	2.56	0.41
1:B:171:LEU:HD12	1:B:171:LEU:HA	1.84	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:259:PRO:HA	1:B:374:HIS:CD2	2.56	0.40
1:A:403:ASN:O	1:A:407:GLU:HB2	2.20	0.40
1:B:155:LYS:HD2	1:B:155:LYS:HA	1.73	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:339:ASP:OD2	6:B:601:HOH:O[4_545]	1.95	0.25

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	463/482 (96%)	448 (97%)	15 (3%)	0	100	100
1	B	440/482 (91%)	431 (98%)	9 (2%)	0	100	100
All	All	903/964 (94%)	879 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	383/416 (92%)	368 (96%)	15 (4%)	32	30

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	362/416 (87%)	354 (98%)	8 (2%)	52	55
All	All	745/832 (90%)	722 (97%)	23 (3%)	43	40

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	HIS
1	A	46	GLU
1	A	84	SER
1	A	107	ASP
1	A	108	TYR
1	A	132[A]	SER
1	A	132[B]	SER
1	A	162	THR
1	A	186	VAL
1	A	199	THR
1	A	222	ASP
1	A	241	MET
1	A	323[A]	GLU
1	A	323[B]	GLU
1	A	429	ARG
1	B	1	SER
1	B	3[A]	MET
1	B	3[B]	MET
1	B	84	SER
1	B	294	SER
1	B	359	LEU
1	B	394	LYS
1	B	429	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	161	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	TTD	C	7	2	42,45,46	1.25	6 (14%)	62,74,77	1.76	14 (22%)
2	TTD	E	7	2	42,45,46	1.24	6 (14%)	62,74,77	1.76	12 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TTD	C	7	2	-	10/22/109/110	0/5/6/6
2	TTD	E	7	2	-	10/22/109/110	0/5/6/6

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	7	TTD	C2T-N1T	3.26	1.43	1.36
2	E	7	TTD	C2T-N1T	3.12	1.42	1.36
2	E	7	TTD	C4T-N3T	-2.93	1.32	1.37
2	C	7	TTD	C4T-N3T	-2.85	1.32	1.37
2	E	7	TTD	C2T-N3T	-2.73	1.33	1.38
2	C	7	TTD	C2-N3	-2.71	1.33	1.38
2	E	7	TTD	C2-N3	-2.63	1.33	1.38
2	C	7	TTD	C1R-N1T	2.62	1.49	1.45
2	E	7	TTD	C4-N3	-2.61	1.33	1.37
2	C	7	TTD	C2T-N3T	-2.59	1.33	1.38
2	C	7	TTD	C4-N3	-2.58	1.33	1.37
2	E	7	TTD	C1R-N1T	2.40	1.48	1.45

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	7	TTD	N3T-C2T-N1T	5.23	122.12	116.69
2	E	7	TTD	N3T-C2T-N1T	4.89	121.77	116.69
2	E	7	TTD	C2'-C1'-N1	-4.60	109.37	115.59
2	C	7	TTD	O4R-C1R-N1T	4.32	113.77	108.65
2	E	7	TTD	O4R-C1R-N1T	4.30	113.75	108.65
2	C	7	TTD	C2'-C1'-N1	-4.22	109.89	115.59
2	E	7	TTD	O4-C4-C5	-4.00	119.68	122.88
2	E	7	TTD	C4T-N3T-C2T	-3.96	120.58	126.67
2	E	7	TTD	O2T-C2T-N1T	-3.88	117.47	123.49
2	C	7	TTD	O4-C4-C5	-3.83	119.81	122.88
2	C	7	TTD	O2T-C2T-N1T	-3.70	117.74	123.49
2	C	7	TTD	C4-N3-C2	-3.67	121.03	126.67
2	C	7	TTD	O4T-C4T-C5T	-3.60	120.00	122.88
2	E	7	TTD	O4'-C1'-N1	3.17	112.41	108.65
2	E	7	TTD	C4-N3-C2	-3.16	121.82	126.67
2	C	7	TTD	C4T-N3T-C2T	-3.05	121.97	126.67
2	C	7	TTD	O4'-C1'-N1	2.77	111.94	108.65
2	E	7	TTD	N3-C2-N1	2.75	119.55	116.69
2	C	7	TTD	N3-C2-N1	2.61	119.40	116.69
2	E	7	TTD	O4T-C4T-C5T	-2.57	120.83	122.88
2	E	7	TTD	C5-C4-N3	2.51	118.24	116.06
2	C	7	TTD	C5M-C5T-C5	-2.23	109.94	116.39
2	C	7	TTD	O2-C2-N1	-2.19	120.10	123.49
2	E	7	TTD	O2-C2-N1	-2.11	120.22	123.49
2	C	7	TTD	C6-C6T-N1T	-2.05	110.00	118.20
2	C	7	TTD	C5-C4-N3	2.05	117.84	116.06

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	7	TTD	O4'-C4R-C5'-O5'
2	C	7	TTD	C3R-C4R-C5'-O5'
2	E	7	TTD	C3R-C4R-C5'-O5'
2	C	7	TTD	C2R-C1R-N1T-C6T
2	C	7	TTD	C2'-C1'-N1-C6
2	E	7	TTD	C2'-C1'-N1-C6
2	C	7	TTD	O4R-C4'-C5R-O5R
2	E	7	TTD	O4R-C4'-C5R-O5R
2	C	7	TTD	C3'-C4'-C5R-O5R
2	E	7	TTD	C2R-C1R-N1T-C6T
2	E	7	TTD	C3'-C4'-C5R-O5R
2	C	7	TTD	O4R-C1R-N1T-C6T

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	C	7	TTD	C2'-C1'-N1-C2
2	E	7	TTD	C2'-C1'-N1-C2
2	E	7	TTD	O4'-C4R-C5'-O5'
2	C	7	TTD	O4'-C1'-N1-C6
2	E	7	TTD	O4'-C1'-N1-C6
2	E	7	TTD	O4R-C1R-N1T-C6T
2	C	7	TTD	O4'-C1'-N1-C2
2	E	7	TTD	O4'-C1'-N1-C2

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	7	TTD	4	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	FAD	A	502	-	53,58,58	0.63	0	68,89,89	1.07	6 (8%)
5	FAD	B	501	-	53,58,58	0.46	0	68,89,89	0.52	2 (2%)
4	SO4	A	501	-	4,4,4	0.14	0	6,6,6	0.05	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	FAD	A	502	-	-	6/30/50/50	0/6/6/6
5	FAD	B	501	-	-	5/30/50/50	0/6/6/6

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	502	FAD	C4-C4X-N5	3.68	123.46	118.23
5	A	502	FAD	C9A-C5X-N5	-3.50	118.63	122.43
5	A	502	FAD	P-O3P-PA	-2.59	123.95	132.83
5	B	501	FAD	P-O3P-PA	-2.55	124.09	132.83
5	A	502	FAD	C10-C4X-N5	-2.53	119.49	124.86
5	A	502	FAD	C5A-C6A-N6A	2.32	123.88	120.35
5	B	501	FAD	C5A-C6A-N6A	2.28	123.82	120.35
5	A	502	FAD	C4-N3-C2	-2.08	121.80	125.64

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	502	FAD	C5B-O5B-PA-O1A
5	A	502	FAD	N10-C1'-C2'-O2'
5	A	502	FAD	N10-C1'-C2'-C3'
5	B	501	FAD	C5B-O5B-PA-O1A
5	B	501	FAD	C5B-O5B-PA-O2A
5	B	501	FAD	C5B-O5B-PA-O3P
5	B	501	FAD	N10-C1'-C2'-C3'
5	B	501	FAD	C4'-C5'-O5'-P
5	A	502	FAD	C5B-O5B-PA-O3P
5	A	502	FAD	C4'-C5'-O5'-P
5	A	502	FAD	C5B-O5B-PA-O2A


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	501	FAD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	456/482 (94%)	0.13	2 (0%) 92 92	17, 22, 40, 62	0
1	B	434/482 (90%)	0.04	0 100 100	17, 27, 43, 53	0
2	C	10/14 (71%)	-0.65	0 100 100	18, 38, 47, 59	0
2	E	9/14 (64%)	-0.62	0 100 100	23, 40, 49, 55	0
3	D	14/14 (100%)	-0.60	0 100 100	25, 41, 59, 61	0
3	F	13/14 (92%)	-0.63	0 100 100	32, 49, 58, 59	0
All	All	936/1020 (91%)	0.05	2 (0%) 95 94	17, 25, 46, 62	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	188	PRO	2.8
1	A	359	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	TTD	E	7	40/41	0.90	0.11	18,23,36,42	0
2	TTD	C	7	40/41	0.91	0.12	17,17,22,27	0

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	FAD	A	502	53/53	0.89	0.16	16,17,17,19	0
5	FAD	B	501	53/53	0.90	0.14	18,19,25,30	0
4	SO4	A	501	5/5	0.95	0.10	19,20,27,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.