



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 11, 2021 – 12:15 PM GMT

PDB ID : 6YE8
Title : E.coli's Putrescine receptor PotF complexed with Spermidine
Authors : Shanmugaratnam, S.; Kroeger, P.; Hocker, B.
Deposited on : 2020-03-24
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.16
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.16


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	352	 5% 90% 7%
1	B	352	 5% 89% 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SPD	B	401	-	-	X	-
3	JFN	A	402	X	-	-	-
3	JFN	A	404	X	-	-	-
3	JFN	A	405	X	-	-	-
3	JFN	B	402	X	-	-	-
3	JFN	B	403	X	-	-	-
3	JFN	B	405	X	-	-	-
3	JFN	B	406	X	-	-	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 11846 atoms, of which 5643 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Putrescine-binding periplasmic protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	342	5570	1794	2792	460	513	11	0	19	0
1	B	340	5438	1757	2706	450	517	8	0	11	0

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	371	LEU	-	expression tag	UNP P31133
A	372	GLU	-	expression tag	UNP P31133
A	373	HIS	-	expression tag	UNP P31133
A	374	HIS	-	expression tag	UNP P31133
A	375	HIS	-	expression tag	UNP P31133
A	376	HIS	-	expression tag	UNP P31133
A	377	HIS	-	expression tag	UNP P31133
A	378	HIS	-	expression tag	UNP P31133
B	371	LEU	-	expression tag	UNP P31133
B	372	GLU	-	expression tag	UNP P31133
B	373	HIS	-	expression tag	UNP P31133
B	374	HIS	-	expression tag	UNP P31133
B	375	HIS	-	expression tag	UNP P31133
B	376	HIS	-	expression tag	UNP P31133
B	377	HIS	-	expression tag	UNP P31133
B	378	HIS	-	expression tag	UNP P31133

- Molecule 2 is SPERMIDINE (three-letter code: SPD) (formula: C₇H₁₉N₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	N	0	0
			29	7	19	3		
2	B	1	Total	C	H	N	0	0
			29	7	19	3		

- Molecule 3 is (2R)-1-methoxypropan-2-amine (three-letter code: JFN) (formula: $C_4H_{11}NO$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	0	0
			17	4	11	1	1		
3	A	1	Total	C	H	N	O	0	0
			17	4	11	1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
3	A	1	Total	C	H	N	O	0	0
			17	4	11	1	1		
3	B	1	Total	C	H	N	O	0	0
			17	4	11	1	1		
3	B	1	Total	C	H	N	O	0	0
			17	4	11	1	1		
3	B	1	Total	C	H	N	O	0	0
			17	4	11	1	1		
3	B	1	Total	C	H	N	O	0	0
			17	4	11	1	1		

- Molecule 4 is (2 {S})-1-(2-methoxyethoxy)propan-2-amine (three-letter code: ONT) (formula: C₆H₁₅NO₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
4	A	1	Total	C	H	N	O	0	0
			24	6	15	1	2		
4	B	1	Total	C	H	N	O	0	0
			24	6	15	1	2		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Cl	0	0
			1	1		
5	A	1	Total	Cl	0	0
			1	1		

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	311	Total	O	0	4
			315	315		
7	B	288	Total	O	0	3
			291	291		


3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putrescine-binding periplasmic protein



- Molecule 1: Putrescine-binding periplasmic protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	70.83Å 70.83Å 272.08Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.71 – 1.50 45.55 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.9 (40.71-1.50) 99.9 (45.55-1.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.95 (at 1.50Å)	Xtrriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.162 , 0.183 0.162 , 0.182	Depositor DCC
R_{free} test set	2101 reflections (1.64%)	wwPDB-VP
Wilson B-factor (Å ²)	25.3	Xtrriage
Anisotropy	0.038	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 53.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.019 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	11846	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, SPD, JFN, ONT, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/2907	0.67	0/3952
1	B	0.46	0/2835	0.66	0/3858
All	All	0.47	0/5742	0.67	0/7810

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2778	2792	2794	27	0
1	B	2732	2706	2700	20	0
2	A	10	19	19	5	0
2	B	10	19	19	6	0
3	A	18	33	0	1	0
3	B	24	44	0	2	0
4	A	9	15	0	0	0
4	B	9	15	0	1	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	B	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	315	0	0	7	0
7	B	291	0	0	6	0
All	All	6203	5643	5532	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:185:GLU:OE2	2:A:401:SPD:C9	2.32	0.78
1:A:185:GLU:OE2	2:A:401:SPD:H91	1.87	0.73
1:B:340:ASP:OD1	7:B:501:HOH:O	2.13	0.66
1:A:185:GLU:OE2	2:A:401:SPD:H92	1.96	0.65
1:B:276:PHE:CD1	2:B:401:SPD:H71	2.35	0.62
1:B:114:PRO:O	1:B:118:LYS:HG2	2.00	0.62
3:A:402:JFN:N04	7:A:504:HOH:O	2.31	0.60
1:B:276:PHE:CE1	2:B:401:SPD:H71	2.37	0.59
1:A:303:ASP:OD1	7:A:501:HOH:O	2.17	0.58
1:A:51:LYS:HD3	7:A:508:HOH:O	2.05	0.56
1:B:185:GLU:OE2	2:B:401:SPD:N10	2.41	0.54
1:A:131[B]:MET:CE	1:A:300:LEU:HD11	2.40	0.52
1:B:36:ASN:O	1:B:61:VAL:HA	2.10	0.52
1:A:99:GLN:OE1	1:A:289:LYS:HE3	2.09	0.52
1:A:276:PHE:CD1	2:A:401:SPD:H82	2.45	0.52
1:B:52[A]:GLU:HG3	7:B:512:HOH:O	2.11	0.51
1:A:36:ASN:O	1:A:61:VAL:HA	2.12	0.50
1:A:216:LYS:NZ	7:A:511:HOH:O	2.46	0.48
1:A:33:HIS:HE1	7:A:744:HOH:O	1.96	0.48
1:A:177:VAL:HG22	1:A:240[B]:VAL:CG2	2.43	0.48
1:A:298:TYR:HA	1:A:301[B]:ARG:HD3	1.95	0.47
1:A:276:PHE:CG	2:A:401:SPD:H82	2.50	0.47
1:B:276:PHE:CB	2:B:401:SPD:H92	2.44	0.47
1:A:47:ALA:O	1:A:51:LYS:HG2	2.15	0.46
1:B:274[B]:MET:CE	1:B:348:LEU:HG	2.46	0.46
1:B:185:GLU:OE2	2:B:401:SPD:C9	2.64	0.46
1:B:183:PRO:HB3	1:B:363:TRP:CG	2.51	0.45
3:B:405:JFN:C15	7:B:771:HOH:O	2.64	0.45
1:A:131[B]:MET:HE2	1:A:300:LEU:HD11	1.98	0.45
1:A:297:ASN:O	1:A:301[B]:ARG:HD2	2.17	0.45
1:B:357:ARG:HE	3:B:405:JFN:C08	2.29	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:118:LYS:HD3	7:B:594:HOH:O	2.17	0.44
1:A:301[A]:ARG:HG2	1:A:303:ASP:OD1	2.18	0.43
1:A:177:VAL:HG22	1:A:240[A]:VAL:CG1	2.49	0.43
1:A:289:LYS:HE2	7:A:743:HOH:O	2.19	0.43
1:B:30:LYS:HB2	7:B:539:HOH:O	2.19	0.43
1:B:342:ARG:HD2	4:B:404:ONT:C06	2.48	0.43
1:B:249:TRP:CZ2	1:B:266:PHE:HB3	2.54	0.42
1:A:169:LEU:HD11	1:A:177:VAL:HG11	2.01	0.42
1:B:31[B]:THR:HA	1:B:56:LYS:O	2.20	0.42
1:B:235[B]:ASN:ND2	7:B:504:HOH:O	2.36	0.42
1:A:131[B]:MET:HE1	1:A:300:LEU:CD1	2.50	0.41
1:B:178:SER:O	1:B:241:ALA:HA	2.20	0.41
1:A:107:PRO:O	7:A:502:HOH:O	2.22	0.41
1:B:276:PHE:CG	2:B:401:SPD:H92	2.56	0.41
1:A:47:ALA:O	1:A:51:LYS:HE2	2.21	0.41
1:A:360:THR:O	1:A:364:THR:HG23	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	359/352 (102%)	354 (99%)	5 (1%)	0	100	100
1	B	349/352 (99%)	343 (98%)	6 (2%)	0	100	100
All	All	708/704 (101%)	697 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	308/298 (103%)	306 (99%)	2 (1%)	86	74
1	B	299/298 (100%)	294 (98%)	5 (2%)	60	33
All	All	607/596 (102%)	600 (99%)	7 (1%)	76	48

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	TYR
1	A	91	ARG
1	B	33	HIS
1	B	40	TYR
1	B	328	GLU
1	B	330[A]	ARG
1	B	330[B]	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 14 ligands modelled in this entry, 2 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	JFN	B	405	-	1/1/1/1	-	-
3	JFN	B	402	-	1/1/1/1	-	-
3	JFN	A	402	-	1/1/1/1	-	-
3	JFN	B	403	-	1/1/1/1	-	-
3	JFN	B	406	-	1/1/1/1	-	-
3	JFN	A	404	-	1/1/1/1	-	-
3	JFN	A	405	-	1/1/1/1	-	-

There are no bond length outliers.

There are no bond angle outliers.

All (7) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	B	405	JFN	C08
3	B	402	JFN	C08
3	A	402	JFN	C08
3	B	403	JFN	C08
3	B	406	JFN	C08
3	A	404	JFN	C08
3	A	405	JFN	C08

There are no torsion outliers.


There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	405	JFN	2	0
3	A	402	JFN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	342/352 (97%)	0.07	2 (0%) 89 91	20, 29, 55, 96	0
1	B	340/352 (96%)	0.29	16 (4%) 31 34	19, 29, 58, 84	0
All	All	682/704 (96%)	0.18	18 (2%) 56 61	19, 29, 57, 96	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	31[A]	THR	3.7
1	B	202	LYS	3.6
1	B	194	LEU	3.0
1	B	203	ALA	3.0
1	B	29	GLN	2.8
1	B	72	LEU	2.7
1	B	284	ALA	2.7
1	B	98	PHE	2.7
1	B	93	LEU	2.6
1	B	201	THR	2.5
1	B	82	VAL	2.4
1	A	350[A]	VAL	2.3
1	B	204	ASP	2.3
1	B	355	ILE	2.2
1	B	215	LEU	2.2
1	B	354	LYS	2.2
1	B	75	GLY	2.1
1	B	281	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	JFN	B	403	6/6	0.37	0.25	51,67,86,86	0
3	JFN	B	406	6/6	0.56	0.44	61,73,87,87	0
3	JFN	A	405	6/6	0.60	0.30	65,81,96,96	0
4	ONT	A	403	9/9	0.63	0.18	61,77,106,106	0
3	JFN	B	405	6/6	0.68	0.34	75,90,113,113	0
3	JFN	A	404	6/6	0.72	0.24	71,97,117,117	0
5	CL	A	406	1/1	0.82	0.10	65,65,65,65	0
3	JFN	A	402	6/6	0.84	0.17	55,66,77,77	0
2	SPD	A	401	10/10	0.86	0.18	29,36,72,75	0
4	ONT	B	404	9/9	0.86	0.14	49,59,65,67	0
6	SO4	B	407	5/5	0.91	0.24	102,103,106,117	0
2	SPD	B	401	10/10	0.92	0.14	26,37,61,67	0
3	JFN	B	402	6/6	0.95	0.08	48,58,64,64	0
5	CL	B	408	1/1	0.99	0.06	29,29,29,29	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around SPD A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around SPD B 401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers

There are no such residues in this entry.