



wwPDB X-ray Structure Validation Summary Report

Feb 6, 2024 – 04:37 AM EST

PDB ID : 1YFG
Title : YEAST INITIATOR TRNA
Authors : Basavappa, R.; Sigler, P.B.
Deposited on : 1997-05-08
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2416 (3.00-3.00)
RNA backbone	3102	1173 (3.30-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	75	 • 24% 40% 35%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1639 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called YEAST INITIATOR TRNA.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	A	75	1639	734	298	531	76	0	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: YEAST INITIATOR TRNA



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value
Space group	P 64 2 2
Cell constants a, b, c, α , β , γ	113.72Å 113.72Å 136.54Å 90.00° 90.00° 120.00°
Resolution (Å)	10.00 – 3.00
% Data completeness (in resolution range)	81.7 (10.00-3.00)
R_{merge}	(Not available)
R_{sym}	0.08
Refinement program	NUCLIN/PROFFT, MODIFIED BY Z.OTWINOWSKI, NUCLIN, PROFFT
R, R_{free}	0.215 , (Not available)
Estimated twinning fraction	No twinning to report.
Total number of atoms	1639
Average B, all atoms (Å ²)	57.0

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: H2U, 1MA, RIA, M2G, 7MG, 2MG, T6A, 1MG, 5MC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.77	16/1530 (1.0%)	3.19	245/2380 (10.3%)

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	41	C	O3'-P	-8.59	1.50	1.61
1	A	18	G	O3'-P	7.16	1.69	1.61
1	A	65	G	P-O5'	-6.79	1.52	1.59
1	A	54	A	O3'-P	6.37	1.68	1.61
1	A	21	A	N9-C4	6.29	1.41	1.37

The worst 5 of 245 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	37	T6A	P-O3'-C3'	26.06	150.97	119.70
1	A	38	A	P-O3'-C3'	20.33	144.09	119.70
1	A	76	A	O4'-C1'-N9	20.29	124.44	108.20
1	A	29	G	P-O3'-C3'	17.69	140.92	119.70
1	A	19	G	O4'-C1'-N9	-16.88	94.70	108.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1639	0	848	72	0
All	All	1639	0	848	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 29.

The worst 5 of 72 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:41:C:H2'	1:A:42:U:C6	2.07	0.90
1:A:47:H2U:O2	1:A:47:H2U:H2'	1.75	0.86
1:A:50:U:H5'	1:A:51:C:OP2	1.77	0.83
1:A:59:A:O2'	1:A:60:A:H5'	1.84	0.78
1:A:38:A:H2'	1:A:39:C:H6	1.48	0.77

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	73/75 (97%)	37 (50%)	7 (9%)

5 of 37 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	5	C
1	A	9	1MG
1	A	11	C
1	A	12	G

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	13	C

5 of 7 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	38	A
1	A	39	C
1	A	74	C
1	A	47	H2U
1	A	37	T6A

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

11 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	2MG	A	10	1	18,26,27	1.41	4 (22%)	16,38,41	1.88	1 (6%)
1	5MC	A	49	1	18,22,23	0.74	0	26,32,35	1.27	2 (7%)
1	T6A	A	37	1	27,34,35	1.74	8 (29%)	29,49,52	3.99	10 (34%)
1	1MA	A	58	1	16,25,26	2.48	6 (37%)	18,37,40	2.27	6 (33%)
1	M2G	A	26	1	20,27,28	1.43	5 (25%)	22,40,43	1.53	2 (9%)
1	H2U	A	47	1	18,21,22	1.04	0	21,30,33	2.16	5 (23%)
1	5MC	A	48	1	18,22,23	0.82	1 (5%)	26,32,35	1.18	2 (7%)
1	H2U	A	16	1	18,21,22	0.95	0	21,30,33	1.27	3 (14%)
1	7MG	A	46	1	22,26,27	3.81	2 (9%)	29,39,42	1.79	4 (13%)
1	RIA	A	64	1	31,38,39	1.20	3 (9%)	39,57,60	2.50	11 (28%)
1	1MG	A	9	1	18,26,27	1.01	1 (5%)	19,39,42	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	2MG	A	10	1	-	0/5/27/28	0/3/3/3
1	5MC	A	49	1	-	4/7/25/26	0/2/2/2
1	T6A	A	37	1	-	9/19/41/42	0/3/3/3
1	1MA	A	58	1	-	2/3/25/26	0/3/3/3
1	M2G	A	26	1	-	2/7/29/30	0/3/3/3
1	H2U	A	47	1	-	5/7/38/39	0/2/2/2
1	5MC	A	48	1	-	1/7/25/26	0/2/2/2
1	H2U	A	16	1	-	2/7/38/39	0/2/2/2
1	7MG	A	46	1	-	4/7/37/38	0/3/3/3
1	RIA	A	64	1	-	4/13/51/52	0/4/4/4
1	1MG	A	9	1	-	0/3/25/26	0/3/3/3

The worst 5 of 30 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	46	7MG	C8-N9	-17.28	1.36	1.46
1	A	58	1MA	O4'-C1'	5.86	1.49	1.41
1	A	58	1MA	O2'-C2'	4.02	1.52	1.43
1	A	64	RIA	P'-O3X	-3.67	1.40	1.54
1	A	58	1MA	C2-N3	3.54	1.33	1.29

The worst 5 of 46 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	37	T6A	O14-C14-C15	11.38	143.46	109.74
1	A	64	RIA	O2A-C1'-O1'	-8.58	102.15	111.43
1	A	37	T6A	O14-C14-C12	-8.31	92.45	109.13
1	A	37	T6A	N6-C10-N11	7.95	124.86	113.76
1	A	37	T6A	ODA-C13-C12	-7.04	97.47	121.70

There are no chirality outliers.

5 of 33 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	37	T6A	O4'-C4'-C5'-O5'
1	A	37	T6A	N11-C12-C13-ODA
1	A	37	T6A	N11-C12-C13-ODB
1	A	37	T6A	N11-C12-C14-O14
1	A	37	T6A	C13-C12-C14-O14

There are no ring outliers.

6 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	10	2MG	1	0
1	A	49	5MC	3	0
1	A	58	1MA	1	0
1	A	26	M2G	3	0
1	A	47	H2U	1	0
1	A	9	1MG	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.