



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 26, 2024 – 07:21 AM EDT

PDB ID : 4YGX  
Title : Crystal Structure of D. melanogaster Ssu72+Symplekin bound to cis peptidomimetic CTD phospho-Ser5 peptide  
Authors : Mayfield, J.E.; Zhang, Y.  
Deposited on : 2015-02-26  
Resolution : 2.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1044 (2.98-2.94)
Clashscore	180529	1097 (2.98-2.94)
Ramachandran outliers	177936	1049 (2.98-2.94)
Sidechain outliers	177891	1049 (2.98-2.94)
RSRZ outliers	164620	1044 (2.98-2.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	339	 85% 12% ..
1	C	339	 76% 15% 9%
2	B	200	 72% 20% 5%
2	D	200	 81% 12% ..
3	E	12	 33% 8% 58%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8165 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Symplekin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	333	2581	1618	447	496	20	0	0	0
1	C	309	2398	1505	416	459	18	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	13	GLY	-	expression tag	UNP Q8MSU4
A	14	PRO	-	expression tag	UNP Q8MSU4
A	15	GLY	-	expression tag	UNP Q8MSU4
A	16	SER	-	expression tag	UNP Q8MSU4
A	17	GLY	-	expression tag	UNP Q8MSU4
A	18	MET	-	expression tag	UNP Q8MSU4
C	13	GLY	-	expression tag	UNP Q8MSU4
C	14	PRO	-	expression tag	UNP Q8MSU4
C	15	GLY	-	expression tag	UNP Q8MSU4
C	16	SER	-	expression tag	UNP Q8MSU4
C	17	GLY	-	expression tag	UNP Q8MSU4
C	18	MET	-	expression tag	UNP Q8MSU4

- Molecule 2 is a protein called LD40846p.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	190	1557	977	269	300	11	0	0	0
2	D	192	1577	988	272	305	12	0	1	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-4	GLY	-	expression tag	UNP Q9VWE4
B	-3	PRO	-	expression tag	UNP Q9VWE4
B	-2	GLY	-	expression tag	UNP Q9VWE4
B	-1	SER	-	expression tag	UNP Q9VWE4
B	0	GLY	-	expression tag	UNP Q9VWE4
B	13	ASP	CYS	engineered mutation	UNP Q9VWE4
B	144	ASN	ASP	engineered mutation	UNP Q9VWE4
D	-4	GLY	-	expression tag	UNP Q9VWE4
D	-3	PRO	-	expression tag	UNP Q9VWE4
D	-2	GLY	-	expression tag	UNP Q9VWE4
D	-1	SER	-	expression tag	UNP Q9VWE4
D	0	GLY	-	expression tag	UNP Q9VWE4
D	13	ASP	CYS	engineered mutation	UNP Q9VWE4
D	144	ASN	ASP	engineered mutation	UNP Q9VWE4

- Molecule 3 is a protein called cis peptidomimetic CTD phospho-Ser5 peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	E	5	42	24	5	12	1	0	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	3	3	3	0	0
4	B	2	2	2	0	0
4	C	2	2	2	0	0
4	D	3	3	3	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Symplekin





- Molecule 1: Symplekin




- Molecule 2: LD40846p



- Molecule 2: LD40846p



- Molecule 3: cis peptidomimetic CTD phospho-Ser5 peptide



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 4	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	127.88Å 127.88Å 105.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.95 50.00 – 2.95	Depositor EDS
% Data completeness (in resolution range)	99.2 (50.00-2.95) 99.2 (50.00-2.95)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.77 (at 2.96Å)	Xtrriage
Refinement program	REFMAC 5.8.0073	Depositor
R, $R_{free}$	0.206 , 0.254 0.204 , 0.248	Depositor DCC
$R_{free}$ test set	1843 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	73.8	Xtrriage
Anisotropy	0.102	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 30.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.032 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8165	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	88.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.93% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 4CG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.67	0/2610	0.88	2/3522 (0.1%)
1	C	0.61	0/2425	0.85	2/3269 (0.1%)
2	B	0.68	1/1582 (0.1%)	0.90	2/2125 (0.1%)
2	D	0.62	0/1603	0.91	2/2154 (0.1%)
3	E	0.68	0/25	0.57	0/32
All	All	0.65	1/8245 (0.0%)	0.88	8/11102 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
3	E	0	1
All	All	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	15	SER	CB-OG	6.51	1.50	1.42

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	184	ARG	NE-CZ-NH1	6.71	123.65	120.30
2	B	19	ARG	NE-CZ-NH2	-6.61	117.00	120.30
2	B	15	SER	N-CA-CB	-6.06	101.41	110.50
2	D	88	ARG	NE-CZ-NH1	5.86	123.23	120.30
1	A	303	ARG	NE-CZ-NH1	5.84	123.22	120.30

*Continued on next page...*



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	174	ARG	NE-CZ-NH1	5.47	123.03	120.30
1	C	68	LEU	CA-CB-CG	5.44	127.82	115.30
1	A	20	ASP	CB-CG-OD1	5.03	122.83	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	112	LEU	Mainchain
3	E	5	4CG	Mainchain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2581	0	2653	23	0
1	C	2398	0	2470	31	0
2	B	1557	0	1530	26	0
2	D	1577	0	1546	25	0
3	E	42	0	34	1	0
4	A	3	0	0	0	0
4	B	2	0	0	0	0
4	C	2	0	0	0	0
4	D	3	0	0	0	0
All	All	8165	0	8233	104	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (104) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:196:LEU:HB3	1:C:257:MET:CE	2.11	0.80
1:C:196:LEU:HB3	1:C:257:MET:HE1	1.69	0.75
2:D:163:MET:HE1	2:D:177:ILE:HD11	1.70	0.72

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:141[B]:ASP:O	2:D:186:ARG:NH2	2.25	0.69
1:C:196:LEU:HB3	1:C:257:MET:HE2	1.79	0.63
2:B:159:MET:O	2:B:163:MET:HG3	1.98	0.63
1:C:313:ARG:HD2	1:C:313:ARG:C	2.20	0.61
1:A:262:MET:HE1	1:A:299:LEU:HB3	1.83	0.60
2:D:141[A]:ASP:O	2:D:186:ARG:NH2	2.35	0.58
1:A:276:LEU:HD13	1:A:285:VAL:HG13	1.85	0.57
2:B:70:GLU:OE2	2:B:74:LYS:HB2	2.04	0.57
1:A:262:MET:HE3	1:A:299:LEU:HD13	1.87	0.57
2:B:141:ASP:O	2:B:186:ARG:NH2	2.37	0.57
2:D:88:ARG:HG2	2:D:88:ARG:HH11	1.69	0.57
2:B:137:VAL:HG21	2:B:193:LEU:HD12	1.85	0.57
1:A:198:ARG:CZ	1:A:257:MET:HE3	2.36	0.56
2:D:62:TYR:CE1	2:D:96:PRO:HD3	2.41	0.54
1:C:57:LEU:HD13	1:C:92:GLN:CG	2.37	0.53
2:D:88:ARG:NH1	2:D:89:ASN:OD1	2.41	0.53
2:B:10:ALA:HB2	2:B:106:PHE:CE1	2.45	0.52
2:D:113:GLU:HA	2:D:141[A]:ASP:OD1	2.09	0.52
1:A:42:PRO:HB3	1:A:79:ASN:HD22	1.74	0.52
2:B:69:LEU:HB3	2:B:77:TYR:CD2	2.45	0.52
2:B:75:GLU:HA	2:B:75:GLU:OE1	2.11	0.51
1:C:281:THR:HG22	1:C:284:GLN:CG	2.40	0.51
1:A:255:ALA:HB2	1:A:265:VAL:HG21	1.92	0.51
1:C:42:PRO:HA	1:C:45:LYS:HD3	1.93	0.51
2:B:15:SER:HB2	2:B:17:MET:HB2	1.92	0.50
2:B:69:LEU:HD22	2:B:77:TYR:CE2	2.47	0.50
1:C:57:LEU:HD22	1:C:96:VAL:HG21	1.93	0.50
2:D:55:VAL:C	2:D:56:TYR:CD1	2.85	0.50
2:B:98:ARG:HD3	2:B:101:ASP:OD2	2.13	0.49
2:B:87:ASP:HA	2:B:90:ARG:NE	2.27	0.49
1:C:294:MET:SD	2:D:121:VAL:HG12	2.53	0.49
2:D:54:ASN:HB3	2:D:56:TYR:HE1	1.78	0.49
2:D:137:VAL:HG21	2:D:193:LEU:CD1	2.42	0.49
1:C:83:ARG:O	1:C:87:VAL:HG23	2.13	0.48
2:D:184:ARG:O	2:D:186:ARG:HG3	2.12	0.48
1:A:48:LEU:O	1:A:52:VAL:HG23	2.13	0.48
1:A:194:ASP:CG	1:A:341:ARG:HH12	2.17	0.48
2:B:137:VAL:CG2	2:B:193:LEU:HD12	2.43	0.48
2:B:117:TYR:OH	2:B:191:SER:HB2	2.14	0.47
2:B:137:VAL:HG21	2:B:193:LEU:CD1	2.44	0.47
2:D:56:TYR:CD1	2:D:56:TYR:N	2.83	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:137:VAL:CG2	2:D:193:LEU:HD12	2.44	0.47
1:C:114:ASP:OD1	1:C:115:ASN:N	2.47	0.47
2:B:177:ILE:O	2:B:181:GLU:HG3	2.14	0.47
1:C:57:LEU:HD13	1:C:92:GLN:HB3	1.96	0.47
2:B:124:MET:HB2	2:B:193:LEU:HD22	1.96	0.47
1:C:281:THR:HG22	1:C:284:GLN:CD	2.35	0.47
1:C:238:HIS:HA	1:C:279:THR:HG21	1.97	0.47
2:B:10:ALA:HB2	2:B:106:PHE:CZ	2.50	0.47
2:D:14:SER:HB3	3:E:5:4CG:O06	2.15	0.46
1:C:150:GLN:OE1	1:C:150:GLN:HA	2.15	0.46
2:D:137:VAL:HG21	2:D:193:LEU:HD12	1.98	0.45
1:C:162:ILE:HG21	1:C:180:PHE:CD2	2.51	0.45
1:A:276:LEU:CD1	1:A:285:VAL:HG13	2.45	0.45
2:B:99:PHE:CE2	2:B:120:VAL:HG13	2.52	0.45
1:A:262:MET:CE	1:A:299:LEU:HB3	2.46	0.45
2:B:111:THR:HG21	2:B:117:TYR:HA	2.00	0.44
1:C:277:PRO:HA	1:C:278:PRO:HD2	1.85	0.44
1:A:72:LEU:HD22	1:A:107:VAL:HG21	1.99	0.44
1:A:182:GLU:OE1	1:A:250:SER:HA	2.18	0.44
1:A:183:GLY:O	1:A:187:LEU:HG	2.18	0.44
2:D:20:SER:HB2	2:D:36:SER:HB2	1.99	0.44
2:D:163:MET:CE	2:D:177:ILE:HD11	2.45	0.44
2:B:60:THR:O	2:B:96:PRO:HD2	2.18	0.44
1:A:119:VAL:O	1:A:123:VAL:HG23	2.18	0.43
2:D:124:MET:HE2	2:D:124:MET:HB3	1.95	0.43
2:B:42:ARG:CZ	2:B:57:GLU:OE2	2.66	0.43
1:A:27:ARG:O	1:A:31:VAL:HG23	2.18	0.43
1:C:101:LEU:HB3	1:C:102:PRO:HD3	2.00	0.43
1:A:83:ARG:NH1	1:A:114:ASP:OD2	2.52	0.43
1:A:227:LEU:CD2	1:A:261:PHE:CD1	3.02	0.43
1:A:198:ARG:NH1	1:A:257:MET:HE3	2.34	0.43
1:C:57:LEU:HD13	1:C:92:GLN:HG3	2.00	0.43
1:C:68:LEU:HG	1:C:69:GLU:H	1.83	0.43
1:C:333:MET:HE2	1:C:338:MET:CG	2.50	0.42
1:A:86:VAL:O	1:A:90:VAL:HG23	2.19	0.42
1:C:316:LEU:HD23	1:C:316:LEU:HA	1.86	0.42
2:D:17:MET:HE1	2:D:86:LEU:HD21	2.01	0.42
1:A:94:CYS:O	1:A:98:VAL:HG23	2.19	0.42
2:B:131:ASP:O	2:B:132:ASN:HB2	2.19	0.42
2:B:184:ARG:O	2:B:186:ARG:HG3	2.20	0.42
1:C:270:LYS:HA	1:C:315:MET:SD	2.59	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:16:ASN:HD22	2:D:16:ASN:HA	1.70	0.42
1:A:281:THR:O	1:A:285:VAL:HG23	2.19	0.42
1:C:189:SER:O	1:C:203:SER:HB2	2.20	0.41
1:A:117:ALA:O	1:A:121:LYS:HG3	2.21	0.41
2:B:39:THR:O	2:B:100:GLN:NE2	2.49	0.41
2:B:117:TYR:O	2:B:121:VAL:HG23	2.20	0.41
2:D:61:LYS:HA	2:D:95:CYS:HB3	2.02	0.41
2:D:60:THR:O	2:D:96:PRO:HD2	2.20	0.41
1:A:105:ILE:HG23	1:A:154:ILE:HD12	2.03	0.41
1:C:196:LEU:CB	1:C:257:MET:HE2	2.48	0.41
2:D:173:ILE:O	2:D:177:ILE:HG12	2.21	0.41
1:C:63:LEU:HD13	1:C:66:GLU:OE1	2.21	0.41
1:C:86:VAL:O	1:C:90:VAL:HG23	2.21	0.41
1:C:105:ILE:HD11	1:C:155:LEU:HD23	2.03	0.40
1:C:120:ILE:O	1:C:124:ILE:HG13	2.21	0.40
1:C:324:ASN:O	1:C:328:LYS:HG3	2.20	0.40
1:C:48:LEU:O	1:C:52:VAL:HG23	2.21	0.40
2:D:82:LEU:O	2:D:86:LEU:HD12	2.21	0.40
2:B:112:VAL:HA	2:B:140:VAL:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	331/339 (98%)	322 (97%)	9 (3%)	0	100	100
1	C	307/339 (91%)	299 (97%)	8 (3%)	0	100	100
2	B	186/200 (93%)	179 (96%)	7 (4%)	0	100	100
2	D	191/200 (96%)	184 (96%)	7 (4%)	0	100	100
3	E	2/12 (17%)	2 (100%)	0	0	100	100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	1017/1090 (93%)	986 (97%)	31 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	293/296 (99%)	285 (97%)	8 (3%)	40	64
1	C	273/296 (92%)	265 (97%)	8 (3%)	37	61
2	B	175/181 (97%)	167 (95%)	8 (5%)	23	47
2	D	177/181 (98%)	172 (97%)	5 (3%)	38	63
3	E	4/9 (44%)	4 (100%)	0	100	100
All	All	922/963 (96%)	893 (97%)	29 (3%)	35	59

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	98	VAL
1	A	129	SER
1	A	164	ASP
1	A	210	HIS
1	A	281	THR
1	A	282	ASP
1	A	283	SER
1	A	303	ARG
2	B	16	ASN
2	B	68	ASP
2	B	74	LYS
2	B	86	LEU
2	B	124	MET
2	B	141	ASP
2	B	167	THR
2	B	175	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	46	CYS
1	C	68	LEU
1	C	109	SER
1	C	143	GLU
1	C	260	ILE
1	C	298	THR
1	C	311	THR
1	C	346	LEU
2	D	16	ASN
2	D	56	TYR
2	D	115	ARG
2	D	124	MET
2	D	147	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	4CG	E	5	3	14,16,17	1.67	3 (21%)	11,22,24	2.15	3 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	4CG	E	5	3	-	3/9/22/24	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	5	4CG	C12-C11	3.81	1.67	1.52
3	E	5	4CG	C13-C14	-2.67	1.48	1.55
3	E	5	4CG	C09-C10	2.27	1.37	1.32

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	5	4CG	C12-C11-C10	-4.80	96.89	104.54
3	E	5	4CG	O08-P05-O04	2.99	114.46	106.67
3	E	5	4CG	C12-C13-C14	2.33	107.66	103.82

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	E	5	4CG	C03-O04-P05-O07
3	E	5	4CG	O04-C03-CA-N
3	E	5	4CG	C10-C09-CA-C03

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	5	4CG	1	0

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	333/339 (98%)	-0.23	5 (1%) 71 69	41, 71, 144, 180	0
1	C	309/339 (91%)	0.02	14 (4%) 39 36	49, 85, 171, 207	0
2	B	190/200 (95%)	-0.03	6 (3%) 50 47	48, 79, 159, 179	0
2	D	192/200 (96%)	-0.32	2 (1%) 79 77	31, 80, 117, 148	1 (0%)
3	E	4/12 (33%)	0.73	0 100 100	101, 120, 129, 138	0
All	All	1028/1090 (94%)	-0.13	27 (2%) 57 54	31, 79, 154, 207	1 (0%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	4	PRO	5.8
2	D	4	PRO	4.9
1	C	48	LEU	4.4
1	C	42	PRO	3.7
1	C	89	PHE	3.2
1	A	350	ALA	3.0
1	A	37	LEU	2.9
1	C	64	ALA	2.9
1	C	71	VAL	2.8
2	B	5	SER	2.8
1	C	68	LEU	2.7
1	C	74	LEU	2.7
2	B	46	PRO	2.7
1	C	67	PHE	2.6
1	C	61	ALA	2.5
1	C	138	LEU	2.5
2	D	56	TYR	2.4
1	C	72	LEU	2.4
1	C	86	VAL	2.4
2	B	15	SER	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	38	VAL	2.3
2	B	78	THR	2.3
1	A	30	VAL	2.2
1	C	350	ALA	2.2
1	A	31	VAL	2.1
1	C	141	LEU	2.1
2	B	152	GLY	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	4CG	E	5	16/17	0.97	0.07	80,86,98,103	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.