

# Full wwPDB X-ray Structure Validation Report (i)

#### Jun 12, 2024 – 02:26 AM EDT

PDB ID	:	1YHK
Title	:	Trypanosoma cruzi farnesyl diphosphate synthase
Authors	:	Gabelli, S.B.; McLellan, J.S.; Montalvetti, A.; Oldfield, E.; Docampo, R.;
		Amzel, L.M.
Deposited on	:	2005-01-09
Resolution	:	2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain		
1	А	362	81%	16%	•••



#### 1YHK

# 2 Entry composition (i)


There are 3 unique types of molecules in this entry. The entry contains 3163 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called farmesyl pyrophosphate synthase.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	А	360	Total 2873	C 1845	N 469	O 536	S 23	0	0	0

• Molecule 2 is SULFATE ION (three-letter code: SO4) (formula:  $O_4S$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
2	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	280	Total         O           280         280	0	0




# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: farnesyl pyrophosphate synthase





## 4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	P 61 2 2	Depositor	
Cell constants	57.92Å 57.92Å 397.70Å	Depositor	
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $120.00^{\circ}$	Depositor	
Resolution (Å)	65.94 - 2.10	Depositor	
% Data completeness	94.0 (65.94-2.10)	Depositor	
(in resolution range)	34.0 (05.34-2.10)	Depositor	
$R_{merge}$	(Not available)	Depositor	
R <sub>sym</sub>	(Not available)	Depositor	
Refinement program	REFMAC 5.1.24	Depositor	
$R, R_{free}$	0.173 , $0.227$	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	3163	wwPDB-VP	
Average B, all atoms $(Å^2)$	31.0	wwPDB-VP	



# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section:  $\mathrm{SO4}$ 

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	1.42	20/2938~(0.7%)	1.15	19/3982~(0.5%)	

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	Ideal(Å)
1	А	231	GLU	CD-OE1	12.11	1.39	1.25
1	А	186	GLN	CG-CD	8.26	1.70	1.51
1	А	171	VAL	CB-CG2	-7.16	1.37	1.52
1	А	264	LYS	CD-CE	6.90	1.68	1.51
1	А	11	TYR	CE1-CZ	6.57	1.47	1.38
1	А	5	GLU	CG-CD	6.32	1.61	1.51
1	А	264	LYS	CE-NZ	6.29	1.64	1.49
1	А	157	GLN	CB-CG	-6.20	1.35	1.52
1	А	343	SER	CB-OG	5.98	1.50	1.42
1	А	295	ALA	CA-CB	5.88	1.64	1.52
1	А	67	GLN	CG-CD	5.78	1.64	1.51
1	А	225	GLU	CD-OE1	5.71	1.31	1.25
1	А	348	GLU	CD-OE1	5.54	1.31	1.25
1	А	114	TYR	CE1-CZ	5.52	1.45	1.38
1	А	158	LYS	CD-CE	5.44	1.64	1.51
1	А	220	GLY	CA-C	5.34	1.60	1.51
1	А	288	ALA	CA-CB	5.34	1.63	1.52
1	А	236	GLU	CD-OE2	5.27	1.31	1.25
1	А	168	MET	SD-CE	-5.17	1.49	1.77
1	А	26	TYR	CE2-CZ	-5.02	1.32	1.38

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	107	ARG	NE-CZ-NH2	-10.41	115.10	120.30

Continued on next page...



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	181	ASP	CB-CG-OD1	9.10	126.49	118.30
1	А	107	ARG	NE-CZ-NH1	8.11	124.35	120.30
1	А	219	MET	CG-SD-CE	7.51	112.21	100.20
1	А	251	ASP	CB-CG-OD1	7.31	124.88	118.30
1	А	271	ASP	CB-CG-OD2	7.29	124.86	118.30
1	А	20	ASP	CB-CG-OD2	7.03	124.63	118.30
1	А	99	ASP	CB-CG-OD2	7.02	124.61	118.30
1	А	69	ASP	CB-CG-OD2	7.01	124.61	118.30
1	А	264	LYS	CD-CE-NZ	6.50	126.64	111.70
1	А	106	MET	CG-SD-CE	-6.38	89.99	100.20
1	А	138	MET	CG-SD-CE	-6.28	90.16	100.20
1	А	51	ARG	NE-CZ-NH1	5.96	123.28	120.30
1	А	191	ASP	CB-CG-OD2	5.58	123.32	118.30
1	А	115	ARG	NE-CZ-NH2	-5.54	117.53	120.30
1	А	108	ARG	NE-CZ-NH2	-5.50	117.55	120.30
1	А	95	LEU	CB-CG-CD1	5.48	120.32	111.00
1	А	107	ARG	CG-CD-NE	-5.43	100.40	111.80
1	А	320	ASP	CB-CG-OD2	5.34	123.11	118.30

Continued from previous page...

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2873	0	2830	24	0
2	А	10	0	0	0	0
3	А	280	0	0	9	1
All	All	3163	0	2830	24	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:106:MET:CE	1:A:106:MET:SD	2.02	1.46
1:A:175:CYS:HB3	3:A:618:HOH:O	1.51	1.07
1:A:145:ASP:HB3	3:A:643:HOH:O	1.79	0.83
1:A:180:LEU:HD22	3:A:579:HOH:O	1.81	0.80
1:A:106:MET:CE	1:A:106:MET:CG	2.69	0.71
1:A:204:VAL:HG22	1:A:247:GLN:HG2	1.79	0.64
1:A:249:GLN:OE1	3:A:649:HOH:O	2.18	0.55
1:A:226:ALA:O	1:A:229:SER:HB3	2.07	0.54
1:A:203:ILE:HG13	3:A:622:HOH:O	2.07	0.54
1:A:197:PRO:O	1:A:201:LYS:HG3	2.08	0.53
1:A:203:ILE:O	1:A:207:LYS:HB3	2.08	0.52
1:A:67:GLN:HB3	3:A:632:HOH:O	2.11	0.51
1:A:93:HIS:NE2	1:A:132:LYS:HE3	2.27	0.49
1:A:25:LYS:HE2	3:A:633:HOH:O	2.12	0.49
1:A:115:ARG:NH2	3:A:464:HOH:O	2.35	0.48
1:A:93:HIS:CD2	1:A:132:LYS:HE3	2.49	0.48
1:A:1:MET:HG3	1:A:3:SER:HB2	1.96	0.47
1:A:318:GLN:CG	3:A:628:HOH:O	2.62	0.47
1:A:63:LEU:HD11	1:A:77:LEU:HG	1.99	0.43
1:A:99:ASP:OD2	1:A:107:ARG:HD2	2.18	0.43
1:A:272:ALA:HA	1:A:297:TYR:CE2	2.54	0.42
1:A:265:VAL:HG13	1:A:267:THR:HG23	2.02	0.42
1:A:35:TYR:C	1:A:35:TYR:CD1	2.92	0.41
1:A:245:TYR:CE2	1:A:325:GLU:HG3	2.55	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)	
3:A:495:HOH:O	3:A:647:HOH:O[1_655]	2.11	0.09	

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	А	358/362~(99%)	350~(98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	307/309~(99%)	300~(98%)	7~(2%)	50 55	

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	5	GLU
1	А	95	LEU
1	А	107	ARG
1	А	115	ARG
1	А	238	VAL
1	А	273	LYS
1	А	358	HIS

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such side chains are listed below:

Mol	Chain	Res	Type
1	А	316	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type	Trune	Chain	Dec	Link	Bond lengths Bon			ond ang	gles	
	туре	Ullalli	$\operatorname{Res}$		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z >2
2	SO4	А	370	-	4,4,4	0.48	0	$6,\!6,\!6$	0.95	0
2	SO4	А	371	-	4,4,4	0.76	0	$6,\!6,\!6$	0.41	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

#### 5.7 Other polymers (i)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

#### 6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

## 6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

#### 6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

