



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 28, 2024 – 10:09 AM JST

PDB ID : 8YIV
Title : N17.1.2 recognition of NRAS neoantigens
Authors : Wu, D.C.; Mariuzza, R.A.
Deposited on : 2024-02-29
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.21
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.004 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	6234 (2.10-2.10)
Clashscore	180529	6893 (2.10-2.10)
Ramachandran outliers	177936	6839 (2.10-2.10)
Sidechain outliers	177891	6840 (2.10-2.10)
RSRZ outliers	164620	6234 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	 6% (red), 81% (green), 17% (yellow), 0% (orange), 0% (grey)
2	C	10	 0% (red), 80% (green), 20% (yellow), 0% (orange), 0% (grey)
3	B	100	 4% (red), 86% (green), 14% (yellow), 0% (orange), 0% (grey)
4	D	207	 2% (red), 79% (green), 16% (yellow), 0% (orange), 0% (grey)
5	E	247	 0% (red), 87% (green), 13% (yellow), 0% (orange), 0% (grey)

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	GOL	E	305	-	-	X	-
7	EDO	E	301	-	-	X	-

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 7286 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MHC class I antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	275	2227	1384	408	425	10	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP F6IQR9

- Molecule 2 is a protein called ILE-LEU-ASP-THR-ALA-GLY-LYS-GLU-GLU-TYR.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	10	80	50	11	19	0	0	0

- Molecule 3 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	B	100	824	525	139	156	4	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	expression tag	UNP P61769


- Molecule 4 is a protein called TCR ALPHA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	202	1567	990	258	310	9	0	0	0

- Molecule 5 is a protein called TCR beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	E	247	1985	1256	341	379	9	0	0	0

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
6	A	1	6	3	3	0	0
6	A	1	6	3	3	0	0
6	A	1	6	3	3	0	0
6	D	1	6	3	3	0	0
6	E	1	6	3	3	0	0
6	E	1	6	3	3	0	0
6	E	1	6	3	3	0	0
6	E	1	6	3	3	0	0

- Molecule 7 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C O 4 2 2	0	0
7	E	1	Total C O 4 2 2	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	139	Total O 139 139	0	0
8	C	10	Total O 10 10	0	0
8	B	45	Total O 45 45	0	0
8	D	129	Total O 129 129	0	0
8	E	224	Total O 224 224	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MHC class I antigen




- Molecule 2: ILE-LEU-ASP-THR-ALA-GLY-LYS-GLU-GLU-TYR




- Molecule 3: Beta-2-microglobulin



- Molecule 4: TCR ALPHA



- Molecule 5: TCR beta

Chain E:  87% 13%



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	71.14Å 133.22Å 222.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.69 – 2.10 42.69 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.7 (42.69-2.10) 99.7 (42.69-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.93 (at 2.10Å)	Xtrriage
Refinement program	PHENIX (1.16_3549: ???)	Depositor
R, R_{free}	0.191 , 0.226 0.196 , 0.228	Depositor DCC
R_{free} test set	3124 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	33.9	Xtrriage
Anisotropy	0.285	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 39.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7286	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.55	2/2287 (0.1%)	0.79	4/3100 (0.1%)
2	C	0.58	0/80	0.66	0/105
3	B	0.46	0/847	0.62	0/1149
4	D	0.55	0/1601	0.71	1/2171 (0.0%)
5	E	0.59	0/2039	0.67	0/2777
All	All	0.55	2/6854 (0.0%)	0.72	5/9302 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	108	ARG	CZ-NH2	6.74	1.41	1.33
1	A	108	ARG	NE-CZ	-6.36	1.24	1.33

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	108	ARG	NE-CZ-NH2	-15.28	112.66	120.30
1	A	226	GLN	CB-CA-C	-11.84	86.73	110.40
1	A	226	GLN	N-CA-C	11.22	141.30	111.00
1	A	108	ARG	NE-CZ-NH1	-9.06	115.77	120.30
4	D	93	LEU	CA-CB-CG	5.68	128.37	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	108	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2227	0	2082	44	1
2	C	80	0	77	1	0
3	B	824	0	777	11	0
4	D	1567	0	1503	37	0
5	E	1985	0	1919	33	1
6	A	18	0	24	2	0
6	D	6	0	8	1	0
6	E	24	0	32	11	0
7	A	4	0	6	0	0
7	E	4	0	6	11	0
8	A	139	0	0	2	0
8	B	45	0	0	3	0
8	C	10	0	0	0	0
8	D	129	0	0	5	0
8	E	224	0	0	6	0
All	All	7286	0	6434	122	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 122 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:E:228:ASP:H	6:E:305:GOL:H31	1.32	0.94
5:E:240:GLU:H	7:E:301:EDO:H22	1.36	0.89
1:A:84:TYR:HA	6:A:302:GOL:H12	1.56	0.88
1:A:231:VAL:O	1:A:243:LYS:NZ	2.09	0.86
1:A:195:SER:OG	1:A:198:GLU:N	2.11	0.83

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1:GLY:N	5:E:246:ASP:OXT[7_554]	2.05	0.15

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/276 (99%)	262 (96%)	11 (4%)	0	100	100
2	C	8/10 (80%)	7 (88%)	1 (12%)	0	100	100
3	B	98/100 (98%)	96 (98%)	2 (2%)	0	100	100
4	D	200/207 (97%)	189 (94%)	11 (6%)	0	100	100
5	E	245/247 (99%)	242 (99%)	3 (1%)	0	100	100
All	All	824/840 (98%)	796 (97%)	28 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	229/233 (98%)	222 (97%)	7 (3%)	35	39
2	C	8/8 (100%)	8 (100%)	0	100	100
3	B	92/95 (97%)	91 (99%)	1 (1%)	70	77

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	D	174/184 (95%)	169 (97%)	5 (3%)	37	41
5	E	222/222 (100%)	218 (98%)	4 (2%)	54	61
All	All	725/742 (98%)	708 (98%)	17 (2%)	45	51

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
5	E	102	LYS
5	E	195	ARG
3	B	70	PHE
4	D	50	ARG
4	D	74	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	141	GLN
1	A	218	GLN
4	D	187	ASN
4	D	190	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	EDO	A	304	-	3,3,3	0.62	0	2,2,2	0.14	0
6	GOL	E	303	-	5,5,5	1.22	0	5,5,5	0.77	0
6	GOL	E	305	-	5,5,5	0.69	0	5,5,5	0.98	0
6	GOL	A	302	-	5,5,5	0.89	0	5,5,5	0.93	0
6	GOL	A	301	-	5,5,5	0.93	0	5,5,5	1.21	1 (20%)
6	GOL	E	304	-	5,5,5	0.34	0	5,5,5	0.66	0
6	GOL	A	303	-	5,5,5	0.18	0	5,5,5	0.52	0
6	GOL	E	302	-	5,5,5	0.13	0	5,5,5	0.30	0
6	GOL	D	301	-	5,5,5	1.03	0	5,5,5	1.23	1 (20%)
7	EDO	E	301	-	3,3,3	0.58	0	2,2,2	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	EDO	A	304	-	-	0/1/1/1	-
6	GOL	E	303	-	-	2/4/4/4	-
6	GOL	E	305	-	-	0/4/4/4	-
6	GOL	A	302	-	-	0/4/4/4	-
6	GOL	A	301	-	-	0/4/4/4	-
6	GOL	E	304	-	-	2/4/4/4	-
6	GOL	A	303	-	-	3/4/4/4	-
6	GOL	E	302	-	-	4/4/4/4	-
6	GOL	D	301	-	-	0/4/4/4	-
7	EDO	E	301	-	-	1/1/1/1	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	D	301	GOL	C3-C2-C1	-2.52	101.90	111.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	301	GOL	C3-C2-C1	-2.03	103.82	111.70

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	303	GOL	O1-C1-C2-C3
6	E	302	GOL	C1-C2-C3-O3
6	E	303	GOL	O1-C1-C2-C3
6	A	303	GOL	O1-C1-C2-O2
6	E	302	GOL	O2-C2-C3-O3

There are no ring outliers.

6 monomers are involved in 25 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	E	303	GOL	1	0
6	E	305	GOL	9	0
6	A	302	GOL	2	0
6	E	304	GOL	1	0
6	D	301	GOL	1	0
7	E	301	EDO	11	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/276 (99%)	0.40	16 (5%) 30 32	25, 40, 69, 107	0
2	C	10/10 (100%)	-0.66	0 100 100	26, 28, 30, 31	0
3	B	100/100 (100%)	0.42	4 (4%) 43 45	28, 44, 65, 69	0
4	D	202/207 (97%)	0.32	5 (2%) 58 60	27, 38, 60, 75	0
5	E	247/247 (100%)	-0.31	1 (0%) 89 90	22, 29, 48, 119	0
All	All	834/840 (99%)	0.16	26 (3%) 51 53	22, 36, 63, 119	0

The worst 5 of 26 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	17	ARG	6.2
4	D	130	SER	4.4
1	A	227	ASP	3.8
1	A	222	GLU	3.7
1	A	16	GLY	3.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	GOL	E	302	6/6	0.59	0.25	46,49,54,57	0
6	GOL	E	304	6/6	0.67	0.20	38,47,55,56	0
6	GOL	E	303	6/6	0.72	0.17	44,44,52,57	0
6	GOL	E	305	6/6	0.73	0.13	55,56,59,59	0
6	GOL	A	303	6/6	0.77	0.15	60,65,77,77	0
6	GOL	A	302	6/6	0.78	0.12	44,52,59,61	0
6	GOL	A	301	6/6	0.80	0.15	45,51,54,59	0
7	EDO	A	304	4/4	0.81	0.16	49,50,50,54	0
7	EDO	E	301	4/4	0.83	0.20	43,44,48,53	0
6	GOL	D	301	6/6	0.96	0.07	32,34,37,37	0

6.5 Other polymers [i](#)

There are no such residues in this entry.