



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 04:45 pm BST

PDB ID : 2YJ1
Title : Puma BH3 foldamer in complex with Bcl-xL
Authors : Lee, E.F.; Smith, B.J.; Horne, W.S.; Mayer, K.N.; Evangelista, M.; Colman, P.M.; Gellman, S.H.; Fairlie, W.D.
Deposited on : 2011-05-18
Resolution : 2.24 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2391 (2.26-2.22)
Clashscore	141614	2539 (2.26-2.22)
Ramachandran outliers	138981	2489 (2.26-2.22)
Sidechain outliers	138945	2490 (2.26-2.22)
RSRZ outliers	127900	2353 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	158	<div style="display: flex; align-items: center;"> <div style="width: 5%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 77%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 16%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 2%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 2%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 5px;">% 77% 16% • •</p>
1	C	158	<div style="display: flex; align-items: center;"> <div style="width: 4%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 73%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 13%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 5%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 11%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 5px;">4% 73% 13% • 11%</p>
2	B	23	<div style="display: flex; align-items: center;"> <div style="width: 57%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 30%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 2%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 5px;">57% 30% 9% •</p>
2	D	23	<div style="display: flex; align-items: center;"> <div style="width: 4%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 57%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 26%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 13%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 2%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 5px;">4% 57% 26% 13% •</p>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2761 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BCL-2-LIKE PROTEIN 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	151	1201	763	204	229	5	0	0	0
1	C	140	1128	724	191	208	5	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	cloning artifact	UNP Q07817
A	-3	PRO	-	cloning artifact	UNP Q07817
A	-2	LEU	-	cloning artifact	UNP Q07817
A	-1	GLY	-	cloning artifact	UNP Q07817
A	0	SER	-	cloning artifact	UNP Q07817
C	-4	GLY	-	cloning artifact	UNP Q07817
C	-3	PRO	-	cloning artifact	UNP Q07817
C	-2	LEU	-	cloning artifact	UNP Q07817
C	-1	GLY	-	cloning artifact	UNP Q07817
C	0	SER	-	cloning artifact	UNP Q07817

- Molecule 2 is a protein called ALPHA-BETA-PUMA BH3 FOLDAMER.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	22	182	113	35	33	1	0	0	1
2	D	22	182	114	34	33	1	0	0	1

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	37	Total	O	0	0
			37	37		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total 2	O 2	0	0
3	C	26	Total 26	O 26	0	0
3	D	3	Total 3	O 3	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: BCL-2-LIKE PROTEIN 1



- Molecule 1: BCL-2-LIKE PROTEIN 1



- Molecule 2: ALPHA-BETA-PUMA BH3 FOLDAMER



- Molecule 2: ALPHA-BETA-PUMA BH3 FOLDAMER



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.92Å 71.38Å 75.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.44 – 2.24 45.44 – 2.24	Depositor EDS
% Data completeness (in resolution range)	99.3 (45.44-2.24) 99.3 (45.44-2.24)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.09 (at 2.24Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.190 , 0.253 0.186 , 0.248	Depositor DCC
R_{free} test set	787 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	29.2	Xtrriage
Anisotropy	0.147	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.44 , 54.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2761	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACE, HR7, B3Q, HT7, NH2, B3D, B3E, B3A

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/1224	0.51	0/1658
1	C	0.46	0/1156	0.53	0/1564
2	B	0.39	0/114	0.62	0/143
2	D	0.42	0/114	0.60	0/143
All	All	0.43	0/2608	0.53	0/3508

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	5
2	D	0	5
All	All	0	10

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (10) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	102	ASN	Peptide
2	B	88	ARG	Peptide
2	B	92	ALA	Peptide
2	B	95	ARG	Peptide
2	B	99	ASP	Peptide
2	D	102	ASN	Peptide
2	D	88	ARG	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
2	D	92	ALA	Peptide
2	D	95	ARG	Peptide
2	D	99	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1201	0	1135	27	0
1	C	1128	0	1079	21	0
2	B	182	0	172	6	0
2	D	182	0	170	5	0
3	A	37	0	0	0	0
3	B	2	0	0	0	0
3	C	26	0	0	1	0
3	D	3	0	0	0	0
All	All	2761	0	2556	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:194:LEU:HD13	1:C:195:TYR:CD1	2.04	0.91
1:A:3:GLN:NE2	1:A:3:GLN:CG	2.41	0.83
1:A:201:ALA:O	1:A:202:GLU:HB2	1.79	0.81
1:A:121:GLN:CG	1:A:121:GLN:NE2	2.43	0.81
1:A:3:GLN:NE2	1:C:26:GLN:HE21	1.79	0.80
1:C:194:LEU:CD1	1:C:195:TYR:HD1	1.97	0.78
1:C:96:GLU:O	1:C:100:ARG:HB2	1.85	0.77
1:C:194:LEU:HD13	1:C:195:TYR:HD1	1.45	0.75
1:A:158:GLU:CG	1:A:158:GLU:OE1	2.37	0.72
1:A:158:GLU:CG	1:A:158:GLU:OE2	2.38	0.72
1:A:3:GLN:CG	1:A:3:GLN:OE1	2.39	0.71
1:A:3:GLN:NE2	1:A:3:GLN:OE1	2.24	0.70

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:121:GLN:CG	1:A:121:GLN:OE1	2.40	0.70
1:A:121:GLN:NE2	1:A:121:GLN:OE1	2.25	0.70
1:A:3:GLN:NE2	1:C:26:GLN:NE2	2.40	0.68
1:A:153:GLU:O	1:A:157:LYS:HG2	1.98	0.63
2:D:95:ARG:NH1	2:D:96:HR7:NZ	2.48	0.61
1:C:112:LEU:HD13	1:C:114:ILE:HG12	1.81	0.61
1:A:153:GLU:HG2	1:A:157:LYS:HD3	1.82	0.59
1:A:158:GLU:OE2	1:A:158:GLU:OE1	2.20	0.59
1:C:112:LEU:O	1:C:112:LEU:HD12	2.04	0.57
1:A:108:LEU:HD22	2:B:90:ILE:HB	1.88	0.54
1:A:26:GLN:O	1:A:86:VAL:HG23	2.08	0.53
1:C:189:ASP:HB2	3:C:2007:HOH:O	2.08	0.53
1:A:129:GLU:HG2	2:B:88:ARG:O	2.09	0.53
1:C:194:LEU:CD1	1:C:195:TYR:CD1	2.77	0.52
1:C:105:PHE:O	1:C:109:THR:HG23	2.11	0.51
1:A:105:PHE:HB2	2:B:97:MET:HE3	1.91	0.51
2:B:86:HT7:HC2	2:B:86:HT7:CE2	2.40	0.51
1:A:175:ASN:OD1	1:C:2:SER:HA	2.11	0.50
2:B:86:HT7:HC2	2:B:86:HT7:CZ3	2.42	0.49
2:D:102:ASN:O	2:D:106:GLU:HB3	2.13	0.48
1:C:16:LYS:HD3	1:C:16:LYS:HA	1.65	0.48
1:A:98:GLU:O	1:A:102:ARG:HA	2.14	0.47
1:A:88:GLN:O	1:A:92:GLU:HG3	2.15	0.47
1:C:132:ARG:HG3	1:C:133:ASP:N	2.30	0.47
1:C:179:GLU:O	1:C:183:GLN:HG2	2.15	0.46
1:A:108:LEU:O	1:A:111:GLN:HB3	2.16	0.45
2:D:88:ARG:CG	2:D:89:B3E:HD2	2.46	0.45
1:C:133:ASP:OD1	1:C:133:ASP:N	2.48	0.44
2:D:88:ARG:HG2	2:D:89:B3E:HD2	1.98	0.44
1:A:172:THR:O	1:A:176:ASP:HB2	2.17	0.44
1:A:174:LEU:HA	1:A:178:LEU:HB2	2.00	0.43
1:C:102:ARG:O	1:C:106:SER:HB3	2.18	0.43
1:A:116:PRO:HA	1:A:162:LEU:HD21	2.01	0.42
1:A:16:LYS:HD3	1:A:16:LYS:HA	1.77	0.42
1:C:160:GLN:H	1:C:160:GLN:CD	2.22	0.42
2:B:94:LEU:HD23	2:B:94:LEU:HA	1.90	0.42
2:D:93:B3Q:H8	2:D:93:B3Q:HAA	1.74	0.41
1:C:174:LEU:HA	1:C:178:LEU:HB2	2.02	0.41
1:C:112:LEU:CD1	1:C:114:ILE:HG12	2.50	0.40
1:A:152:VAL:HG11	1:C:20:LYS:HG3	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	149/158 (94%)	145 (97%)	4 (3%)	0	100	100
1	C	138/158 (87%)	137 (99%)	1 (1%)	0	100	100
2	B	15/23 (65%)	14 (93%)	1 (7%)	0	100	100
2	D	14/23 (61%)	13 (93%)	1 (7%)	0	100	100
All	All	316/362 (87%)	309 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	122/131 (93%)	116 (95%)	6 (5%)	25	25
1	C	117/131 (89%)	109 (93%)	8 (7%)	16	13
2	B	11/11 (100%)	10 (91%)	1 (9%)	9	6
2	D	11/11 (100%)	11 (100%)	0	100	100
All	All	261/284 (92%)	246 (94%)	15 (6%)	20	19

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	ARG
1	A	103	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	150	LEU
1	A	157	LYS
1	A	176	ASP
1	A	178	LEU
2	B	90	ILE
1	C	4	SER
1	C	90	LEU
1	C	99	LEU
1	C	106	SER
1	C	133	ASP
1	C	160	GLN
1	C	178	LEU
1	C	194	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	128	ASN
1	C	26	GLN
1	C	128	ASN
1	C	175	ASN
1	C	177	HIS
1	C	183	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

12 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	B3Q	D	93	2	9,9,10	0.60	0	9,10,12	1.52	2 (22%)
2	B3E	B	89	2	6,9,10	0.70	0	6,10,12	0.58	0
2	B3Q	B	93	2	9,9,10	0.70	0	9,10,12	1.03	0
2	B3A	B	103	2	5,5,6	0.83	0	5,5,7	1.51	1 (20%)
2	B3A	D	103	2	5,5,6	0.69	0	5,5,7	2.70	2 (40%)
2	B3D	B	100	2	5,8,9	0.45	0	4,9,11	1.58	1 (25%)
2	HT7	D	86	2	15,16,17	0.66	0	14,21,23	0.78	0
2	HR7	B	96	2	11,11,12	0.88	1 (9%)	10,12,14	1.33	1 (10%)
2	HR7	D	96	2	11,11,12	0.77	0	10,12,14	1.10	0
2	HT7	B	86	2	15,16,17	0.59	0	14,21,23	0.91	0
2	B3E	D	89	2	6,9,10	0.70	0	6,10,12	1.21	0
2	B3D	D	100	2	5,8,9	0.68	0	4,9,11	2.62	3 (75%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	B3Q	D	93	2	-	3/8/8/9	-
2	B3E	B	89	2	-	4/6/8/9	-
2	B3Q	B	93	2	-	2/8/8/9	-
2	B3A	B	103	2	-	1/3/3/4	-
2	B3A	D	103	2	-	1/3/3/4	-
2	B3D	B	100	2	-	2/5/7/8	-
2	HT7	D	86	2	-	0/6/7/8	0/2/2/2
2	HR7	B	96	2	-	4/8/10/11	-
2	HR7	D	96	2	-	2/8/10/11	-
2	HT7	B	86	2	-	3/6/7/8	0/2/2/2
2	B3E	D	89	2	-	3/6/8/9	-
2	B3D	D	100	2	-	1/5/7/8	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	96	HR7	CH-NZ	2.06	1.33	1.28

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	103	B3A	CG-CB-CA	-5.39	105.42	112.07
2	D	100	B3D	CA-CB-CG	-3.90	105.17	110.81
2	D	93	B3Q	CB-CA-C	-3.08	107.73	112.25
2	B	103	B3A	CG-CB-CA	-3.02	108.35	112.07
2	B	96	HR7	CD-CE-NZ	2.87	115.89	110.66
2	B	100	B3D	CA-CB-CG	-2.76	106.82	110.81
2	D	100	B3D	CB-CA-C	2.36	115.72	112.25
2	D	93	B3Q	CG-CB-N	2.35	115.87	109.03
2	D	103	B3A	CB-CA-C	-2.11	109.16	113.37
2	D	100	B3D	CD-CG-CB	2.05	115.13	112.95

There are no chirality outliers.

All (26) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	93	B3Q	C-CA-CB-CG
2	B	89	B3E	N-CB-CG-CD
2	B	89	B3E	CA-CB-CG-CD
2	B	96	HR7	CE-CD-CG-CB
2	B	86	HT7	C-CA-CB-N
2	B	86	HT7	C-CA-CB-CG
2	B	96	HR7	CG-CD-CE-NZ
2	B	89	B3E	CE-CD-CG-CB
2	B	93	B3Q	O-C-CA-CB
2	B	103	B3A	O-C-CA-CB
2	D	103	B3A	O-C-CA-CB
2	B	100	B3D	O-C-CA-CB
2	B	96	HR7	O-C-CA-CB
2	D	96	HR7	O-C-CA-CB
2	B	96	HR7	CD-CE-NZ-CH
2	D	93	B3Q	C-CA-CB-N
2	B	89	B3E	C-CA-CB-N
2	B	93	B3Q	C-CA-CB-N
2	B	100	B3D	C-CA-CB-N
2	D	96	HR7	C-CA-CB-N
2	D	89	B3E	C-CA-CB-N
2	D	93	B3Q	O-C-CA-CB
2	D	89	B3E	O-C-CA-CB
2	D	100	B3D	O-C-CA-CB
2	B	86	HT7	CA-CB-CG-CD
2	D	89	B3E	C-CA-CB-CG

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	93	B3Q	1	0
2	D	96	HR7	1	0
2	B	86	HT7	2	0
2	D	89	B3E	2	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	151/158 (95%)	-0.06	2 (1%) 77 78	20, 32, 50, 71	3 (1%)
1	C	140/158 (88%)	0.01	6 (4%) 35 34	18, 29, 45, 69	1 (0%)
2	B	15/23 (65%)	-0.07	0 100 100	27, 31, 43, 49	0
2	D	15/23 (65%)	0.09	1 (6%) 17 17	23, 27, 44, 58	0
All	All	321/362 (88%)	-0.02	9 (2%) 53 53	18, 30, 49, 71	4 (1%)

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	3	GLN	5.3
2	D	105	TYR	3.5
1	C	2	SER	3.1
1	A	112	LEU	3.0
1	C	194	LEU	2.7
1	C	1	MET	2.6
1	C	195	TYR	2.4
1	C	112	LEU	2.4
1	A	200	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	HT7	B	86	15/16	0.68	0.27	38,57,68,68	0
2	B3Q	D	93	10/11	0.80	0.29	23,30,45,49	0
2	HR7	B	96	12/13	0.88	0.21	28,30,50,58	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	HR7	D	96	12/13	0.91	0.22	25,31,51,58	0
2	B3E	B	89	10/11	0.91	0.18	32,41,58,65	0
2	B3Q	B	93	10/11	0.92	0.15	26,30,37,38	0
2	B3E	D	89	10/11	0.92	0.17	25,29,48,54	0
2	HT7	D	86	15/16	0.94	0.14	29,38,46,47	0
2	B3D	B	100	9/10	0.95	0.11	30,33,43,43	0
2	B3A	B	103	6/7	0.96	0.13	34,36,38,44	0
2	B3A	D	103	6/7	0.96	0.11	29,32,33,34	0
2	B3D	D	100	9/10	0.97	0.12	24,27,34,40	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.