



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 13, 2023 – 11:08 PM JST

PDB ID : 5YNL  
Title : Crystal structure of a cold-adapted arginase from psychrophilic yeast, *Glaciozyma antarctica*  
Authors : Yusof, N.Y.; Quay, D.H.X.; Illias, R.M.; Firdaus-Raih, M.; Abu-Bakar, F.D.; Mahadi, N.M.; Murad, A.M.A.  
Deposited on : 2017-10-24  
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2096 (2.36-2.32)
Clashscore	141614	2193 (2.36-2.32)
Ramachandran outliers	138981	2159 (2.36-2.32)
Sidechain outliers	138945	2160 (2.36-2.32)
RSRZ outliers	127900	2067 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	362	

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 2092 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Arginase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	272	2037	1295	351	379	12	0	3	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	55	Total	O	0	0
			55	55		

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Arginase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	91.19Å 91.19Å 107.78Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.77 – 2.35 28.77 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.0 (28.77-2.35) 99.0 (28.77-2.35)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.57 (at 2.34Å)	Xtriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, $R_{free}$	0.224 , 0.266 0.224 , 0.263	Depositor DCC
$R_{free}$ test set	692 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	39.0	Xtriage
Anisotropy	0.181	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.25 , 29.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.074 for $-2/3^*h-1/3^*k+2/3^*l,-1/3^*h-2/3^*k-2/3^*l,2/3^*h-2/3^*k+1/3^*l$ 0.056 for $-h,1/3^*h-1/3^*k+2/3^*l,2/3^*h+4/3^*k+1/3^*l$ 0.067 for $-1/3^*h+1/3^*k-2/3^*l,-k,-4/3^*h-2/3^*k+1/3^*l$ 0.048 for $-h,2/3^*h+1/3^*k-2/3^*l,-2/3^*h-4/3^*k-1/3^*l$ 0.045 for $1/3^*h+2/3^*k+2/3^*l,-k,4/3^*h+2/3^*k-1/3^*l$ 0.065 for $-1/3^*h-2/3^*k-2/3^*l,-2/3^*h-1/3^*k+2/3^*l,-2/3^*h+2/3^*k-1/3^*l$ 0.086 for $h,-h-k,-l$	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	2092	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.76% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.47	1/2086 (0.0%)	0.63	0/2836

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	312	CYS	CB-SG	-5.37	1.73	1.81

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	109	GLY	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2037	0	1987	46	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	55	0	0	6	0
All	All	2092	0	1987	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:115:ALA:HA	1:A:118:THR:HG22	1.56	0.86
1:A:282:VAL:O	2:A:401:HOH:O	1.98	0.81
1:A:32:HIS:CE1	1:A:110:GLY:HA2	2.21	0.74
1:A:273:GLU:OE2	2:A:402:HOH:O	2.07	0.72
1:A:118:THR:HG21	1:A:285:ASP:OD2	1.94	0.67
1:A:32:HIS:HE1	1:A:288:GLU:HA	1.60	0.67
1:A:157:SER:HB2	1:A:162:LEU:HD21	1.77	0.66
1:A:48:GLU:OE2	2:A:403:HOH:O	2.16	0.62
1:A:138:ALA:HB3	1:A:196:GLY:HA3	1.81	0.62
1:A:269:HIS:CE1	1:A:312:CYS:HG	2.19	0.61
1:A:197:GLU:HG3	1:A:199:LYS:H	1.67	0.59
1:A:170:ILE:HG23	1:A:171:PRO:HD3	1.86	0.57
1:A:137:HIS:ND1	1:A:194:ASP:HB3	2.22	0.54
1:A:248:ASP:OD1	2:A:404:HOH:O	2.18	0.54
1:A:98:HIS:ND1	1:A:103:GLN:OE1	2.31	0.53
1:A:153:GLY:HA3	1:A:195:ALA:HB3	1.91	0.53
1:A:18:VAL:HG22	1:A:107:THR:HG23	1.91	0.52
1:A:191:ARG:HG2	1:A:192:ASP:H	1.75	0.52
1:A:32:HIS:CD2	1:A:290:ASN:H	2.29	0.50
1:A:193:ILE:HG12	1:A:198:ARG:HG2	1.95	0.49
1:A:96:TYR:CZ	1:A:100:SER:HB3	2.48	0.49
1:A:139:ASP:HB3	1:A:155:PRO:HD2	1.95	0.48
1:A:21:PRO:O	1:A:32:HIS:N	2.47	0.48
1:A:140:ILE:HG22	1:A:196:GLY:HA2	1.94	0.48
1:A:32:HIS:HD2	1:A:34:GLY:H	1.61	0.47
1:A:238:ILE:HD12	1:A:280:LEU:HB2	1.95	0.47
1:A:267:GLU:O	1:A:271:ILE:HG13	2.15	0.47
1:A:115:ALA:HA	1:A:118:THR:CG2	2.35	0.46
1:A:140:ILE:HG13	1:A:156:VAL:HB	1.97	0.46
1:A:124:LYS:HE3	1:A:124:LYS:HB2	1.79	0.45
1:A:57:ALA:HB3	1:A:61:ALA:HB2	1.99	0.45
1:A:6:SER:HA	1:A:7:PRO:HD3	1.78	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:38:LEU:HA	1:A:303:VAL:HG13	2.00	0.43
1:A:224[A]:GLU:HG2	2:A:409:HOH:O	2.18	0.43
1:A:2:SER:HB2	2:A:434:HOH:O	2.18	0.42
1:A:191:ARG:HG2	1:A:192:ASP:N	2.34	0.42
1:A:79:LYS:HB3	1:A:81:PRO:HD3	2.01	0.42
1:A:140:ILE:H	1:A:196:GLY:HA2	1.85	0.42
1:A:192:ASP:O	1:A:194:ASP:N	2.54	0.41
1:A:14:THR:HB	1:A:103:GLN:HG2	2.02	0.41
1:A:269:HIS:NE2	1:A:312:CYS:SG	2.81	0.41
1:A:9:LEU:HD12	1:A:9:LEU:HA	1.88	0.41
1:A:45:ASN:ND2	1:A:49:ASN:HD21	2.19	0.41
1:A:193:ILE:HG22	1:A:193:ILE:O	2.21	0.40
1:A:207:LYS:HB2	1:A:229:HIS:CD2	2.57	0.40
1:A:248:ASP:HA	1:A:249:PRO:HD2	1.87	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	263/362 (73%)	242 (92%)	15 (6%)	6 (2%)	<b>6</b> <b>3</b>

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	171	PRO
1	A	193	ILE
1	A	260	ARG
1	A	100	SER
1	A	290	ASN
1	A	154	CYS



### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	214/298 (72%)	194 (91%)	20 (9%)	<b>9</b> <b>8</b>

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	9	LEU
1	A	38	LEU
1	A	62	LEU
1	A	83	LEU
1	A	90	ASP
1	A	114	LEU
1	A	124	LYS
1	A	162	LEU
1	A	163	ASP
1	A	170	ILE
1	A	174	SER
1	A	184	ARG
1	A	191	ARG
1	A	194	ASP
1	A	239	HIS
1	A	245	ASP
1	A	275	ILE
1	A	302	THR
1	A	317	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	45	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	272/362 (75%)	-0.22	11 (4%)	38 49	18, 50, 106, 175	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	194	ASP	5.1
1	A	193	ILE	4.2
1	A	195	ALA	4.1
1	A	318	LEU	3.5
1	A	149	GLY	3.4
1	A	164	GLY	3.2
1	A	196	GLY	3.0
1	A	170	ILE	3.0
1	A	152	HIS	2.8
1	A	11	THR	2.5
1	A	22	PHE	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.