



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 5, 2024 – 02:00 am GMT

PDB ID : 4YP9  
Title : Crystal Structure of LuxP In Complex With a Formose Derived AI-2 Analogue  
Authors : McDonough, M.A.; Sattin, S.; Gardner, P.M.; Liu, C.; Davis, B.G.  
Deposited on : 2015-03-12  
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	345	86% 11% ..
1	B	345	89% 9% ..

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5637 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Autoinducer 2-binding periplasmic protein LuxP.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	337	Total	C	N	O	S	0	0	0
			2712	1729	457	520	6			
1	B	342	Total	C	N	O	S	0	0	0
			2755	1755	465	529	6			


There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	21	GLY	-	expression tag	UNP P54300
A	22	SER	-	expression tag	UNP P54300
A	23	MET	-	expression tag	UNP P54300
B	21	GLY	-	expression tag	UNP P54300
B	22	SER	-	expression tag	UNP P54300
B	23	MET	-	expression tag	UNP P54300

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mg	0	0
			2	2		

- Molecule 3 is 1-deoxy-alpha-L-xylulofuranose-1,2-borate (three-letter code: A1B) (formula: C<sub>5</sub>H<sub>10</sub>BO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	B	C	O		
3	A	1	12	1	5	6	0	0
3	B	1	12	1	5	6	0	0

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
4	B	1	1	1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	83	83	83	0	0
5	B	60	60	60	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	104.73Å 104.73Å 69.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.80 – 2.70 46.84 – 2.70	Depositor EDS
% Data completeness (in resolution range)	97.7 (46.80-2.70) 97.8 (46.84-2.70)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	13.38 (at 2.69Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, $R_{free}$	0.167 , 0.218 0.169 , 0.220	Depositor DCC
$R_{free}$ test set	1049 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.9	Xtrriage
Anisotropy	0.081	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 30.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.035 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5637	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.38% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, A1B, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.39	0/2776	0.56	0/3764
1	B	0.38	0/2820	0.56	1/3824 (0.0%)
All	All	0.39	0/5596	0.56	1/7588 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	225	ARG	NE-CZ-NH1	7.53	124.06	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2712	0	2631	16	0
1	B	2755	0	2676	17	0
2	A	2	0	0	0	0
3	A	12	0	0	0	0
3	B	12	0	0	0	0
4	B	1	0	0	0	0
5	A	83	0	0	0	0
5	B	60	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5637	0	5307	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:76:GLN:NE2	1:B:112:ASP:OD2	2.21	0.74
1:A:111:ALA:O	1:A:117:SER:N	2.31	0.63
1:A:108:ARG:O	1:A:110:ASN:N	2.37	0.57
1:B:40:GLU:N	1:B:40:GLU:OE1	2.36	0.55
1:A:178:PHE:CD2	1:A:310:ARG:HB2	2.42	0.55
1:A:223:VAL:HG13	1:A:229:PHE:CD2	2.41	0.54
1:B:225:ARG:CG	1:B:225:ARG:HH11	2.21	0.54
1:A:136:ASP:H	1:A:140:HIS:HD2	1.57	0.52
1:B:223:VAL:HG13	1:B:229:PHE:HB2	1.94	0.49
1:B:178:PHE:CD2	1:B:310:ARG:HB2	2.46	0.49
1:B:63:THR:OG1	1:B:65:ARG:O	2.21	0.48
1:B:178:PHE:HB3	1:B:340:PHE:CZ	2.49	0.48
1:B:158:GLN:HE21	1:B:319:MET:HE1	1.79	0.47
1:B:350:GLU:OE2	1:B:350:GLU:N	2.42	0.47
1:B:111:ALA:HB1	1:B:112:ASP:HA	1.95	0.47
1:B:158:GLN:HE21	1:B:319:MET:CE	2.28	0.47
1:A:106:PHE:N	1:A:106:PHE:CD1	2.82	0.46
1:B:34:PHE:CE2	1:B:41:GLN:HG2	2.50	0.46
1:B:133:PHE:CD2	1:B:134:THR:O	2.69	0.45
1:B:178:PHE:CE2	1:B:310:ARG:HB2	2.51	0.45
1:A:106:PHE:N	1:A:106:PHE:HD1	2.13	0.45
1:A:82:TRP:CZ3	1:A:158:GLN:HG2	2.52	0.44
1:A:132:ILE:CG2	1:A:319:MET:HE1	2.47	0.44
1:A:118:LEU:HD21	1:A:139:ARG:HB3	2.00	0.44
1:A:106:PHE:CE2	1:B:110:ASN:HB3	2.54	0.43
1:A:223:VAL:HG13	1:A:229:PHE:HB2	2.00	0.43
1:B:121:MET:CE	1:B:124:LEU:HD12	2.49	0.43
1:B:178:PHE:CE2	1:B:338:GLY:HA3	2.54	0.42
1:A:133:PHE:CG	1:A:134:THR:N	2.86	0.42
1:A:156:ILE:HG21	1:A:319:MET:HE3	2.00	0.42
1:A:205:TYR:HB2	1:A:236:TYR:HA	2.01	0.42

There are no symmetry-related clashes.



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	333/345 (96%)	316 (95%)	13 (4%)	4 (1%)	13	32
1	B	340/345 (99%)	325 (96%)	14 (4%)	1 (0%)	41	66
All	All	673/690 (98%)	641 (95%)	27 (4%)	5 (1%)	22	46

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	110	ASN
1	A	109	PRO
1	A	118	LEU
1	B	97	ASN
1	A	196	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	291/299 (97%)	272 (94%)	19 (6%)	17	38
1	B	296/299 (99%)	282 (95%)	14 (5%)	26	54
All	All	587/598 (98%)	554 (94%)	33 (6%)	21	45

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	40	GLU
1	A	65	ARG
1	A	85	ASN
1	A	97	ASN
1	A	102	LEU
1	A	105	VAL
1	A	106	PHE
1	A	118	LEU
1	A	120	LEU
1	A	131	LEU
1	A	138	THR
1	A	155	LEU
1	A	221	HIS
1	A	223	VAL
1	A	267	ASP
1	A	289	TRP
1	A	351	ARG
1	A	358	ARG
1	B	25	LEU
1	B	48	LEU
1	B	65	ARG
1	B	94	TYR
1	B	108	ARG
1	B	118	LEU
1	B	121	MET
1	B	127	LYS
1	B	152	ASN
1	B	223	VAL
1	B	267	ASP
1	B	289	TRP
1	B	351	ARG
1	B	357	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	97	ASN
1	A	103	ASN
1	A	140	HIS
1	A	224	ASN
1	A	312	ASN
1	B	115	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	224	ASN
1	B	228	ASN
1	B	242	GLN
1	B	312	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	A1B	A	403	-	11,13,13	1.74	3 (27%)	12,22,22	2.45	5 (41%)
3	A1B	B	402	-	11,13,13	1.61	2 (18%)	12,22,22	2.30	5 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A1B	A	403	-	-	-	0/2/2/2
3	A1B	B	402	-	-	-	0/2/2/2

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	403	A1B	O5-C6	3.41	1.51	1.42
3	B	402	A1B	O5-C6	2.98	1.50	1.42
3	A	403	A1B	O5-C4	2.82	1.49	1.42
3	B	402	A1B	O5-C4	2.44	1.48	1.42
3	A	403	A1B	C7-C8	-2.07	1.48	1.52

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	403	A1B	B-O1-C8	-5.53	104.90	110.39
3	B	402	A1B	B-O1-C8	-4.62	105.81	110.39
3	B	402	A1B	C6-C7-C8	3.55	106.41	102.19
3	A	403	A1B	O1-C8-C4	3.45	107.98	103.85
3	B	402	A1B	O1-C8-C4	3.44	107.97	103.85
3	B	402	A1B	O1-B-O3	3.26	107.29	104.53
3	A	403	A1B	O1-B-O3	3.19	107.23	104.53
3	A	403	A1B	O3-C4-C11	2.91	111.75	107.91
3	A	403	A1B	C6-C7-C8	2.50	105.16	102.19
3	B	402	A1B	O3-C4-C11	2.06	110.63	107.91

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	337/345 (97%)	-0.37	1 (0%) 94 95	13, 23, 59, 80	0
1	B	342/345 (99%)	-0.29	2 (0%) 89 91	14, 27, 66, 87	0
All	All	679/690 (98%)	-0.33	3 (0%) 92 93	13, 24, 64, 87	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	106	PHE	2.2
1	B	64	GLN	2.2
1	B	111	ALA	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	A1B	B	402	12/12	0.92	0.21	26,33,40,44	0
3	A1B	A	403	12/12	0.93	0.18	19,25,31,32	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MG	A	401	1/1	0.95	0.09	19,19,19,19	0
4	CA	B	401	1/1	0.96	0.06	37,37,37,37	0
2	MG	A	402	1/1	0.97	0.06	21,21,21,21	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.