



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 6, 2021 – 04:11 am BST

PDB ID : 6YT3
Title : Structure of the MoStoNano fusion protein
Authors : Benoit, R.M.; Bierig, T.; Collu, C.; Engilberge, S.; Olieric, V.
Deposited on : 2020-04-23
Resolution : 2.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3168 (2.90-2.82)
Clashscore	141614	3438 (2.90-2.82)
Ramachandran outliers	138981	3348 (2.90-2.82)
Sidechain outliers	138945	3351 (2.90-2.82)
RSRZ outliers	127900	3103 (2.90-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	293	 11% 74% 9% 17%
2	B	379	 11% 87% 11% ..

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 4923 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Molybdenum storage protein subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	243	1831	1160	342	326	3	0	9	0

There are 17 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	277	GLU	-	expression tag	UNP P84308
A	278	ASN	-	expression tag	UNP P84308
A	279	LEU	-	expression tag	UNP P84308
A	280	TYR	-	expression tag	UNP P84308
A	281	PHE	-	expression tag	UNP P84308
A	282	GLN	-	expression tag	UNP P84308
A	283	GLY	-	expression tag	UNP P84308
A	284	SER	-	expression tag	UNP P84308
A	285	ALA	-	expression tag	UNP P84308
A	286	TRP	-	expression tag	UNP P84308
A	287	SER	-	expression tag	UNP P84308
A	288	HIS	-	expression tag	UNP P84308
A	289	PRO	-	expression tag	UNP P84308
A	290	GLN	-	expression tag	UNP P84308
A	291	PHE	-	expression tag	UNP P84308
A	292	GLU	-	expression tag	UNP P84308
A	293	LYS	-	expression tag	UNP P84308


- Molecule 2 is a protein called Thioredoxin, Molybdenum storage protein subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	376	2818	1793	485	529	11	67	11	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	MET	-	initiating methionine	UNP A0A0U0X1R7
B	0	GLY	-	expression tag	UNP A0A0U0X1R7
B	108	GLU	-	linker	UNP A0A0U0X1R7
B	109	GLU	-	linker	UNP A0A0U0X1R7
B	110	GLU	-	linker	UNP A0A0U0X1R7
B	111	LYS	-	linker	UNP A0A0U0X1R7
B	112	ARG	-	linker	UNP A0A0U0X1R7
B	113	LYS	-	linker	UNP A0A0U0X1R7
B	114	ARG	-	linker	UNP A0A0U0X1R7
B	115	GLU	-	linker	UNP A0A0U0X1R7

- Molecule 3 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
3	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	100	Total 103	O 103	0	3
5	B	103	Total 108	O 108	0	5

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Molybdenum storage protein subunit alpha

Chain A: 



- Molecule 2: Thioredoxin, Molybdenum storage protein subunit beta

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	165.70Å 165.70Å 352.16Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.89 – 2.85 47.89 – 2.85	Depositor EDS
% Data completeness (in resolution range)	76.3 (47.89-2.85) 76.3 (47.89-2.85)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.28 (at 2.86Å)	Xtrriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.233 , 0.263 0.259 , 0.286	Depositor DCC
R_{free} test set	1695 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	51.8	Xtrriage
Anisotropy	0.110	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	4923	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.46	0/1880	0.69	0/2563
2	B	0.46	0/2868	0.68	0/3892
All	All	0.46	0/4748	0.68	0/6455

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1831	0	1881	16	0
2	B	2818	0	2872	20	0
3	A	31	0	12	0	0
3	B	31	0	12	1	0
4	A	1	0	0	0	0
5	A	103	0	0	3	0
5	B	108	0	0	0	0
All	All	4923	0	4777	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:78:THR:HG21	1:A:113:GLY:CA	2.12	0.78
2:B:174:ARG:HD3	2:B:248:VAL:O	1.88	0.73
1:A:42:GLN:NE2	5:A:401:HOH:O	2.25	0.68
1:A:255:ARG:NH1	5:A:402:HOH:O	2.27	0.67
2:B:332:ILE:HG12	2:B:352:VAL:HG11	1.80	0.63
1:A:78:THR:HG21	1:A:113:GLY:HA2	1.81	0.61
2:B:211:SER:HA	2:B:256:MET:HG3	1.87	0.56
2:B:60:ILE:HD12	2:B:67:ALA:HB2	1.87	0.56
1:A:51:MET:HG3	1:A:112:ASN:HB3	1.89	0.55
2:B:206:LEU:HA	2:B:209:LEU:HD12	1.90	0.54
1:A:159:GLU:CD	1:A:167:PRO:HD2	2.33	0.48
2:B:230:ILE:HG12	2:B:250:ALA:HB3	1.96	0.48
2:B:79:LEU:HD22	2:B:89:THR:HG22	1.95	0.48
2:B:357:VAL:HB	2:B:360:ASN:ND2	2.29	0.48
1:A:255:ARG:HD3	1:A:265:VAL:HG22	1.96	0.47
2:B:281:PHE:HA	2:B:292:MET:HE3	1.95	0.47
1:A:237:ALA:HB2	5:A:403:HOH:O	2.15	0.46
1:A:248:VAL:HG11	1:A:259:ALA:HB2	1.97	0.46
1:A:88:PHE:CE1	1:A:105:ALA:HB2	2.53	0.44
2:B:334:GLU:HB2	2:B:337:VAL:HG23	1.99	0.44
1:A:252:VAL:HB	1:A:255:ARG:HG3	2.00	0.43
2:B:245:LEU:HD23	2:B:245:LEU:HA	1.86	0.42
2:B:297:ASP:HA	2:B:356:LEU:HD13	2.01	0.42
2:B:72:ILE:HD11	2:B:79:LEU:HD11	2.01	0.42
1:A:76:ILE:O	1:A:148:VAL:HA	2.19	0.42
1:A:77:LEU:HB3	1:A:151:ALA:HA	2.02	0.42
3:B:401:ATP:H2'	3:B:401:ATP:N3	2.35	0.42
1:A:103:PRO:HG3	1:A:156:HIS:HA	2.02	0.41
2:B:119:LEU:HG	2:B:119:LEU:O	2.20	0.41
2:B:26:ASP:HB2	2:B:42:LEU:HD21	2.02	0.41
2:B:7:LEU:HB2	2:B:58:LEU:HD12	2.03	0.41
2:B:77:THR:HG22	2:B:91:VAL:HG13	2.02	0.41
2:B:229:GLY:C	2:B:248:VAL:HG21	2.41	0.41
1:A:156:HIS:HD2	1:A:157:HIS:CD2	2.38	0.41
2:B:255:GLY:O	2:B:279:GLY:HA3	2.21	0.41
2:B:67:ALA:HB3	2:B:68:PRO:HD3	2.03	0.40
1:A:34:PRO:O	1:A:241:HIS:HE1	2.05	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	244/293 (83%)	234 (96%)	10 (4%)	0	100	100
2	B	374/379 (99%)	360 (96%)	14 (4%)	0	100	100
All	All	618/672 (92%)	594 (96%)	24 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	189/231 (82%)	185 (98%)	4 (2%)	53	79
2	B	293/296 (99%)	279 (95%)	14 (5%)	25	55
All	All	482/527 (92%)	464 (96%)	18 (4%)	33	65

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	87	VAL
1	A	128	VAL
1	A	214	THR
1	A	265	VAL
2	B	2	ASP
2	B	41	ILE
2	B	47	ASP
2	B	110	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	146	THR
2	B	155	VAL
2	B	174	ARG
2	B	211	SER
2	B	238	LEU
2	B	239	SER
2	B	248	VAL
2	B	320	VAL
2	B	329	HIS
2	B	362	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	GLN
1	A	156	HIS
1	A	241	HIS
2	B	130	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ATP	A	301	4	26,33,33	0.62	0	31,52,52	0.98	2 (6%)
3	ATP	B	401	-	26,33,33	0.61	0	31,52,52	0.94	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ATP	A	301	4	-	3/18/38/38	0/3/3/3
3	ATP	B	401	-	-	8/18/38/38	0/3/3/3

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	401	ATP	C1'-N9-C4	2.50	131.04	126.64
3	B	401	ATP	C5-C6-N6	2.26	123.79	120.35
3	A	301	ATP	C5-C6-N6	2.25	123.77	120.35
3	A	301	ATP	PB-O3B-PG	2.19	140.35	132.83

There are no chirality outliers.

All (11) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	401	ATP	PB-O3B-PG-O2G
3	B	401	ATP	PB-O3B-PG-O3G
3	B	401	ATP	C5'-O5'-PA-O1A
3	B	401	ATP	C5'-O5'-PA-O2A
3	B	401	ATP	C4'-C5'-O5'-PA
3	A	301	ATP	C3'-C4'-C5'-O5'
3	B	401	ATP	C5'-O5'-PA-O3A
3	A	301	ATP	O4'-C4'-C5'-O5'
3	B	401	ATP	PB-O3A-PA-O1A
3	B	401	ATP	O4'-C4'-C5'-O5'
3	A	301	ATP	C5'-O5'-PA-O2A


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	401	ATP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	243/293 (82%)	-0.36	1 (0%) 92 92	19, 40, 82, 106	0
2	B	376/379 (99%)	0.32	43 (11%) 5 3	30, 73, 124, 139	16 (4%)
All	All	619/672 (92%)	0.06	44 (7%) 16 12	19, 58, 120, 139	16 (2%)

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	88	ALA	4.6
2	B	42	LEU	4.4
2	B	53	LEU	4.3
2	B	45	ILE	4.3
2	B	56	ALA	4.2
2	B	81	PHE	3.9
2	B	329	HIS	3.9
2	B	46	ALA	3.7
2	B	325	ALA	3.6
2	B	43	ASP	3.5
2	B	99	LEU	3.5
2	B	103	LEU	3.4
2	B	14	THR	3.3
2	B	12	PHE	3.1
2	B	55	VAL	3.0
2	B	7	LEU	3.0
2	B	26	ASP	3.0
2	B	4	ILE	2.9
2	B	39	ALA	2.9
2	B	95	SER	2.8
2	B	40	PRO	2.8
2	B	38	ILE	2.7
2	B	41	ILE	2.7
2	B	47	ASP	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	207	GLN	2.7
2	B	49	TYR	2.7
2	B	44	GLU	2.6
2	B	23	ILE	2.5
2	B	22	ALA	2.5
2	B	37	MET	2.4
2	B	63	ASN	2.4
2	B	5	ILE	2.4
2	B	6	HIS	2.4
2	B	13	ASP	2.3
2	B	25	VAL	2.2
2	B	91	VAL	2.2
2	B	10	ASP	2.2
2	B	234	GLY	2.2
2	B	96	LYS	2.2
2	B	58	LEU	2.2
2	B	313	THR	2.2
2	B	303	THR	2.1
2	B	94	LEU	2.1
2	B	62	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ATP	B	401	31/31	0.90	0.19	93,108,116,117	0
3	ATP	A	301	31/31	0.96	0.14	45,58,60,61	0
4	MG	A	302	1/1	0.98	0.12	47,47,47,47	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.