



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 23, 2023 – 02:51 AM JST

PDB ID : 7YU2
Title : Structure of 6-aminohexanoate-oligomer hydrolase NylC, D122G/H130Y/T267C mutant, hydroxylamine-treated
Authors : Negoro, S.; Higuchi, Y.
Deposited on : 2022-08-16
Resolution : 1.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1232 (1.24-1.20)
Clashscore	141614	1294 (1.24-1.20)
Ramachandran outliers	138981	1251 (1.24-1.20)
Sidechain outliers	138945	1250 (1.24-1.20)
RSRZ outliers	127900	1209 (1.24-1.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	355	 5% 80% 12% • 7%
1	B	355	 4% 80% 12% • 6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	B	401	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 11206 atoms, of which 5274 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 6-aminohexanoate-oligomer endohydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	330	5178	1629	2585	468	484	12	64	27	0
1	B	334	5274	1660	2625	477	500	12	64	30	0

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	122	GLY	ASP	engineered mutation	UNP Q79F77
A	130	TYR	HIS	engineered mutation	UNP Q79F77
A	267	CYS	THR	engineered mutation	UNP Q79F77
B	122	GLY	ASP	engineered mutation	UNP Q79F77
B	130	TYR	HIS	engineered mutation	UNP Q79F77
B	267	CYS	THR	engineered mutation	UNP Q79F77

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	2	0
			14	3	8	3		
2	A	1	Total	C	H	O	2	0
			14	3	8	3		
2	A	1	Total	C	H	O	2	0
			14	3	8	3		
2	A	1	Total	C	H	O	2	0
			14	3	8	3		
2	B	1	Total	C	H	O	2	0
			14	3	8	3		
2	B	1	Total	C	H	O	2	0
			14	3	8	3		
2	B	1	Total	C	H	O	2	0
			14	3	8	3		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	O S	0	0
			5	4 1		
3	A	1	Total	O S	0	0
			5	4 1		
3	B	1	Total	O S	0	0
			5	4 1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	317	Total 317	O 317	0	0
4	B	310	Total 310	O 310	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 6-aminohexanoate-oligomer endohydrolase



- Molecule 1: 6-aminohexanoate-oligomer endohydrolase



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	70.27Å 144.02Å 128.26Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.14 – 1.21 27.14 – 1.21	Depositor EDS
% Data completeness (in resolution range)	99.2 (27.14-1.21) 91.0 (27.14-1.21)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.17 (at 1.21Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.116 , 0.143 0.115 , 0.141	Depositor DCC
R_{free} test set	9819 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	10.7	Xtrriage
Anisotropy	0.657	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.46 , 55.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	11206	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.94	8/2690 (0.3%)	1.02	10/3650 (0.3%)
1	B	1.02	13/2732 (0.5%)	1.12	18/3709 (0.5%)
All	All	0.98	21/5422 (0.4%)	1.07	28/7359 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	66[A]	GLU	CD-OE1	11.26	1.38	1.25
1	B	66[B]	GLU	CD-OE1	11.26	1.38	1.25
1	A	136	GLU	CD-OE2	9.54	1.36	1.25
1	A	30	PHE	C-O	8.20	1.39	1.23
1	B	208[A]	GLY	C-O	7.93	1.36	1.23
1	B	208[B]	GLY	C-O	7.93	1.36	1.23
1	A	149	SER	CB-OG	7.14	1.51	1.42
1	A	239	GLN	CD-NE2	6.97	1.50	1.32
1	B	136	GLU	CD-OE1	6.56	1.32	1.25
1	B	344	GLU	CD-OE2	-6.05	1.19	1.25
1	B	292	SER	CA-CB	5.99	1.61	1.52
1	A	317	GLU	CD-OE1	-5.88	1.19	1.25
1	B	66[A]	GLU	CD-OE2	5.87	1.32	1.25
1	B	66[B]	GLU	CD-OE2	5.87	1.32	1.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	129	GLU	CD-OE2	-5.80	1.19	1.25
1	B	173	GLU	CD-OE1	5.73	1.31	1.25
1	A	344	GLU	CD-OE2	-5.71	1.19	1.25
1	A	49	GLU	CB-CG	-5.22	1.42	1.52
1	A	66	GLU	CD-OE2	5.18	1.31	1.25
1	B	130	TYR	CG-CD1	5.09	1.45	1.39
1	B	65	TYR	CZ-OH	-5.09	1.29	1.37

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	277[A]	ARG	NE-CZ-NH1	11.87	126.24	120.30
1	B	277[B]	ARG	NE-CZ-NH1	11.87	126.24	120.30
1	B	65	TYR	CD1-CE1-CZ	7.49	126.54	119.80
1	A	88[A]	ARG	NE-CZ-NH1	-6.75	116.92	120.30
1	A	88[B]	ARG	NE-CZ-NH1	-6.75	116.92	120.30
1	A	267[A]	CYS	CB-CA-C	-6.45	97.50	110.40
1	A	267[B]	CYS	CB-CA-C	-6.45	97.50	110.40
1	B	88[A]	ARG	NE-CZ-NH1	-6.14	117.23	120.30
1	B	88[B]	ARG	NE-CZ-NH1	-6.14	117.23	120.30
1	B	127	ARG	NE-CZ-NH2	-5.96	117.32	120.30
1	B	151[A]	ARG	NE-CZ-NH1	5.92	123.26	120.30
1	B	151[B]	ARG	NE-CZ-NH1	5.92	123.26	120.30
1	B	65	TYR	CE1-CZ-CE2	-5.73	110.63	119.80
1	B	159[A]	LYS	CB-CA-C	-5.69	99.02	110.40
1	B	159[B]	LYS	CB-CA-C	-5.69	99.02	110.40
1	B	163	ARG	NE-CZ-NH1	5.66	123.13	120.30
1	A	330	ARG	NE-CZ-NH2	-5.63	117.49	120.30
1	B	233	ARG	NE-CZ-NH1	-5.42	117.59	120.30
1	B	277[A]	ARG	CG-CD-NE	-5.26	100.75	111.80
1	B	277[B]	ARG	CG-CD-NE	-5.26	100.75	111.80
1	B	236	TYR	CB-CG-CD1	-5.25	117.85	121.00
1	A	163[A]	ARG	NE-CZ-NH2	-5.25	117.68	120.30
1	A	163[B]	ARG	NE-CZ-NH2	-5.25	117.68	120.30
1	A	82[A]	ARG	NE-CZ-NH2	-5.25	117.68	120.30
1	A	82[B]	ARG	NE-CZ-NH2	-5.25	117.68	120.30
1	A	226	ASP	CB-CG-OD2	-5.13	113.69	118.30
1	B	277[A]	ARG	CB-CG-CD	5.05	124.73	111.60
1	B	277[B]	ARG	CB-CG-CD	5.05	124.73	111.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	239	GLN	Sidechain
1	B	86	ASP	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2593	2585	2586	28	0
1	B	2649	2625	2617	33	0
2	A	30	40	40	2	0
2	B	18	24	24	5	0
3	A	10	0	0	1	0
3	B	5	0	0	0	0
4	A	317	0	0	7	0
4	B	310	0	0	12	1
All	All	5932	5274	5267	64	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:30:PHE:CE1	1:B:149[B]:SER:OG	2.00	1.12
1:B:30:PHE:HE1	1:B:149[B]:SER:OG	1.32	1.06
1:B:206:ARG:NH1	1:B:209[B]:ASP:OD1	2.10	0.85
1:B:277[A]:ARG:HG2	1:B:316:ASP:HA	1.61	0.82
1:B:208[A]:GLY:O	1:B:209[A]:ASP:HB2	1.81	0.79
1:A:288[A]:LYS:HE3	4:A:643:HOH:O	1.82	0.78
1:A:221[B]:VAL:HG11	1:A:305[B]:MET:HB2	1.71	0.73
2:B:401:GOL:O3	2:B:402:GOL:H32	1.91	0.71
1:A:82[B]:ARG:HD2	1:A:314:THR:CG2	2.22	0.70
1:A:284[B]:ASN:OD1	4:A:502:HOH:O	2.10	0.69
2:B:401:GOL:O3	2:B:402:GOL:C3	2.40	0.69
1:A:82[B]:ARG:HD2	1:A:314:THR:HG23	1.76	0.68
1:A:67[A]:GLU:OE1	4:A:503:HOH:O	2.12	0.67
1:B:82[B]:ARG:CG	4:B:722:HOH:O	2.42	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:170:VAL:O	4:B:503:HOH:O	2.15	0.64
1:A:163[A]:ARG:NH2	4:A:501:HOH:O	2.10	0.63
1:A:50[B]:MET:SD	1:A:52:ARG:NH2	2.79	0.56
1:B:277[A]:ARG:HG2	1:B:316:ASP:CA	2.35	0.55
1:B:76:HIS:HE1	4:B:729:HOH:O	1.89	0.55
1:B:119:GLY:HA2	1:B:159[B]:LYS:HG3	1.88	0.54
1:A:82[B]:ARG:HG2	1:A:283:LEU:HD22	1.91	0.52
1:B:76:HIS:HD2	4:B:741:HOH:O	1.93	0.52
1:A:38:PHE:CD1	1:A:67[B]:GLU:HG2	2.46	0.51
1:B:50[B]:MET:SD	1:B:52:ARG:CZ	2.98	0.51
1:A:163[B]:ARG:NH1	3:A:407:SO4:O2	2.36	0.51
1:A:244[B]:ARG:HG3	4:A:702:HOH:O	2.10	0.51
1:B:82[B]:ARG:NH2	4:B:507:HOH:O	2.43	0.51
1:B:110:ALA:HB1	2:B:401:GOL:H12	1.92	0.50
2:B:401:GOL:HO3	2:B:402:GOL:C3	2.24	0.50
1:B:114:LEU:HB3	4:B:511:HOH:O	2.11	0.50
1:A:82[B]:ARG:CD	1:A:314:THR:CG2	2.90	0.50
1:A:146:TYR:HB3	2:A:402:GOL:H11	1.92	0.50
1:A:82[B]:ARG:HG2	1:A:283:LEU:CD2	2.42	0.50
1:B:126:GLU:HG2	1:B:130:TYR:OH	2.12	0.49
1:B:267[A]:CYS:N	1:B:306:ASP:HB3	2.27	0.49
1:B:50[B]:MET:SD	1:B:52:ARG:NH2	2.86	0.49
1:B:82[B]:ARG:HG3	4:B:705:HOH:O	2.13	0.49
1:A:69:PRO:HB2	1:A:183[B]:MET:CG	2.43	0.48
1:B:82[B]:ARG:CD	4:B:722:HOH:O	2.60	0.48
1:A:77[B]:ILE:HG21	1:A:81:ALA:HB2	1.96	0.48
1:B:82[B]:ARG:HD3	4:B:722:HOH:O	2.15	0.47
1:A:82[B]:ARG:HG3	4:A:522:HOH:O	2.14	0.47
1:A:111:GLY:N	2:A:401:GOL:H12	2.30	0.47
1:A:281:VAL:O	1:A:285[A]:GLN:HG2	2.14	0.47
1:B:69:PRO:HB2	1:B:183[B]:MET:HG3	1.96	0.47
1:B:221:VAL:HG21	1:B:305[B]:MET:HB2	1.96	0.46
1:B:82[B]:ARG:HG3	4:B:722:HOH:O	2.10	0.46
1:B:281:VAL:O	1:B:285[A]:GLN:HG2	2.16	0.46
1:A:35:ASN:OD1	1:A:155:VAL:HA	2.17	0.45
1:A:50[B]:MET:SD	1:A:52:ARG:CZ	3.05	0.45
1:A:69:PRO:HB2	1:A:183[B]:MET:HG3	1.99	0.44
1:B:69:PRO:HB2	1:B:183[B]:MET:CG	2.48	0.44
1:A:288[A]:LYS:CE	4:A:643:HOH:O	2.53	0.44
1:B:284[A]:ASN:HB3	4:B:515:HOH:O	2.19	0.43
1:A:105:CYS:O	1:A:142:SER:HA	2.18	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:218:PRO:C	1:A:220:PRO:HD3	2.40	0.42
1:B:206:ARG:NE	1:B:208[B]:GLY:O	2.50	0.42
1:B:146:TYR:HB3	2:B:403:GOL:H32	2.02	0.42
1:B:128:LEU:HD21	1:B:136:GLU:HB3	2.01	0.42
1:B:167:GLU:OE1	4:B:504:HOH:O	2.22	0.41
1:B:35:ASN:OD1	1:B:155:VAL:HA	2.20	0.41
1:A:267[B]:CYS:N	1:A:306:ASP:HB3	2.36	0.41
1:B:18:VAL:HG11	1:B:232:VAL:HB	2.01	0.41
1:A:277:ARG:HD2	1:A:319:ASP:OD1	2.20	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:531:HOH:O	4:B:531:HOH:O[3_554]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	352/355 (99%)	340 (97%)	12 (3%)	0	100	100
1	B	359/355 (101%)	348 (97%)	11 (3%)	0	100	100
All	All	711/710 (100%)	688 (97%)	23 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	264/261 (101%)	258 (98%)	6 (2%)	50	13
1	B	268/261 (103%)	263 (98%)	5 (2%)	57	19
All	All	532/522 (102%)	521 (98%)	11 (2%)	67	16

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	55	PHE
1	A	78	PRO
1	A	151[A]	ARG
1	A	151[B]	ARG
1	A	267[A]	CYS
1	A	267[B]	CYS
1	B	55	PHE
1	B	267[A]	CYS
1	B	267[B]	CYS
1	B	318[A]	ILE
1	B	318[B]	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	257	GLN
1	B	76	HIS
1	B	251	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	B	402	-	5,5,5	1.02	1 (20%)	5,5,5	0.41	0
2	GOL	B	403	-	5,5,5	0.83	0	5,5,5	0.53	0
3	SO4	B	404	-	4,4,4	0.50	0	6,6,6	0.23	0
3	SO4	A	406	-	4,4,4	0.32	0	6,6,6	0.26	0
2	GOL	A	403	-	5,5,5	0.66	0	5,5,5	0.74	0
2	GOL	A	405	-	5,5,5	0.27	0	5,5,5	0.58	0
2	GOL	B	401	-	5,5,5	0.39	0	5,5,5	0.92	0
2	GOL	A	404	-	5,5,5	0.20	0	5,5,5	0.84	0
3	SO4	A	407	-	4,4,4	0.26	0	6,6,6	0.18	0
2	GOL	A	402	-	5,5,5	0.44	0	5,5,5	0.54	0
2	GOL	A	401	-	5,5,5	1.06	0	5,5,5	1.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	402	-	-	2/4/4/4	-
2	GOL	B	403	-	-	3/4/4/4	-
2	GOL	A	403	-	-	1/4/4/4	-
2	GOL	A	405	-	-	0/4/4/4	-
2	GOL	B	401	-	-	0/4/4/4	-
2	GOL	A	404	-	-	0/4/4/4	-
2	GOL	A	402	-	-	1/4/4/4	-
2	GOL	A	401	-	-	1/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	402	GOL	O3-C3	2.14	1.51	1.42

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	GOL	O1-C1-C2-C3
2	A	403	GOL	C1-C2-C3-O3
2	B	403	GOL	O1-C1-C2-C3
2	B	403	GOL	O1-C1-C2-O2
2	B	402	GOL	O1-C1-C2-O2
2	A	402	GOL	O1-C1-C2-C3
2	B	402	GOL	O1-C1-C2-C3
2	B	403	GOL	C1-C2-C3-O3

There are no ring outliers.

6 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	402	GOL	3	0
2	B	403	GOL	1	0
2	B	401	GOL	4	0
3	A	407	SO4	1	0
2	A	402	GOL	1	0
2	A	401	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	330/355 (92%)	-0.15	17 (5%) 27 25	9, 14, 34, 54	0
1	B	334/355 (94%)	-0.18	15 (4%) 33 32	9, 13, 32, 73	0
All	All	664/710 (93%)	-0.16	32 (4%) 30 29	9, 14, 33, 73	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	260	PRO	8.4
1	B	29	VAL	6.3
1	A	322	THR	6.2
1	B	16	ILE	6.0
1	A	30	PHE	5.9
1	A	258	VAL	5.8
1	A	259	PRO	5.7
1	B	30	PHE	5.4
1	B	17	ALA	5.2
1	A	323	THR	5.0
1	B	18	VAL	4.5
1	B	322	THR	4.5
1	B	28	PRO	4.1
1	B	259	PRO	4.1
1	B	267[A]	CYS	3.9
1	A	330	ARG	3.7
1	A	29	VAL	3.7
1	B	323	THR	3.6
1	A	28	PRO	3.5
1	A	325	GLY	3.4
1	B	330	ARG	3.3
1	A	19	ASP	2.8
1	A	324	PRO	2.8
1	A	31	GLY	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	258	VAL	2.7
1	B	27	PRO	2.6
1	B	31	GLY	2.5
1	A	321	PRO	2.5
1	A	267[A]	CYS	2.3
1	A	326	SER	2.3
1	A	39	ASP	2.2
1	A	272[A]	ILE	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	B	403	6/6	0.69	0.19	29,39,59,60	2
2	GOL	B	402	6/6	0.73	0.13	49,61,66,67	2
2	GOL	A	402	6/6	0.85	0.15	29,34,52,52	2
2	GOL	A	401	6/6	0.89	0.14	18,40,67,68	2
3	SO4	B	404	5/5	0.90	0.27	47,51,70,82	0
2	GOL	B	401	6/6	0.91	0.13	20,41,45,46	2
2	GOL	A	405	6/6	0.91	0.09	40,53,60,61	2
3	SO4	A	407	5/5	0.95	0.22	41,58,88,101	0
2	GOL	A	403	6/6	0.95	0.13	18,22,26,26	2
3	SO4	A	406	5/5	0.96	0.27	50,53,72,81	0
2	GOL	A	404	6/6	0.97	0.09	19,27,45,50	2

6.5 Other polymers [i](#)

There are no such residues in this entry.