



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 24, 2024 – 11:48 AM EDT

PDB ID : 6YWA
Title : Human REF STING in complex with 3',3'-c-[2'FdAMP-2'FdAM(PS)]
Authors : Boura, E.; Smola, M.
Deposited on : 2020-04-29
Resolution : 2.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5974 (2.34-2.30)
Clashscore	141614	6604 (2.34-2.30)
Ramachandran outliers	138981	6523 (2.34-2.30)
Sidechain outliers	138945	6523 (2.34-2.30)
RSRZ outliers	127900	5855 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	204	 4% 83% 6% 10%
1	B	204	 4% 80% 8% 11%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6005 atoms, of which 2878 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Stimulator of interferon genes protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	183	2931	942	1451	254	277	7	0	6	0
1	B	181	2797	900	1373	244	273	7	0	4	0

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
2	A	1	14	3	8	3	0	0

- Molecule 3 is 3',3'-c-[2'FdAMP-2'FdAM(PS)] (three-letter code: PWB) (formula: C₂₀H₂₄F₂N₁₀O₉P₂S) (labeled as "Ligand of Interest" by depositor).

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Stimulator of interferon genes protein

Chain A: 



- Molecule 1: Stimulator of interferon genes protein

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, α , β , γ	35.96Å 94.24Å 116.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.96 – 2.31 35.96 – 2.31	Depositor EDS
% Data completeness (in resolution range)	99.3 (35.96-2.31) 99.3 (35.96-2.31)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.54 (at 2.31Å)	Xtrriage
Refinement program	PHENIX 1.14_3260	Depositor
R, R_{free}	0.194 , 0.244 0.196 , 0.244	Depositor DCC
R_{free} test set	901 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	31.9	Xtrriage
Anisotropy	0.338	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 57.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6005	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PWB, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/1527	0.42	0/2070
1	B	0.30	0/1463	0.42	0/1986
All	All	0.30	0/2990	0.42	0/4056

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1480	1451	1449	9	0
1	B	1424	1373	1371	11	0
2	A	6	8	8	0	0
3	B	88	46	0	0	0
4	A	70	0	0	2	0
4	B	59	0	0	2	0
All	All	3127	2878	2828	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:157:HIS:HE1	4:B:502:HOH:O	1.72	0.72
1:B:272:SER:HA	1:B:279:PHE:HB3	1.84	0.59
1:A:260:GLU:HA	1:B:234:GLY:HA3	1.91	0.53
1:B:302:ALA:HB1	1:B:304:GLU:OE2	2.11	0.50
1:B:211:ASN:HB3	1:B:214:MET:HG2	1.95	0.47
1:A:158:GLY:HA3	1:B:155:VAL:HG13	1.97	0.47
1:A:222[B]:LEU:HG	1:A:246:GLU:HB2	1.98	0.45
1:A:161:TRP:CH2	1:B:276:GLN:HG3	2.52	0.44
1:A:237:ASP:OD2	4:A:601:HOH:O	2.21	0.44
1:B:194:VAL:HG23	1:B:257:CYS:HA	1.99	0.44
1:A:232:HIS:CD2	4:B:555:HOH:O	2.72	0.43
1:B:304:GLU:HG2	1:B:305:SER:N	2.33	0.42
1:A:239:VAL:HB	1:B:239:VAL:HB	2.00	0.42
1:B:197:ARG:NH2	1:B:340:GLU:OE2	2.52	0.42
1:A:218:ASN:ND2	4:A:619:HOH:O	2.51	0.41
1:B:336:GLU:O	1:B:340:GLU:HG3	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	185/204 (91%)	178 (96%)	7 (4%)	0	100	100
1	B	181/204 (89%)	176 (97%)	5 (3%)	0	100	100
All	All	366/408 (90%)	354 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	158/176 (90%)	155 (98%)	3 (2%)	57	73
1	B	150/176 (85%)	149 (99%)	1 (1%)	84	92
All	All	308/352 (88%)	304 (99%)	4 (1%)	69	81

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	162	SER
1	A	213	SER
1	A	321	SER
1	B	301	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	157	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	501	-	5,5,5	0.12	0	5,5,5	0.30	0
3	PWB	B	401[A]	-	41,50,50	1.64	10 (24%)	49,78,78	1.39	7 (14%)
3	PWB	B	401[B]	-	41,50,50	1.65	10 (24%)	49,78,78	1.39	8 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	501	-	-	4/4/4/4	-
3	PWB	B	401[A]	-	-	1/20/62/62	0/6/7/7
3	PWB	B	401[B]	-	-	1/20/62/62	0/6/7/7

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	401[A]	PWB	C3-N11	3.65	1.47	1.34
3	B	401[B]	PWB	C13-N10	3.64	1.47	1.34
3	B	401[A]	PWB	C13-N10	3.44	1.46	1.34
3	B	401[B]	PWB	C3-N11	3.41	1.46	1.34
3	B	401[B]	PWB	F2-C6	-2.99	1.34	1.40
3	B	401[A]	PWB	F1-C16	-2.98	1.34	1.40
3	B	401[A]	PWB	F2-C6	-2.91	1.34	1.40
3	B	401[B]	PWB	F1-C16	-2.85	1.34	1.40
3	B	401[A]	PWB	P-O	-2.78	1.40	1.46
3	B	401[B]	PWB	P-O	-2.76	1.40	1.46
3	B	401[B]	PWB	C16-C17	-2.60	1.50	1.53
3	B	401[A]	PWB	C6-C7	-2.44	1.50	1.53
3	B	401[A]	PWB	C16-C17	-2.24	1.50	1.53
3	B	401[B]	PWB	C6-C7	-2.23	1.50	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	401[A]	PWB	C12-N7	-2.16	1.32	1.39
3	B	401[B]	PWB	C2-N3	-2.15	1.32	1.39
3	B	401[A]	PWB	P1-O22	-2.13	1.43	1.50
3	B	401[A]	PWB	C2-N3	-2.10	1.32	1.39
3	B	401[B]	PWB	C12-N7	-2.08	1.32	1.39
3	B	401[B]	PWB	P1-O22	-2.07	1.43	1.50

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	401[A]	PWB	N1-C-N	-3.64	123.74	128.67
3	B	401[B]	PWB	N5-C10-N4	-3.60	123.79	128.67
3	B	401[B]	PWB	N1-C-N	-3.31	124.19	128.67
3	B	401[A]	PWB	N5-C10-N4	-3.25	124.26	128.67
3	B	401[B]	PWB	C16-C17-N6	-3.13	105.93	112.59
3	B	401[A]	PWB	C6-C7-N2	-2.98	106.26	112.59
3	B	401[A]	PWB	C1-C2-N3	-2.78	106.40	109.34
3	B	401[B]	PWB	C11-C12-N7	-2.75	106.43	109.34
3	B	401[A]	PWB	C16-C17-N6	-2.56	107.15	112.59
3	B	401[B]	PWB	C6-C7-N2	-2.49	107.30	112.59
3	B	401[A]	PWB	O1-C7-N2	2.11	111.55	108.75
3	B	401[B]	PWB	C1-C2-N3	-2.10	107.12	109.34
3	B	401[B]	PWB	O5-C17-N6	2.09	111.52	108.75
3	B	401[B]	PWB	C6-C5-C8	-2.06	99.78	102.41
3	B	401[A]	PWB	C11-C12-N7	-2.06	107.16	109.34

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	GOL	C1-C2-C3-O3
2	A	501	GOL	O2-C2-C3-O3
2	A	501	GOL	O1-C1-C2-C3
2	A	501	GOL	O1-C1-C2-O2
3	B	401[B]	PWB	C9-O2-P-O
3	B	401[A]	PWB	C9-O2-P-O

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	183/204 (89%)	0.15	3 (1%) 72 78	22, 38, 66, 81	0
1	B	181/204 (88%)	0.25	9 (4%) 28 36	21, 39, 69, 97	1 (0%)
All	All	364/408 (89%)	0.20	12 (3%) 46 53	21, 38, 68, 97	1 (0%)

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	152	ASN	4.6
1	B	232	HIS	4.6
1	B	194	VAL	4.3
1	B	153	PHE	3.9
1	A	153	PHE	3.4
1	B	304	GLU	3.3
1	B	184	GLN	3.1
1	B	181	THR	2.8
1	B	235	ILE	2.3
1	B	226	PRO	2.3
1	A	250	ASN	2.3
1	B	193	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	A	501	6/6	0.78	0.21	38,67,96,104	0
3	PWB	B	401[A]	44/44	0.97	0.15	17,28,39,43	67
3	PWB	B	401[B]	44/44	0.97	0.15	18,28,39,43	67

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around PWB B 401 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.