



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 16, 2022 – 01:16 pm BST

PDB ID : 7Z4M  
Title : Plasmodium falciparum pyruvate kinase complexed with Mg<sup>2+</sup> and K<sup>+</sup>  
Authors : Dillenberger, M.; Rahlfs, S.; Becker, K.; Fritz-Wolf, K.  
Deposited on : 2022-03-04  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.29  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.29


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	519	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	PEG	A	604	-	-	X	-

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Type</b>	<b>Chain</b>	<b>Res</b>	<b>Chirality</b>	<b>Geometry</b>	<b>Clashes</b>	<b>Electron density</b>
5	PEG	A	606	-	-	X	-

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3988 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyruvate kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	497	3782	2381	655	716	30	0	1	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	512	LEU	-	expression tag	UNP C6KTA4
A	513	GLU	-	expression tag	UNP C6KTA4
A	514	HIS	-	expression tag	UNP C6KTA4
A	515	HIS	-	expression tag	UNP C6KTA4
A	516	HIS	-	expression tag	UNP C6KTA4
A	517	HIS	-	expression tag	UNP C6KTA4
A	518	HIS	-	expression tag	UNP C6KTA4
A	519	HIS	-	expression tag	UNP C6KTA4

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is POTASSIUM ION (three-letter code: K) (formula: K) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	K	0	0
			1	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula:  $C_4H_{10}O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	4	3		
5	A	1	Total	C	O	0	0
			7	4	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	178	Total 178	O 178	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pyruvate kinase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	89.20Å 109.67Å 134.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.74 – 1.90 19.93 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.2 (19.74-1.90) 99.1 (19.93-1.80)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.96 (at 1.80Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158, PHENIX 1.19.2_4158	Depositor
R, $R_{free}$	0.203 , 0.242 0.203 , 0.241	Depositor DCC
$R_{free}$ test set	3624 reflections (6.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	47.0	Xtrriage
Anisotropy	0.113	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3988	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.96% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PEG, MG, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.34	0/3830	0.54	0/5170

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	470	ARG	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3782	0	3944	41	3
2	A	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	1	0	0	0	0
4	A	12	0	16	0	0
5	A	14	0	20	6	2
6	A	178	0	0	7	0
All	All	3988	0	3980	41	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:449:ASN:ND2	6:A:701:HOH:O	2.20	0.73
1:A:424:ARG:HE	5:A:606:PEG:H31	1.55	0.71
1:A:313:THR:HG22	1:A:314:GLN:HG3	1.73	0.69
1:A:113:LEU:HD13	1:A:118:VAL:HG13	1.82	0.61
1:A:388:GLU:HG2	1:A:390:PRO:HD2	1.82	0.61
1:A:376:TYR:OH	1:A:452:ARG:NH1	2.35	0.59
1:A:147:LYS:NZ	6:A:706:HOH:O	2.30	0.59
1:A:424:ARG:HD3	5:A:606:PEG:H21	1.87	0.56
1:A:319:MET:SD	1:A:359:ALA:HA	2.48	0.54
1:A:10:ALA:N	6:A:709:HOH:O	2.41	0.54
1:A:436:LEU:HD11	1:A:457:ILE:HG13	1.92	0.52
1:A:165:VAL:HG22	1:A:185:ALA:HB3	1.93	0.51
1:A:159:LEU:HD21	5:A:604:PEG:H42	1.92	0.50
1:A:355:PHE:HB3	1:A:358:GLU:HB2	1.94	0.50
1:A:45:LEU:HD11	1:A:66:CYS:HB3	1.93	0.50
1:A:284:MET:HA	5:A:604:PEG:H31	1.94	0.49
1:A:280:GLY:HA3	1:A:313:THR:HG21	1.94	0.49
1:A:104:LYS:HE3	1:A:107:GLU:CD	2.34	0.48
1:A:266:ASP:OD1	1:A:301:LYS:HE3	2.13	0.48
1:A:380:TYR:OH	1:A:399:ARG:HG2	2.12	0.48
1:A:146:LYS:NZ	6:A:708:HOH:O	2.39	0.48
1:A:371:GLU:HA	1:A:374:ILE:HD12	1.96	0.47
1:A:254:PRO:HD2	1:A:274:GLY:O	2.15	0.47
1:A:83:ASN:O	1:A:87:LYS:HG3	2.16	0.45
1:A:138:GLU:H	1:A:138:GLU:CD	2.19	0.45
1:A:308:PRO:HG3	5:A:606:PEG:H32	1.97	0.45
1:A:439:SER:HB3	1:A:445:VAL:HG23	1.98	0.45
1:A:56:VAL:HG21	1:A:91:LEU:HD12	2.00	0.44
1:A:290:LYS:HG3	1:A:293:LEU:HD12	1.99	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:416:LEU:HD11	1:A:469:ILE:HG12	2.00	0.44
1:A:50:LYS:HE2	1:A:80:GLU:OE1	2.17	0.43
1:A:494:ILE:HA	1:A:503:ASN:OD1	2.19	0.43
1:A:424:ARG:NE	5:A:606:PEG:H31	2.28	0.43
1:A:121:LYS:HE3	6:A:704:HOH:O	2.20	0.42
1:A:319:MET:HE1	1:A:362:ILE:HG21	2.02	0.42
1:A:148:LEU:HB3	1:A:149:PRO:HD3	2.01	0.41
1:A:128:LEU:HA	1:A:141:ILE:O	2.21	0.41
1:A:218:ILE:HB	1:A:219:PRO:HD3	2.03	0.41
1:A:157:ILE:O	1:A:196:PRO:HD2	2.20	0.41
1:A:492:HIS:HB2	6:A:844:HOH:O	2.21	0.41
1:A:267:LYS:HD3	6:A:843:HOH:O	2.21	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:324:ARG:NH1	5:A:604:PEG:O4[2_555]	1.30	0.90
1:A:324:ARG:NH1	5:A:604:PEG:C4[2_555]	1.77	0.43
1:A:133:GLU:OE2	1:A:470:ARG:NH1[8_555]	1.82	0.38

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	494/519 (95%)	482 (98%)	11 (2%)	1 (0%)	47 38

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	313	THR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	426/446 (96%)	422 (99%)	4 (1%)	<a href="#">78</a> <a href="#">79</a>

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	119	HIS
1	A	137	ASP
1	A	290	LYS
1	A	300	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	84	ASN
1	A	480	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GOL	A	603	-	5,5,5	0.82	0	5,5,5	1.03	0
4	GOL	A	605	-	5,5,5	0.89	0	5,5,5	0.90	0
5	PEG	A	604	-	6,6,6	0.49	0	5,5,5	0.23	0
5	PEG	A	606	-	6,6,6	0.53	0	5,5,5	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	603	-	-	1/4/4/4	-
4	GOL	A	605	-	-	3/4/4/4	-
5	PEG	A	604	-	-	1/4/4/4	-
5	PEG	A	606	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	605	GOL	O1-C1-C2-C3
5	A	606	PEG	O1-C1-C2-O2
4	A	605	GOL	O1-C1-C2-O2
5	A	606	PEG	O2-C3-C4-O4
4	A	603	GOL	C1-C2-C3-O3
4	A	605	GOL	O2-C2-C3-O3
5	A	606	PEG	C4-C3-O2-C2
5	A	604	PEG	C1-C2-O2-C3

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	604	PEG	2	2
5	A	606	PEG	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	497/519 (95%)	0.84	77 (15%) <b>2</b> <b>2</b>	32, 49, 75, 103	0

All (77) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	10	ALA	13.5
1	A	11	ALA	8.5
1	A	389	THR	7.2
1	A	501	GLY	6.9
1	A	18	ALA	6.8
1	A	502	THR	6.7
1	A	512	LEU	6.5
1	A	494	ILE	5.2
1	A	390	PRO	5.1
1	A	115	ASN	4.5
1	A	493	GLY	4.2
1	A	128	LEU	4.2
1	A	388	GLU	3.9
1	A	414	ILE	3.9
1	A	245	PRO	3.7
1	A	45	LEU	3.7
1	A	160	ILE	3.6
1	A	256	ILE	3.5
1	A	378	LEU	3.5
1	A	15	MET	3.5
1	A	230	ILE	3.4
1	A	126	LEU	3.4
1	A	322	ASN	3.3
1	A	17	SER	3.2
1	A	119	HIS	3.2
1	A	101	LEU	3.2
1	A	316	LEU	3.2

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	352	GLY	3.1
1	A	93	PRO	3.1
1	A	355	PHE	3.1
1	A	16	GLN	3.0
1	A	72	HIS	2.9
1	A	12	GLY	2.9
1	A	31	ASN	2.9
1	A	77	ASP	2.8
1	A	354	LYS	2.8
1	A	248	ARG	2.8
1	A	385	ASN	2.8
1	A	80	GLU	2.8
1	A	135	LEU	2.7
1	A	142	ALA	2.7
1	A	246	ARG	2.7
1	A	74	SER	2.7
1	A	87	LYS	2.7
1	A	464	GLY	2.6
1	A	76	GLU	2.6
1	A	511	GLU	2.5
1	A	158	ILE	2.5
1	A	360	VAL	2.5
1	A	53	GLU	2.4
1	A	366	ILE	2.4
1	A	330	VAL	2.4
1	A	195	LEU	2.4
1	A	68	PHE	2.4
1	A	116	LYS	2.4
1	A	415	ALA	2.4
1	A	14	SER	2.4
1	A	313	THR	2.4
1	A	489	ILE	2.4
1	A	193	MET	2.3
1	A	437	ALA	2.3
1	A	66	CYS	2.2
1	A	323	PRO	2.2
1	A	19	ALA	2.2
1	A	43	CYS	2.2
1	A	52	VAL	2.2
1	A	391	ILE	2.1
1	A	367	CYS	2.1
1	A	114	LYS	2.1

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	48	ALA	2.1
1	A	117	GLU	2.1
1	A	154	PRO	2.1
1	A	32	VAL	2.0
1	A	337	VAL	2.0
1	A	227	ALA	2.0
1	A	162	ASP	2.0
1	A	438	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	A	605	6/6	0.41	0.34	65,79,89,93	0
5	PEG	A	604	7/7	0.70	0.17	62,73,82,85	0
4	GOL	A	603	6/6	0.74	0.20	48,71,81,82	0
5	PEG	A	606	7/7	0.83	0.20	37,49,66,70	0
2	MG	A	601	1/1	0.93	0.12	51,51,51,51	0
3	K	A	602	1/1	0.94	0.19	68,68,68,68	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around MG A 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.