

# Full wwPDB X-ray Structure Validation Report (i)

#### Mar 15, 2022 – 09:13 pm GMT

PDB ID : 7Z6R

Title: Psychrobacter arcticus ATPPRT (HisGZ) R56A mutant bound to ATP and

PRPP

Authors: Alphey, M.S.; Fisher, G.; da Silva, R.G.

Deposited on : 2022-03-14

Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.27

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0267

CCP4 : 7.1.010 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.27

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\mathring{A})}) \end{array}$
$R_{free}$	130704	1284 (2.56-2.52)
Clashscore	141614	1332 (2.56-2.52)
Ramachandran outliers	138981	1315 (2.56-2.52)
Sidechain outliers	138945	1315 (2.56-2.52)
RSRZ outliers	127900	1272 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	388	85%	9% • 5%
1	В	388	86%	10% • •
2	С	232	78% 9%	• 11%
2	D	232	81%	% 11%



# 2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 8522 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called ATP phosphoribosyltransferase regulatory subunit.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	A	368	Total 2633	C 1652	N 473	O 498	S 10	0	0	0
1	В	377	Total 2809	C 1786	N 499	O 513	S 11	0	1	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP Q4FTX3
В	0	GLY	-	expression tag	UNP Q4FTX3

• Molecule 2 is a protein called ATP phosphoribosyltransferase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	С	206	Total	С	N	О	S	0	0	0
	200	1478	938	260	276	4	U	U	0	
9	D	206	Total	С	N	О	S	0	0	0
	2 D	206	1480	946	253	277	4	0		

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	0	GLY	-	expression tag	UNP Q4FQF7
С	56	ALA	ARG	engineered mutation	UNP Q4FQF7
D	0	GLY	-	expression tag	UNP Q4FQF7
D	56	ALA	ARG	engineered mutation	UNP Q4FQF7


• Molecule 3 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula:  $C_{10}H_{16}N_5O_{13}P_3$ ) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	C	1	Total	С	N	О	Р	0	0
	1	31	10	5	13	3	U		
9	D	1	Total	С	N	О	Р	0 0	
$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	1	31	10	5	13	3	U	0	

• Molecule 4 is 1-O-pyrophosphono-5-O-phosphono-alpha-D-ribofuranose (three-letter code: PRP) (formula:  $C_5H_{13}O_{14}P_3$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	С	1	Total 22	C 5		P 3	0	0



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	D	1	Total 22	C 5		P 3	0	0

#### • Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	8	Total O 8 8	0	0
5	В	4	Total O 4 4	0	0
5	С	3	Total O 3 3	0	0
5	D	1	Total O 1 1	0	0




## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ATP phosphoribosyltransferase regulatory subunit







# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	102.16Å 145.53Å 93.81Å	Donogiton
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 102.48° 90.00°	Depositor
Resolution (Å)	56.97 - 2.55	Depositor
Resolution (A)	56.97 - 2.55	EDS
% Data completeness	99.8 (56.97-2.55)	Depositor
(in resolution range)	99.8 (56.97-2.55)	EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.19 (at 2.55Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
D D.	0.233 , 0.275	Depositor
$R, R_{free}$	0.236 , $0.277$	DCC
$R_{free}$ test set	2156 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	64.7	Xtriage
Anisotropy	0.305	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	(Not available), (Not available)	EDS
L-test for twinning <sup>2</sup>	$  <  L  > = 0.50, < L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	8522	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.06% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ATP, PRP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI		RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.66	0/2682	0.78	0/3631	
1	В	0.68	0/2867	0.79	0/3894	
2	С	0.69	0/1500	0.79	0/2040	
2	D	0.68	0/1503	0.78	0/2047	
All	All	0.68	0/8552	0.78	0/11612	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	2633	0	2406	21	0
1	В	2809	0	2726	26	0
2	С	1478	0	1441	19	0
2	D	1480	0	1446	10	0
3	С	31	0	12	1	0
3	D	31	0	12	0	0
4	С	22	0	8	1	0
4	D	22	0	8	2	0
5	A	8	0	0	0	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	В	4	0	0	0	0
5	С	3	0	0	0	0
5	D	1	0	0	0	0
All	All	8522	0	8059	73	0

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 4.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A.11	A	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
2:C:117:LEU:CD1	2:C:177:VAL:HG12	2.26	0.66
1:B:275:ILE:HD11	1:B:282:LEU:HD22	1.78	0.65
1:B:82:LEU:HD21	1:B:290:GLY:HA2	1.80	0.64
2:C:78:PRO:HG2	2:C:95:VAL:HG12	1.83	0.60
1:A:63:ILE:HD12	1:A:63:ILE:N	2.19	0.58
2:C:62:THR:HG22	2:C:67:VAL:O	2.03	0.58
2:C:46:VAL:HG13	2:C:62:THR:HG21	1.85	0.57
2:D:78:PRO:HG2	2:D:95:VAL:HG12	1.87	0.56
2:D:62:THR:HG22	2:D:67:VAL:O	2.06	0.55
1:A:62:ILE:HD13	1:A:73:ILE:HD11	1.88	0.54
2:C:117:LEU:HD11	2:C:177:VAL:HG12	1.90	0.53
1:B:155:ASP:OD1	1:B:254:SER:OG	2.26	0.52
2:C:165:ALA:HB3	2:C:166:PRO:HD3	1.92	0.52
2:D:165:ALA:HB3	2:D:166:PRO:HD3	1.92	0.52
1:A:102:VAL:HG23	1:A:116:PRO:HG2	1.91	0.52
1:A:30:THR:HB	1:B:352:ARG:HG2	1.93	0.51
1:B:153:HIS:CE1	1:B:276:ASN:O	2.65	0.50
2:C:107:LEU:HD12	2:C:204:VAL:HG21	1.94	0.50
2:C:130:ARG:NH2	2:C:172:ASP:OD1	2.43	0.50
2:C:116:LYS:HE2	2:C:118:MET:HE2	1.93	0.49
1:B:275:ILE:CD1	1:B:282:LEU:HD22	2.43	0.49
1:B:21:ARG:HD3	1:B:117:LEU:HD21	1.93	0.49
1:B:140:PHE:HZ	1:B:251:CYS:HG	1.59	0.49
1:A:287:ARG:NH1	1:A:303:GLN:HG2	2.28	0.48
2:C:179:ASP:N	4:C:402:PRP:O1P	2.46	0.48
1:B:136:ILE:HD12	1:B:245:LEU:HD21	1.95	0.48
1:A:42:GLU:HG2	1:B:2:LEU:CD2	2.43	0.48
1:B:60:PHE:CZ	1:B:105:THR:HA	2.48	0.48
2:C:177:VAL:HG21	3:C:401:ATP:C8	2.49	0.48
1:A:224:ASN:OD1	1:A:224:ASN:N	2.46	0.47



 $Continued\ from\ previous\ page...$ 

Continued from prev		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
2:D:217:GLU:HB3	2:D:218:PRO:HD3	1.96	0.47
1:A:60:PHE:CZ	1:A:105:THR:HA	2.48	0.47
1:B:102:VAL:HG23	1:B:116:PRO:HG2	1.95	0.47
1:A:102:VAL:CG2	1:A:116:PRO:HG2	2.43	0.47
1:B:28:LEU:HD13	1:B:97:CYS:HB2	1.96	0.47
1:A:156:LEU:HB2	1:A:255:ILE:HD13	1.97	0.47
2:C:48:LEU:CD2	2:C:60:PHE:HB3	2.45	0.47
2:C:46:VAL:HG12	2:C:46:VAL:O	2.15	0.46
1:A:129:ILE:HD12	1:A:237:GLU:HG2	1.97	0.46
1:B:124:PHE:HB3	1:B:302:ARG:HE	1.80	0.46
1:B:275:ILE:CG1	1:B:282:LEU:HD22	2.46	0.46
2:D:178:VAL:HG11	2:D:184:LEU:HD11	1.98	0.46
1:B:171:LEU:HD11	1:B:204:PHE:CZ	2.51	0.46
2:D:179:ASP:HB3	4:D:501:PRP:O2P	2.16	0.45
1:B:18:GLU:OE1	1:B:21:ARG:NH1	2.50	0.45
1:B:102:VAL:CG2	1:B:116:PRO:HG2	2.47	0.44
1:B:241:LEU:HD23	1:B:241:LEU:HA	1.83	0.44
1:B:333:PHE:HD1	1:B:344:LEU:HD23	1.83	0.44
1:A:165:LEU:CB	1:A:204:PHE:HE1	2.31	0.43
1:A:19:VAL:O	1:A:23:GLN:HG2	2.18	0.43
1:A:333:PHE:HD1	1:A:344:LEU:HD23	1.82	0.43
1:A:242:LYS:O	1:A:246:GLN:HG3	2.18	0.43
2:C:117:LEU:HD13	2:C:198:VAL:HG21	2.01	0.43
1:A:129:ILE:O	1:A:133:ILE:HD12	2.18	0.43
1:A:155:ASP:OD1	1:A:254:SER:OG	2.34	0.43
1:A:290:GLY:C	1:A:302:ARG:HB3	2.39	0.43
1:B:387:VAL:HG23	1:B:387:VAL:O	2.19	0.42
1:A:42:GLU:HB2	1:A:47:LEU:HB2	2.00	0.42
1:B:19:VAL:O	1:B:23:GLN:HG2	2.20	0.42
1:B:214:ILE:HB	1:B:255:ILE:HD12	2.02	0.42
2:C:48:LEU:HD21	2:C:60:PHE:HB3	2.00	0.42
2:D:180:THR:N	4:D:501:PRP:O2P	2.41	0.42
1:B:58:GLN:OE1	1:B:104:HIS:HB3	2.19	0.41
1:B:129:ILE:HD12	1:B:237:GLU:HA	2.03	0.41
2:D:145:TYR:CG	2:D:195:ILE:HG12	2.55	0.41
2:C:118:MET:HE3	2:C:178:VAL:HG21	2.02	0.41
2:C:83:HIS:CE1	2:D:156:ILE:CD1	3.03	0.41
2:C:145:TYR:CG	2:C:195:ILE:HG12	2.56	0.41
2:C:117:LEU:HD23	2:C:141:VAL:HG11	2.03	0.40
2:D:77:VAL:HB	2:D:78:PRO:HD3	2.03	0.40
1:A:326:VAL:HA	1:A:373:HIS:O	2.22	0.40



Atom-1	Atom-1 Atom-2		$egin{aligned}  ext{Clash} \  ext{overlap } ( ext{Å}) \end{aligned}$
1:A:23:GLN:HA	1:A:23:GLN:OE1	2.21	0.40
1:B:341:ARG:O	1:B:345:LEU:HG	2.21	0.40

There are no symmetry-related clashes.

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percen	${f tiles}$
1	A	358/388~(92%)	346 (97%)	12 (3%)	0	100	100
1	В	374/388 (96%)	363 (97%)	11 (3%)	0	100	100
2	$\mathbf{C}$	202/232~(87%)	189 (94%)	13 (6%)	0	100	100
2	D	202/232~(87%)	190 (94%)	12 (6%)	0	100	100
All	All	1136/1240 (92%)	1088 (96%)	48 (4%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	otameric Outliers		Percentiles		
1	A	$250/329\ (76\%)$	242 (97%)	8 (3%)	39 53	3		
1	В	283/329~(86%)	276 (98%)	7 (2%)	47 62	2		
2	С	144/194~(74%)	140 (97%)	4 (3%)	43 58	3		



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
2	D	146/194 (75%)	142 (97%)	4 (3%)	44	59	
All	All	823/1046 (79%)	800 (97%)	23 (3%)	43	58	

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	49	SER
1	A	113	SER
1	A	143	ILE
1	A	164	ARG
1	A	196	GLN
1	A	224	ASN
1	A	302	ARG
1	A	331	ASP
1	В	113	SER
1	В	129	ILE
1	В	153	HIS
1	В	205	TYR
1	В	241	LEU
1	В	261	SER
1	В	275	ILE
2	C C C	48	LEU
2	С	88	PHE
2	С	117	LEU
2	С	178	VAL
2	D	88	PHE
2	D	132	LEU
2	D	178	VAL
2	D	227	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	185	ASN
1	A	196	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.



### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	e Chain	Res	Link	Bond lengths			Bond angles		
MIOI	Type	Chain	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	ATP	D	502	-	26,33,33	0.66	0	31,52,52	0.75	1 (3%)
3	ATP	С	401	-	26,33,33	0.65	0	31,52,52	0.86	2 (6%)
4	PRP	D	501	-	19,22,22	0.62	0	33,35,35	1.20	2 (6%)
4	PRP	С	402	-	19,22,22	0.58	0	33,35,35	1.09	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	$\operatorname{Res}$	Link	Chirals	Torsions	Rings
3	ATP	D	502	-	-	1/18/38/38	0/3/3/3
3	ATP	С	401	-	-	6/18/38/38	0/3/3/3
4	PRP	D	501	-	-	8/16/33/33	0/1/1/1
4	PRP	C	402	-	-	1/16/33/33	0/1/1/1

There are no bond length outliers.

All (8) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$Ideal(^{o})$
4	D	501	PRP	PA-O3A-PB	-2.89	122.91	132.83
4	D	501	PRP	O1-C1-C2	2.63	111.17	106.72
4	С	402	PRP	PA-O3A-PB	-2.49	124.30	132.83
3	С	401	ATP	C5-C6-N6	2.25	123.77	120.35
3	D	502	ATP	C5-C6-N6	2.24	123.76	120.35
3	С	401	ATP	C3'-C2'-C1'	2.16	104.24	100.98
4	С	402	PRP	O1-C1-C2	2.07	110.22	106.72
4	С	402	PRP	C1-C2-C3	2.01	104.84	102.30

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	С	401	ATP	C3'-C4'-C5'-O5'
4	D	501	PRP	C5-O5-P-O1P
4	D	501	PRP	C5-O5-P-O2P
4	D	501	PRP	C5-O5-P-O3P
4	D	501	PRP	O4-C4-C5-O5
4	D	501	PRP	C1-O1-PA-O3A
3	С	401	ATP	O4'-C4'-C5'-O5'
4	С	402	PRP	PA-O3A-PB-O3B
4	D	501	PRP	PB-O3A-PA-O2A
3	С	401	ATP	PB-O3B-PG-O1G
4	D	501	PRP	PB-O3A-PA-O1A
3	С	401	ATP	PB-O3B-PG-O2G
3	С	401	ATP	PB-O3B-PG-O3G
3	С	401	ATP	PG-O3B-PB-O2B
3	D	502	ATP	PB-O3B-PG-O1G
4	D	501	PRP	C2-C1-O1-PA

There are no ring outliers.


3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	С	401	ATP	1	0
4	D	501	PRP	2	0
4	С	402	PRP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is




within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











# 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q<0.9
1	A	368/388 (94%)	0.15	6 (1%) 72 78	52, 76, 123, 142	0
1	В	377/388 (97%)	0.07	0 100 100	45, 64, 82, 98	0
2	С	206/232 (88%)	0.15	4 (1%) 66 73	70, 87, 107, 123	0
2	D	206/232 (88%)	0.41	14 (6%) 17 20	65, 86, 113, 142	0
All	All	1157/1240 (93%)	0.17	24 (2%) 63 70	45, 76, 113, 142	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	28	LEU	4.0
2	D	33	ILE	3.7
1	A	263	TYR	3.6
2	D	37	THR	3.0
2	D	90	VAL	2.9
1	A	378	ALA	2.8
2	С	81	VAL	2.7
2	С	23	GLY	2.7
2	D	227	ILE	2.6
1	A	387	VAL	2.6
2	С	101	ALA	2.6
2	D	54	ALA	2.6
2	D	46	VAL	2.5
2	D	191	ALA	2.5
2	D	121	GLY	2.5
1	A	301	PRO	2.4
2	D	74	ALA	2.3
1	A	124	PHE	2.2
1	A	82	LEU	2.2
2	D	175	VAL	2.1
2	D	122	VAL	2.1



Mol	Chain	Res	Type	RSRZ
2	D	184	LEU	2.1
2	С	46	VAL	2.1
2	D	52	PRO	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
3	ATP	D	502	31/31	0.81	0.25	93,95,121,122	31
3	ATP	С	401	31/31	0.82	0.18	89,92,108,111	31
4	PRP	D	501	22/22	0.86	0.12	93,113,139,140	0
4	PRP	С	402	22/22	0.90	0.11	94,106,111,115	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




## Electron density around ATP D 502:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










#### Electron density around ATP C 401:


 $2 \text{mF}_o\text{-DF}_c$  (at 0.7 rmsd) in gray  $\text{mF}_o\text{-DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)














 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)









# 6.5 Other polymers (i)

There are no such residues in this entry.

