



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 14, 2023 – 12:28 PM JST

PDB ID : 5Z9G
Title : Crystal structure of KAI2
Authors : Kim, K.L.; Cha, J.S.; Soh, M.S.; Cho, H.S.
Deposited on : 2018-02-03
Resolution : 1.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	270	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2294 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable esterase KAI2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	267	2065	1316	344	395	10	0	0	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	229	Total	O	0	0
			229	229		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Probable esterase KAI2



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	50.60Å 55.78Å 53.03Å 90.00° 116.07° 90.00°	Depositor
Resolution (Å)	47.64 – 1.49 27.89 – 1.49	Depositor EDS
% Data completeness (in resolution range)	99.4 (47.64-1.49) 99.3 (27.89-1.49)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	13.04 (at 1.49Å)	Xtrriage
Refinement program	REFMAC 5.5.0110	Depositor
R, R_{free}	0.136 , 0.177 0.146 , 0.182	Depositor DCC
R_{free} test set	2183 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	11.0	Xtrriage
Anisotropy	0.128	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 46.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.018 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2294	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.13	8/2111 (0.4%)	1.14	16/2876 (0.6%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	226	GLU	CG-CD	10.81	1.68	1.51
1	A	176	ARG	CZ-NH2	7.74	1.43	1.33
1	A	22	LEU	CG-CD1	-6.41	1.28	1.51
1	A	252	SER	CB-OG	-6.30	1.34	1.42
1	A	262	ARG	CD-NE	-5.39	1.37	1.46
1	A	69	ASN	CB-CG	-5.22	1.39	1.51
1	A	182	ARG	CD-NE	-5.14	1.37	1.46
1	A	71	GLU	CB-CG	-5.05	1.42	1.52

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	262	ARG	NE-CZ-NH2	-17.54	111.53	120.30
1	A	182	ARG	NE-CZ-NH1	14.11	127.35	120.30
1	A	262	ARG	NE-CZ-NH1	12.88	126.74	120.30
1	A	182	ARG	NE-CZ-NH2	-10.69	114.95	120.30
1	A	176	ARG	NE-CZ-NH1	-6.99	116.81	120.30
1	A	63	ASP	CB-CG-OD1	6.79	124.42	118.30
1	A	22	LEU	CB-CG-CD2	6.73	122.44	111.00
1	A	15	SER	C-N-CA	-6.71	108.20	122.30
1	A	138	ASP	CB-CG-OD1	6.60	124.24	118.30
1	A	199	ARG	NE-CZ-NH2	-6.44	117.08	120.30
1	A	176	ARG	NE-CZ-NH2	6.18	123.39	120.30
1	A	63	ASP	CB-CG-OD2	-5.56	113.29	118.30
1	A	182	ARG	CD-NE-CZ	5.43	131.21	123.60
1	A	45	ARG	NE-CZ-NH1	-5.13	117.73	120.30
1	A	226	GLU	CA-CB-CG	5.12	124.67	113.40
1	A	76	ASP	CB-CG-OD1	5.08	122.87	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2065	0	2015	23	0
2	A	229	0	0	7	0
All	All	2294	0	2015	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:182:ARG:HD2	1:A:184:ASP:OD1	1.51	1.07
1:A:136:GLN:HE22	1:A:195:GLN:HE22	1.05	0.98
1:A:182:ARG:CD	1:A:184:ASP:OD1	2.16	0.93
1:A:69:ASN:HB2	2:A:372:HOH:O	1.76	0.84
1:A:16:GLY:HA3	2:A:460:HOH:O	1.96	0.65
1:A:136:GLN:NE2	1:A:195:GLN:HE22	1.87	0.65
1:A:248:PRO:HB2	1:A:256:VAL:HG21	1.83	0.61
1:A:213:GLN:HE21	1:A:225:SER:HB3	1.70	0.57
1:A:136:GLN:HE22	1:A:195:GLN:NE2	1.88	0.55
1:A:25:GLY:H	1:A:94:HIS:HD2	1.54	0.54
1:A:182:ARG:HD3	1:A:184:ASP:OD1	2.02	0.54
1:A:210:HIS:HD2	1:A:263:HIS:ND1	2.06	0.54
1:A:94:HIS:HE1	1:A:246:HIS:O	1.91	0.53
1:A:71:GLU:HG2	1:A:75:PHE:CZ	2.51	0.45
1:A:69:ASN:CG	2:A:372:HOH:O	2.54	0.45
1:A:210:HIS:HE1	2:A:503:HOH:O	1.98	0.45
1:A:201:ILE:C	1:A:201:ILE:HD12	2.40	0.42
1:A:69:ASN:CB	2:A:372:HOH:O	2.49	0.42
1:A:71:GLU:HG2	1:A:75:PHE:CE2	2.55	0.41
1:A:226:GLU:CG	2:A:315:HOH:O	2.68	0.41
1:A:22:LEU:HD13	1:A:92:VAL:HB	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:30:GLN:NE2	2:A:304:HOH:O	2.53	0.41
1:A:15:SER:O	1:A:16:GLY:C	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	265/270 (98%)	261 (98%)	4 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	231/235 (98%)	229 (99%)	2 (1%)	78 61

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	LEU
1	A	142	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	30	GLN
1	A	94	HIS
1	A	136	GLN
1	A	210	HIS
1	A	213	GLN
1	A	231	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	267/270 (98%)	0.06	14 (5%) 27 30	6, 10, 23, 30	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	143	PHE	5.6
1	A	15	SER	4.5
1	A	16	GLY	4.5
1	A	146	ILE	3.8
1	A	17	GLU	3.5
1	A	148	SER	3.5
1	A	166	MET	3.1
1	A	147	ARG	3.1
1	A	167	ASP	2.6
1	A	141	GLN	2.6
1	A	142	LEU	2.4
1	A	165	ASP	2.4
1	A	145	ALA	2.3
1	A	144	GLU	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.