



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 12, 2021 – 08:31 AM EDT

PDB ID : 1ZDT  
Title : The Crystal Structure of Human Steroidogenic Factor-1  
Authors : Wang, W.; Zhang, C.; Marimuthu, A.; Krupka, H.I.; Tabrizizad, M.; Shelloe, R.; Mehra, U.; Eng, K.; Nguyen, H.; Settachatgul, C.; Powell, B.; Milburn, M.V.; West, B.L.  
Deposited on : 2005-04-14  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	241	
1	B	241	
2	P	12	
2	Q	12	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4320 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Steroidogenic factor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	233	1873	1202	320	339	12	0	0	0
1	B	239	1922	1231	330	349	12	0	0	0


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	247	SER	CYS	engineered mutation	UNP Q13285
A	412	SER	CYS	engineered mutation	UNP Q13285
B	247	SER	CYS	engineered mutation	UNP Q13285
B	412	SER	CYS	engineered mutation	UNP Q13285

- Molecule 2 is a protein called Nuclear receptor coactivator 2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	P	12	102	65	17	20	0	0	0
2	Q	11	94	61	16	17	0	0	0

- Molecule 3 is DI-PALMITOYL-3-SN-PHOSPHATIDYLETHANOLAMINE (three-letter code: PEF) (formula: C<sub>37</sub>H<sub>74</sub>NO<sub>8</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	47	37	1	8	1	0	0
3	B	1	47	37	1	8	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	116	116	116	0	0
4	B	114	114	114	0	0
4	P	3	3	3	0	0
4	Q	2	2	2	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Steroidogenic factor 1




- Molecule 1: Steroidogenic factor 1



- Molecule 2: Nuclear receptor coactivator 2



- Molecule 2: Nuclear receptor coactivator 2



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.60Å 73.60Å 195.68Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.10 97.84 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.3 (50.00-2.10) 98.4 (97.84-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.11	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.60 (at 2.10Å)	Xtrriage
Refinement program	REFMAC 5.1.25	Depositor
R, $R_{free}$	0.216 , 0.265 0.221 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	40.5	Xtrriage
Anisotropy	0.298	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 44.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.036 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4320	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.60% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: PEF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.74	1/1904 (0.1%)	0.86	3/2575 (0.1%)
1	B	0.75	0/1955	0.84	6/2646 (0.2%)
2	P	1.16	2/102 (2.0%)	1.07	1/136 (0.7%)
2	Q	0.59	0/94	0.91	0/125
All	All	0.76	3/4055 (0.1%)	0.86	10/5482 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	P	751	LYS	CA-C	5.50	1.67	1.52
2	P	752	ASP	N-CA	5.17	1.56	1.46
1	A	357	GLN	CD-OE1	5.04	1.35	1.24

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	238	ASP	CB-CG-OD2	6.88	124.49	118.30
1	A	278	ASP	CB-CG-OD2	6.29	123.96	118.30
1	B	392	ASP	CB-CG-OD2	5.91	123.62	118.30
1	B	380	ASP	CB-CG-OD2	5.85	123.56	118.30
1	A	364	ASP	CB-CG-OD2	5.75	123.47	118.30
1	B	406	LEU	CA-CB-CG	5.41	127.74	115.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	403	ASP	CB-CG-OD2	5.40	123.16	118.30
1	B	278	ASP	CB-CG-OD2	5.37	123.13	118.30
1	B	309	ASP	CB-CG-OD2	5.09	122.88	118.30
2	P	752	ASP	N-CA-C	5.04	124.62	111.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	244	ILE	Peptide
1	A	257	ASP	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1873	0	1918	15	1
1	B	1922	0	1969	20	0
2	P	102	0	103	3	0
2	Q	94	0	99	2	0
3	A	47	0	73	2	0
3	B	47	0	73	1	0
4	A	116	0	0	3	0
4	B	114	0	0	2	1
4	P	3	0	0	0	0
4	Q	2	0	0	0	0
All	All	4320	0	4235	40	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:453:ILE:O	1:A:457:GLN:HG2	1.88	0.73

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:236:ASP:O	1:B:240:VAL:HG12	1.97	0.64
1:A:304:GLU:HG2	4:A:1034:HOH:O	2.00	0.61
1:B:285:VAL:HG11	1:B:393:ALA:HB2	1.84	0.59
1:A:334:THR:O	1:A:338:THR:HG23	2.03	0.59
1:A:238:ASP:HA	4:A:1025:HOH:O	2.03	0.58
1:A:347:LEU:HD11	1:A:433:ALA:HA	1.85	0.58
2:Q:750:ASP:O	2:Q:751:LYS:C	2.42	0.56
1:A:288:GLU:HG3	4:A:1035:HOH:O	2.04	0.56
1:B:338:THR:HG23	1:B:339:GLN:OE1	2.05	0.56
1:A:324:LEU:HD23	1:A:330:GLU:HG2	1.87	0.56
1:B:402:LEU:O	1:B:406:LEU:HD22	2.06	0.54
1:B:277:VAL:HG11	2:Q:748:LEU:HB3	1.89	0.54
1:A:402:LEU:O	1:A:406:LEU:HB2	2.07	0.53
1:B:225:GLU:O	1:B:229:GLN:HG3	2.10	0.52
1:B:347:LEU:HD13	1:B:432:GLN:HB3	1.90	0.52
1:B:379:LEU:HD11	1:B:383:PHE:CE1	2.46	0.51
2:P:741:GLU:OE2	2:P:746:ARG:NH1	2.45	0.50
1:A:347:LEU:HD13	3:A:1001:PEF:H212	1.94	0.49
1:A:323:ILE:O	1:A:330:GLU:HA	2.12	0.49
1:B:347:LEU:HD21	1:B:433:ALA:HA	1.95	0.49
1:B:305:LEU:HD13	1:B:375:ILE:CG2	2.43	0.49
1:B:255:ARG:HD2	4:B:1060:HOH:O	2.13	0.48
1:A:351:ALA:HA	1:A:429:LEU:HD21	1.97	0.46
1:B:338:THR:CG2	1:B:339:GLN:OE1	2.64	0.45
1:B:386:ASN:OD1	1:B:388:ILE:HG22	2.17	0.44
2:P:747:TYR:OH	2:P:752:ASP:HB3	2.18	0.44
1:B:381:LEU:HD21	1:B:391:LYS:HB2	2.00	0.43
1:B:458:ALA:O	1:B:459:LYS:CB	2.66	0.43
1:B:241:ARG:HG2	1:B:245:LEU:HD22	2.00	0.43
1:A:310:HIS:CD2	3:A:1001:PEF:H241	2.53	0.43
1:A:237:GLU:O	1:A:238:ASP:HB2	2.18	0.43
1:B:305:LEU:HD13	1:B:375:ILE:HG21	2.00	0.42
1:B:252:THR:HG22	1:B:253:LYS:N	2.35	0.42
1:B:289:LEU:HD12	1:B:389:LEU:HD23	2.00	0.42
1:A:307:VAL:O	1:A:311:ILE:HG12	2.19	0.41
2:P:751:LYS:O	2:P:752:ASP:HB2	2.20	0.41
1:B:383:PHE:CD2	1:B:383:PHE:N	2.88	0.41
3:B:1002:PEF:H52	4:B:1115:HOH:O	2.21	0.40
1:A:333:LEU:HD13	1:A:333:LEU:HA	1.89	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:417:GLN:OE1	4:B:1003:HOH:O[5_555]	2.07	0.13

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	229/241 (95%)	220 (96%)	8 (4%)	1 (0%)	34	32
1	B	237/241 (98%)	233 (98%)	4 (2%)	0	100	100
2	P	10/12 (83%)	9 (90%)	0	1 (10%)	0	0
2	Q	9/12 (75%)	8 (89%)	1 (11%)	0	100	100
All	All	485/506 (96%)	470 (97%)	13 (3%)	2 (0%)	34	32

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	459	LYS
2	P	751	LYS

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	208/216 (96%)	183 (88%)	25 (12%)	5	2
1	B	214/216 (99%)	188 (88%)	26 (12%)	5	2
2	P	11/11 (100%)	8 (73%)	3 (27%)	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	Q	10/11 (91%)	6 (60%)	4 (40%)	0	0
All	All	443/454 (98%)	385 (87%)	58 (13%)	4	2

All (58) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	222	ASN
1	A	239	GLN
1	A	243	ARG
1	A	257	ASP
1	A	258	GLN
1	A	264	LEU
1	A	267	ARG
1	A	274	ILE
1	A	288	GLU
1	A	289	LEU
1	A	315	VAL
1	A	324	LEU
1	A	342	SER
1	A	347	LEU
1	A	349	LEU
1	A	357	GLN
1	A	363	LEU
1	A	371	LEU
1	A	402	LEU
1	A	412	SER
1	A	423	LEU
1	A	429	LEU
1	A	437	LEU
1	A	448	ARG
1	A	460	GLN
1	B	222	ASN
1	B	226	LEU
1	B	240	VAL
1	B	245	LEU
1	B	253	LYS
1	B	257	ASP
1	B	284	MET
1	B	305	LEU
1	B	332	GLU
1	B	333	LEU
1	B	338	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	343	LEU
1	B	346	SER
1	B	349	LEU
1	B	383	PHE
1	B	389	LEU
1	B	406	LEU
1	B	412	SER
1	B	414	ASP
1	B	415	LYS
1	B	417	GLN
1	B	423	LEU
1	B	440	LYS
1	B	448	ARG
1	B	451	LEU
1	B	459	LYS
2	P	741	GLU
2	P	746	ARG
2	P	752	ASP
2	Q	741	GLU
2	Q	744	LEU
2	Q	745	LEU
2	Q	751	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PEF	A	1001	-	46,46,46	1.05	4 (8%)	49,51,51	0.98	3 (6%)
3	PEF	B	1002	-	46,46,46	1.02	4 (8%)	49,51,51	0.98	4 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEF	A	1001	-	-	12/50/50/50	-
3	PEF	B	1002	-	-	13/50/50/50	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1001	PEF	O3-C30	3.76	1.44	1.33
3	B	1002	PEF	O3-C30	3.54	1.43	1.33
3	A	1001	PEF	O2-C10	3.54	1.44	1.34
3	B	1002	PEF	O2-C10	3.35	1.43	1.34
3	B	1002	PEF	C5-N	-2.48	1.26	1.46
3	B	1002	PEF	C44-C43	-2.46	1.34	1.51
3	A	1001	PEF	C44-C43	-2.44	1.34	1.51
3	A	1001	PEF	C5-N	-2.36	1.27	1.46

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1001	PEF	O3-C30-O5	-2.94	116.17	123.59
3	B	1002	PEF	O2-C10-C11	2.93	117.81	111.50
3	B	1002	PEF	O3-C30-C31	2.36	119.30	111.91

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1002	PEF	O2-C10-O4	-2.22	118.34	123.70
3	B	1002	PEF	O3-C30-O5	-2.12	118.24	123.59
3	A	1001	PEF	O2P-P-O1P	2.08	122.52	112.24
3	A	1001	PEF	O3-C30-C31	2.00	118.19	111.91

There are no chirality outliers.

All (25) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	1002	PEF	O4P-C4-C5-N
3	A	1001	PEF	C17-C18-C19-C20
3	B	1002	PEF	C17-C18-C19-C20
3	B	1002	PEF	C40-C41-C42-C43
3	A	1001	PEF	C16-C17-C18-C19
3	A	1001	PEF	C14-C15-C16-C17
3	B	1002	PEF	C16-C17-C18-C19
3	B	1002	PEF	C13-C14-C15-C16
3	A	1001	PEF	C19-C20-C21-C22
3	B	1002	PEF	C14-C15-C16-C17
3	A	1001	PEF	C39-C40-C41-C42
3	B	1002	PEF	C19-C20-C21-C22
3	B	1002	PEF	C21-C22-C23-C24
3	B	1002	PEF	C38-C39-C40-C41
3	B	1002	PEF	C15-C16-C17-C18
3	B	1002	PEF	C41-C42-C43-C44
3	B	1002	PEF	C36-C37-C38-C39
3	A	1001	PEF	C32-C33-C34-C35
3	A	1001	PEF	C13-C14-C15-C16
3	B	1002	PEF	C35-C36-C37-C38
3	A	1001	PEF	C31-C30-O3-C3
3	A	1001	PEF	O3-C30-C31-C32
3	A	1001	PEF	C37-C38-C39-C40
3	A	1001	PEF	O5-C30-O3-C3
3	A	1001	PEF	O5-C30-C31-C32

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1001	PEF	2	0
3	B	1002	PEF	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	233/241 (96%)	0.09	10 (4%) 35 41	16, 22, 32, 42	0
1	B	239/241 (99%)	0.01	7 (2%) 51 57	11, 23, 31, 36	0
2	P	12/12 (100%)	0.55	1 (8%) 11 14	16, 23, 27, 36	0
2	Q	11/12 (91%)	1.03	1 (9%) 9 12	20, 22, 26, 26	0
All	All	495/506 (97%)	0.08	19 (3%) 40 46	11, 23, 32, 42	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	256	PRO	7.4
1	B	254	SER	7.0
2	P	752	ASP	5.8
1	B	221	PRO	4.1
1	A	248	LEU	3.9
1	B	383	PHE	3.7
1	A	237	GLU	3.5
1	B	448	ARG	3.4
1	A	245	LEU	3.3
1	A	243	ARG	3.3
2	Q	747	TYR	3.2
1	A	410	PRO	2.9
1	A	411	HIS	2.6
1	A	460	GLN	2.4
1	B	253	LYS	2.2
1	B	255	ARG	2.2
1	A	240	VAL	2.2
1	B	258	GLN	2.1
1	A	459	LYS	2.0



## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	PEF	A	1001	47/47	0.95	0.13	19,24,27,28	0
3	PEF	B	1002	47/47	0.95	0.14	19,24,27,29	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around PEF A 1001:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around PEF B 1002:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.