



Full wwPDB EM Validation Report ⓘ

Dec 24, 2024 – 04:29 PM JST

PDB ID : 8ZDS
EMDB ID : EMD-60007
Title : Structure of the Salmonella flagellar MS-ring with C11 symmetry applied
Authors : Kinoshita, M.; Makino, F.; Miyata, T.; Imada, K.; Minamino, T.; Namba, K.
Deposited on : 2024-05-03
Resolution : 3.10 Å(reported)
Based on initial models : ., 7CIK, 7D84

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev113
MolProbity : 4.02b-467
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	560	
1	B	560	
1	C	560	
1	D	560	
1	E	560	
1	F	560	
1	G	560	
1	H	560	
1	I	560	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	J	560	
1	K	560	
1	L	560	
1	M	560	
1	N	560	
1	O	560	
1	P	560	
1	Q	560	
1	R	560	
1	S	560	
1	T	560	
1	U	560	
1	V	560	
1	W	560	
1	X	560	
1	Y	560	
1	Z	560	
1	a	560	
1	b	560	
1	c	560	
1	d	560	
1	e	560	
1	f	560	
1	g	560	

2 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 69498 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Flagellar M-ring protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	257	1977	1212	365	396	4	1	0
1	B	309	2363	1454	437	467	5	1	0
1	C	257	1978	1213	365	396	4	1	0
1	D	257	1977	1212	365	396	4	1	0
1	E	309	2363	1454	437	467	5	1	0
1	F	257	1978	1213	365	396	4	1	0
1	G	257	1977	1212	365	396	4	1	0
1	H	309	2363	1454	437	467	5	1	0
1	I	257	1978	1213	365	396	4	1	0
1	J	257	1977	1212	365	396	4	1	0
1	K	309	2363	1454	437	467	5	1	0
1	L	257	1978	1213	365	396	4	1	0
1	M	257	1977	1212	365	396	4	1	0
1	N	309	2363	1454	437	467	5	1	0
1	O	257	1978	1213	365	396	4	1	0
1	P	257	1977	1212	365	396	4	1	0
1	Q	309	2363	1454	437	467	5	1	0



Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
1	R	257	Total 1978	C 1213	N 365	O 396	S 4	1	0
1	S	257	Total 1977	C 1212	N 365	O 396	S 4	1	0
1	T	309	Total 2363	C 1454	N 437	O 467	S 5	1	0
1	U	257	Total 1978	C 1213	N 365	O 396	S 4	1	0
1	V	257	Total 1977	C 1212	N 365	O 396	S 4	1	0
1	W	309	Total 2363	C 1454	N 437	O 467	S 5	1	0
1	X	257	Total 1978	C 1213	N 365	O 396	S 4	1	0
1	Y	257	Total 1977	C 1212	N 365	O 396	S 4	1	0
1	Z	309	Total 2363	C 1454	N 437	O 467	S 5	1	0
1	a	257	Total 1978	C 1213	N 365	O 396	S 4	1	0
1	b	257	Total 1977	C 1212	N 365	O 396	S 4	1	0
1	c	309	Total 2363	C 1454	N 437	O 467	S 5	1	0
1	d	257	Total 1978	C 1213	N 365	O 396	S 4	1	0
1	e	257	Total 1977	C 1212	N 365	O 396	S 4	1	0
1	f	309	Total 2363	C 1454	N 437	O 467	S 5	1	0
1	g	257	Total 1978	C 1213	N 365	O 396	S 4	1	0




● Molecule 1: Flagellar M-ring protein






• Molecule 1: Flagellar M-ring protein




• Molecule 1: Flagellar M-ring protein






● Molecule 1: Flagellar M-ring protein






● Molecule 1: Flagellar M-ring protein






• Molecule 1: Flagellar M-ring protein






• Molecule 1: Flagellar M-ring protein





• Molecule 1: Flagellar M-ring protein






● Molecule 1: Flagellar M-ring protein






● Molecule 1: Flagellar M-ring protein







● Molecule 1: Flagellar M-ring protein






● Molecule 1: Flagellar M-ring protein





● Molecule 1: Flagellar M-ring protein





● Molecule 1: Flagellar M-ring protein



ASN	GLU	ALA	PRO	ILE	ALA	THR	PRO	THR	ASN	GLN	ASN	ALA	GLN	ASN	THR	THR	GLN	THR	GLN	THR	THR	THR	ASN	ASN	ASN	ASN	ALA	GLY	P355	R356	E367	V368	D369	R370	R373	H374	N378	V379	G380	D381	I382	E383	R384	L385	A388	V389	V390	V391	N392	Y393	K394	T395	L396				
ALA	ASP	GLY	LYS	P401	L402	P403	L404	T405	A406	D407	Q408	M409	K410	Q411	I412	E413	D414	L415	T416	R417	E418	A419	M420	G421	F422	S423	D424	K425	R426	G427	D428	T429	L430	M431	V432	V433	M434	S435	P436	F437	S438	ALA	VAL	ASP	ASN	THR	GLY	GLY	LEU	PRO	PHE	TRP	GLN	GLN	SER	PHE	ILE
ASP	GLN	LEU	GLN	LEU	ALA	ALA	GLY	ARG	TRP	LEU	LEU	VAL	LEU	VAL	VAL	VAL	TRP	ILE	GLN	TRP	ILE	TRP	ARG	LYS	ALA	VAL	ARG	VAL	PRO	GLN	LEU	THR	ARG	VAL	VAL	ALA	LEU	VAL	GLU	ALA	ILE	ALA	LYS	GLN	ALA	ALA	GLN	GLN	ASP	ALA	HIS	GLU					
GLN	LEU	GLN	GLN	ARG	ARG	ALA	ASN	GLN	ARG	GLY	ALA	GLU	VAL	VAL	MET	SER	GLN	ARG	ILE	ARG	GLU	MET	SER	ASP	ASN	ASP	PRO	PRO	LEU	THR	VAL	VAL	VAL	ALA	VAL	VAL	LEU	VAL	ILE	ALA	ARG	GLN	ALA	ALA	GLN	GLN	SER	ASP	ALA	HIS	GLU						

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C11	Depositor
Number of particles used	70250	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	JEOL CRYO ARM 300	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	50000	Depositor
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.052	Depositor
Minimum map value	-0.013	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.025	Depositor
Map size (\AA)	400.0, 400.0, 400.0	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.0, 1.0, 1.0	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.28	0/2005	0.52	0/2713
1	B	0.27	0/2397	0.51	0/3242
1	C	0.28	0/2006	0.54	0/2715
1	D	0.28	0/2005	0.53	0/2713
1	E	0.27	0/2397	0.51	0/3242
1	F	0.29	0/2006	0.53	0/2715
1	G	0.28	0/2005	0.53	0/2713
1	H	0.27	0/2397	0.52	0/3242
1	I	0.28	0/2006	0.54	0/2715
1	J	0.28	0/2005	0.51	0/2713
1	K	0.27	0/2397	0.51	0/3242
1	L	0.28	0/2006	0.53	0/2715
1	M	0.28	0/2005	0.54	0/2713
1	N	0.27	0/2397	0.51	0/3242
1	O	0.28	0/2006	0.52	0/2715
1	P	0.28	0/2005	0.52	0/2713
1	Q	0.27	0/2397	0.53	0/3242
1	R	0.28	0/2006	0.52	0/2715
1	S	0.27	0/2005	0.52	0/2713
1	T	0.27	0/2397	0.51	0/3242
1	U	0.28	0/2006	0.53	0/2715
1	V	0.28	0/2005	0.54	0/2713
1	W	0.27	0/2397	0.51	0/3242
1	X	0.28	0/2006	0.53	0/2715
1	Y	0.28	0/2005	0.54	0/2713
1	Z	0.27	0/2397	0.51	0/3242
1	a	0.28	0/2006	0.54	0/2715
1	b	0.28	0/2005	0.52	0/2713
1	c	0.27	0/2397	0.51	0/3242
1	d	0.28	0/2006	0.53	0/2715
1	e	0.28	0/2005	0.52	0/2713
1	f	0.27	0/2397	0.52	0/3242
1	g	0.28	0/2006	0.53	0/2715
All	All	0.28	0/70488	0.52	0/95370

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	250/560 (45%)	240 (96%)	10 (4%)	0	100	100
1	B	300/560 (54%)	284 (95%)	16 (5%)	0	100	100
1	C	250/560 (45%)	241 (96%)	9 (4%)	0	100	100
1	D	250/560 (45%)	239 (96%)	11 (4%)	0	100	100
1	E	300/560 (54%)	284 (95%)	16 (5%)	0	100	100
1	F	250/560 (45%)	240 (96%)	10 (4%)	0	100	100
1	G	250/560 (45%)	239 (96%)	11 (4%)	0	100	100
1	H	300/560 (54%)	285 (95%)	15 (5%)	0	100	100
1	I	250/560 (45%)	239 (96%)	11 (4%)	0	100	100
1	J	250/560 (45%)	238 (95%)	12 (5%)	0	100	100
1	K	300/560 (54%)	282 (94%)	18 (6%)	0	100	100
1	L	250/560 (45%)	240 (96%)	10 (4%)	0	100	100
1	M	250/560 (45%)	235 (94%)	15 (6%)	0	100	100
1	N	300/560 (54%)	283 (94%)	17 (6%)	0	100	100
1	O	250/560 (45%)	241 (96%)	9 (4%)	0	100	100
1	P	250/560 (45%)	241 (96%)	9 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	Q	300/560 (54%)	284 (95%)	16 (5%)	0	100	100
1	R	250/560 (45%)	240 (96%)	10 (4%)	0	100	100
1	S	250/560 (45%)	241 (96%)	9 (4%)	0	100	100
1	T	300/560 (54%)	285 (95%)	15 (5%)	0	100	100
1	U	250/560 (45%)	241 (96%)	9 (4%)	0	100	100
1	V	250/560 (45%)	240 (96%)	10 (4%)	0	100	100
1	W	300/560 (54%)	285 (95%)	15 (5%)	0	100	100
1	X	250/560 (45%)	240 (96%)	10 (4%)	0	100	100
1	Y	250/560 (45%)	239 (96%)	11 (4%)	0	100	100
1	Z	300/560 (54%)	285 (95%)	15 (5%)	0	100	100
1	a	250/560 (45%)	240 (96%)	10 (4%)	0	100	100
1	b	250/560 (45%)	239 (96%)	11 (4%)	0	100	100
1	c	300/560 (54%)	286 (95%)	14 (5%)	0	100	100
1	d	250/560 (45%)	242 (97%)	8 (3%)	0	100	100
1	e	250/560 (45%)	241 (96%)	9 (4%)	0	100	100
1	f	300/560 (54%)	284 (95%)	16 (5%)	0	100	100
1	g	250/560 (45%)	239 (96%)	11 (4%)	0	100	100
All	All	8800/18480 (48%)	8402 (96%)	398 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/467 (48%)	221 (99%)	2 (1%)	75	88
1	B	260/467 (56%)	258 (99%)	2 (1%)	79	89
1	C	223/467 (48%)	223 (100%)	0	100	100
1	D	223/467 (48%)	220 (99%)	3 (1%)	65	82

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	260/467 (56%)	259 (100%)	1 (0%)	89	94
1	F	223/467 (48%)	222 (100%)	1 (0%)	89	94
1	G	222/467 (48%)	221 (100%)	1 (0%)	86	92
1	H	260/467 (56%)	259 (100%)	1 (0%)	89	94
1	I	223/467 (48%)	223 (100%)	0	100	100
1	J	223/467 (48%)	220 (99%)	3 (1%)	65	82
1	K	260/467 (56%)	259 (100%)	1 (0%)	89	94
1	L	223/467 (48%)	223 (100%)	0	100	100
1	M	223/467 (48%)	220 (99%)	3 (1%)	65	82
1	N	260/467 (56%)	259 (100%)	1 (0%)	89	94
1	O	223/467 (48%)	223 (100%)	0	100	100
1	P	222/467 (48%)	220 (99%)	2 (1%)	75	88
1	Q	260/467 (56%)	258 (99%)	2 (1%)	79	89
1	R	223/467 (48%)	223 (100%)	0	100	100
1	S	223/467 (48%)	221 (99%)	2 (1%)	75	88
1	T	260/467 (56%)	258 (99%)	2 (1%)	79	89
1	U	223/467 (48%)	223 (100%)	0	100	100
1	V	223/467 (48%)	221 (99%)	2 (1%)	75	88
1	W	260/467 (56%)	259 (100%)	1 (0%)	89	94
1	X	223/467 (48%)	223 (100%)	0	100	100
1	Y	223/467 (48%)	221 (99%)	2 (1%)	75	88
1	Z	260/467 (56%)	259 (100%)	1 (0%)	89	94
1	a	223/467 (48%)	223 (100%)	0	100	100
1	b	223/467 (48%)	221 (99%)	2 (1%)	75	88
1	c	260/467 (56%)	259 (100%)	1 (0%)	89	94
1	d	223/467 (48%)	223 (100%)	0	100	100
1	e	223/467 (48%)	220 (99%)	3 (1%)	65	82
1	f	260/467 (56%)	258 (99%)	2 (1%)	79	89
1	g	223/467 (48%)	223 (100%)	0	100	100
All	All	7764/15411 (50%)	7723 (100%)	41 (0%)	85	92

All (41) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	186	LEU
1	A	236	LYS
1	B	236	LYS
1	B	425	LYS
1	D	186	LEU
1	D	236	LYS
1	D	425	LYS
1	E	236	LYS
1	F	425	LYS
1	G	236	LYS
1	H	236	LYS
1	J	186	LEU
1	J	236	LYS
1	J	425	LYS
1	K	236	LYS
1	M	186	LEU
1	M	236	LYS
1	M	425	LYS
1	N	236	LYS
1	P	236	LYS
1	P	425	LYS
1	Q	236	LYS
1	Q	425	LYS
1	S	186	LEU
1	S	236	LYS
1	T	236	LYS
1	T	425	LYS
1	V	186	LEU
1	V	236	LYS
1	W	236	LYS
1	Y	186	LEU
1	Y	236	LYS
1	Z	236	LYS
1	b	186	LEU
1	b	236	LYS
1	c	236	LYS
1	e	186	LEU
1	e	236	LYS
1	e	425	LYS
1	f	236	LYS
1	f	425	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	103	GLN
1	B	196	HIS
1	E	103	GLN
1	H	94	HIS
1	H	103	GLN
1	I	231	ASN
1	N	103	GLN
1	P	133	GLN
1	Q	103	GLN
1	T	103	GLN
1	Y	133	GLN
1	Z	103	GLN
1	Z	196	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-60007. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 200




Y Index: 200




Z Index: 200


6.2.2 Raw map



X Index: 200



Y Index: 200




Z Index: 200


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 149




Y Index: 251




Z Index: 183


6.3.2 Raw map



X Index: 149



Y Index: 251




Z Index: 216


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.025. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 54 nm³; this corresponds to an approximate mass of 49 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum i




*Reported resolution corresponds to spatial frequency of 0.323 \AA^{-1}

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.323 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.10	-	-
Author-provided FSC curve	3.03	3.43	3.06
Unmasked-calculated*	3.44	4.20	3.47

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.44 differs from the reported value 3.1 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-60007 and PDB model 8ZDS. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.025 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.025).





































































9.4 Atom inclusion [i](#)



At the recommended contour level, 20% of all backbone atoms, 14% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.025) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.1420	 0.4300
A	 0.1580	 0.4660
B	 0.1260	 0.3720
C	 0.1510	 0.4630
D	 0.1510	 0.4650
E	 0.1260	 0.3710
F	 0.1500	 0.4640
G	 0.1500	 0.4650
H	 0.1260	 0.3720
I	 0.1540	 0.4610
J	 0.1570	 0.4670
K	 0.1270	 0.3710
L	 0.1520	 0.4620
M	 0.1510	 0.4630
N	 0.1260	 0.3720
O	 0.1530	 0.4620
P	 0.1540	 0.4660
Q	 0.1240	 0.3750
R	 0.1510	 0.4640
S	 0.1490	 0.4670
T	 0.1230	 0.3760
U	 0.1460	 0.4630
V	 0.1490	 0.4660
W	 0.1240	 0.3780
X	 0.1500	 0.4650
Y	 0.1490	 0.4660
Z	 0.1240	 0.3730
a	 0.1520	 0.4630
b	 0.1530	 0.4670
c	 0.1260	 0.3750
d	 0.1530	 0.4640
e	 0.1520	 0.4650
f	 0.1240	 0.3720
g	 0.1540	 0.4640

