



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 14, 2023 – 02:06 PM JST

PDB ID : 5ZEN
Title : Crystal structure of human topoisomerase II beta in complex with DNA: a new quaternary conformation showing opening of the protein-linked DNA-gate
Authors : Chen, S.F.; Wang, Y.R.; Chan, N.L.
Deposited on : 2018-02-27
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1235 (2.78-2.74)
Clashscore	141614	1277 (2.78-2.74)
Ramachandran outliers	138981	1257 (2.78-2.74)
Sidechain outliers	138945	1257 (2.78-2.74)
RSRZ outliers	127900	1207 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	803	
2	B	8	
3	C	9	

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 5765 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA topoisomerase 2-beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	708	5395	3472	906	994	23	0	8	0

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	419	MET	-	expression tag	UNP Q02880
A	420	ALA	-	expression tag	UNP Q02880
A	421	SER	-	expression tag	UNP Q02880
A	422	TRP	-	expression tag	UNP Q02880
A	423	SER	-	expression tag	UNP Q02880
A	424	HIS	-	expression tag	UNP Q02880
A	425	PRO	-	expression tag	UNP Q02880
A	426	GLN	-	expression tag	UNP Q02880
A	427	PHE	-	expression tag	UNP Q02880
A	428	GLU	-	expression tag	UNP Q02880
A	429	LYS	-	expression tag	UNP Q02880
A	430	GLY	-	expression tag	UNP Q02880
A	431	ALA	-	expression tag	UNP Q02880
A	432	ASP	-	expression tag	UNP Q02880
A	433	ASP	-	expression tag	UNP Q02880
A	434	ASP	-	expression tag	UNP Q02880
A	435	ASP	-	expression tag	UNP Q02880
A	436	LYS	-	expression tag	UNP Q02880
A	437	VAL	-	expression tag	UNP Q02880
A	438	PRO	-	expression tag	UNP Q02880
A	439	ASP	-	expression tag	UNP Q02880
A	440	PRO	-	expression tag	UNP Q02880
A	441	THR	-	expression tag	UNP Q02880
A	442	SER	-	expression tag	UNP Q02880
A	443	VAL	-	expression tag	UNP Q02880
A	444	ASP	-	expression tag	UNP Q02880
A	1202	GLY	-	expression tag	UNP Q02880

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	1203	ALA	-	expression tag	UNP Q02880
A	1204	PRO	-	expression tag	UNP Q02880
A	1205	GLY	-	expression tag	UNP Q02880
A	1206	PHE	-	expression tag	UNP Q02880
A	1207	SER	-	expression tag	UNP Q02880
A	1208	SER	-	expression tag	UNP Q02880
A	1209	ILE	-	expression tag	UNP Q02880
A	1210	SER	-	expression tag	UNP Q02880
A	1211	ALA	-	expression tag	UNP Q02880
A	1212	HIS	-	expression tag	UNP Q02880
A	1213	HIS	-	expression tag	UNP Q02880
A	1214	HIS	-	expression tag	UNP Q02880
A	1215	HIS	-	expression tag	UNP Q02880
A	1216	HIS	-	expression tag	UNP Q02880
A	1217	HIS	-	expression tag	UNP Q02880
A	1218	HIS	-	expression tag	UNP Q02880
A	1219	HIS	-	expression tag	UNP Q02880
A	1220	HIS	-	expression tag	UNP Q02880
A	1221	HIS	-	expression tag	UNP Q02880

- Molecule 2 is a DNA chain called DNA (5'-D(P*AP*GP*CP*CP*GP*AP*GP*C)-3').

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
			Total	C	N	O				P
2	B	8	165	77	34	46	8	0	0	0

- Molecule 3 is a DNA chain called DNA (5'-D(P*AP*GP*CP*TP*CP*GP*GP*CP*T)-3').

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
			Total	C	N	O				P
3	C	9	184	87	33	55	9	0	0	0

- Molecule 4 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mn		
4	A	2	2	2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	19	Total O 19 19	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA topoisomerase 2-beta



- Molecule 2: DNA (5'-D(P*AP*GP*CP*CP*GP*AP*GP*C)-3')



- Molecule 3: DNA (5'-D(P*AP*GP*CP*TP*CP*GP*GP*CP*T)-3')



A12	
G13	
C14	
T15	
C16	T20
G17	

4 Data and refinement statistics i

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	95.76Å 95.76Å 231.06Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.97 – 2.75 19.97 – 2.75	Depositor EDS
% Data completeness (in resolution range)	99.7 (19.97-2.75) 99.7 (19.97-2.75)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.29 (at 2.75Å)	Xtrriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.221 , 0.266 0.217 , 0.264	Depositor DCC
R_{free} test set	2006 reflections (6.17%)	wwPDB-VP
Wilson B-factor (Å ²)	68.1	Xtrriage
Anisotropy	0.345	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 46.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.023 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5765	wwPDB-VP
Average B, all atoms (Å ²)	76.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.32	0/5530	0.54	0/7509
2	B	0.73	0/185	0.77	0/283
3	C	0.70	0/205	0.99	0/314
All	All	0.36	0/5920	0.58	0/8106

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5395	0	5103	89	0
2	B	165	0	89	1	0
3	C	184	0	102	3	0
4	A	2	0	0	0	0
5	A	19	0	0	0	0
All	All	5765	0	5294	91	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (91) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:809:ARG:NH2	1:A:941:GLU:OE1	2.09	0.85
1:A:978:THR:HG22	1:A:980:THR:H	1.41	0.85
1:A:752:ARG:HH12	1:A:758[B]:VAL:HG21	1.45	0.80
1:A:608:ILE:HG23	1:A:609:PRO:HD3	1.74	0.69
1:A:831:ARG:HH21	1:A:836:ALA:HA	1.58	0.69
1:A:1154:VAL:O	1:A:1158:ILE:HG13	1.94	0.68
1:A:678:LYS:NZ	1:A:875:GLY:O	2.26	0.67
1:A:751:LYS:HB3	1:A:770:MET:HE1	1.75	0.66
1:A:750:PHE:CG	1:A:831:ARG:NH1	2.63	0.66
1:A:1071:ILE:HG13	1:A:1149:LEU:HD23	1.79	0.65
1:A:934:ARG:HH11	1:A:934:ARG:HG3	1.63	0.64
1:A:1141:ILE:O	1:A:1144:MET:HG3	1.98	0.63
1:A:489:LEU:HB3	1:A:493:GLY:HA3	1.79	0.63
1:A:558:GLN:HG3	1:A:591:THR:O	1.97	0.63
1:A:1050:LYS:NZ	1:A:1174:LYS:O	2.27	0.63
1:A:670:SER:HB3	1:A:673:LYS:HG2	1.80	0.62
1:A:944:VAL:HG12	1:A:945:ARG:HG2	1.82	0.62
1:A:1068:ALA:O	1:A:1072:LEU:HB2	2.00	0.62
1:A:852:VAL:HG23	1:A:853:GLU:H	1.65	0.61
1:A:630:LYS:O	1:A:634:THR:OG1	2.19	0.61
1:A:657:ALA:N	1:A:661:ASP:OD2	2.37	0.57
1:A:732:PRO:HG2	1:A:869:ALA:HB1	1.86	0.57
3:C:12:DA:H2"	3:C:13:DG:C8	2.39	0.56
1:A:1067:GLN:O	1:A:1071:ILE:HG22	2.05	0.56
1:A:884:ASP:OD2	1:A:887[A]:GLU:HG3	2.05	0.56
1:A:459:ASP:OD2	1:A:548:ARG:NH2	2.39	0.56
1:A:611:PHE:HZ	1:A:626:ILE:HD13	1.71	0.55
1:A:934:ARG:HG3	1:A:934:ARG:NH1	2.22	0.54
1:A:959:MET:HB3	1:A:970:ILE:HG12	1.89	0.54
1:A:831:ARG:NH2	1:A:836:ALA:HA	2.22	0.54
1:A:1071:ILE:O	1:A:1075:ILE:HG22	2.08	0.54
1:A:474:ILE:HA	1:A:553:MET:O	2.09	0.53
1:A:736:ASP:OD2	1:A:743:ARG:HD3	2.09	0.52
1:A:457:LEU:HD22	1:A:529:ILE:HG12	1.91	0.52
1:A:1002:HIS:HB3	1:A:1008:GLN:CG	2.40	0.52
1:A:906:PRO:HG2	1:A:915:ILE:HG21	1.91	0.52
1:A:939:ILE:HB	1:A:984:PHE:HB2	1.91	0.51
1:A:554:ILE:HD12	1:A:566:LYS:HA	1.92	0.51
1:A:750:PHE:HB3	1:A:831:ARG:HH12	1.76	0.51
1:A:560:GLN:HB3	1:A:722:PHE:HA	1.92	0.51
1:A:574:HIS:HB2	1:A:581:LEU:HD22	1.93	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:807:GLY:HA3	1:A:813:GLY:HA2	1.93	0.50
1:A:993:LEU:O	1:A:997:GLU:HG2	2.12	0.49
1:A:1171:LEU:HD13	1:A:1174:LYS:HD2	1.93	0.49
1:A:858:ILE:HG13	1:A:1042:ARG:HD2	1.95	0.49
1:A:678:LYS:HD3	1:A:876:TRP:NE1	2.28	0.48
1:A:1066:ASN:ND2	1:A:1099:GLU:H	2.11	0.48
1:A:557:ASP:CG	1:A:631:GLY:HA2	2.34	0.48
1:A:834:PHE:CE2	1:A:859:PRO:HG3	2.48	0.47
1:A:934:ARG:O	1:A:993:LEU:HD23	2.14	0.47
1:A:752:ARG:NH1	1:A:754:ASP:OD2	2.47	0.47
1:A:1145:SER:OG	1:A:1147:TRP:HB2	2.14	0.47
1:A:978:THR:HG22	1:A:980:THR:N	2.21	0.47
1:A:479:ASP:HA	1:A:482:LYS:HB3	1.98	0.46
1:A:1164:LYS:O	1:A:1168:VAL:HG23	2.16	0.45
1:A:767:VAL:O	1:A:771:SER:OG	2.33	0.45
1:A:964:ASP:OD1	1:A:965:LYS:N	2.49	0.45
3:C:14:DC:H2''	3:C:15:DT:H5'	1.99	0.44
1:A:989:THR:OG1	1:A:992:LYS:HB2	2.18	0.44
1:A:875:GLY:N	3:C:17:DG:H5''	2.32	0.44
1:A:970:ILE:HG22	1:A:988:MET:HG2	1.98	0.44
1:A:1066:ASN:HD22	1:A:1099:GLU:H	1.63	0.44
1:A:750:PHE:HB3	1:A:831:ARG:NH1	2.33	0.44
1:A:555:MET:HE1	1:A:644:PHE:HZ	1.82	0.43
1:A:1063:LYS:O	1:A:1067:GLN:HG3	2.18	0.43
1:A:553:MET:CE	1:A:651:ARG:HD3	2.49	0.43
1:A:978:THR:HB	1:A:981:THR:OG1	2.18	0.43
1:A:752:ARG:NH1	1:A:754:ASP:OD1	2.51	0.43
1:A:482:LYS:O	1:A:486:VAL:HG23	2.19	0.43
1:A:955:VAL:C	1:A:958:PRO:HD2	2.39	0.43
1:A:844:PHE:HA	1:A:854:PRO:HA	2.00	0.43
1:A:677:ARG:O	1:A:680:TRP:HB3	2.18	0.43
1:A:608:ILE:CG2	1:A:609:PRO:HD3	2.47	0.42
1:A:748:THR:O	1:A:752:ARG:HG3	2.18	0.42
1:A:675:ASP:HA	1:A:678:LYS:HG3	2.02	0.42
1:A:586:LEU:HD23	1:A:586:LEU:HA	1.94	0.42
1:A:505:LYS:HG3	2:B:7:DG:H21	1.85	0.42
1:A:553:MET:SD	1:A:589:PHE:HB2	2.60	0.42
1:A:510:ARG:NH2	1:A:571:ASN:OD1	2.45	0.41
1:A:744:LYS:HG2	1:A:771:SER:HB2	2.02	0.41
1:A:842:LEU:HD21	1:A:858:ILE:HG22	2.02	0.41
1:A:1043:LEU:HB2	1:A:1180:TRP:HZ3	1.85	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:883:TYR:OH	1:A:1031[A]:VAL:HG21	2.21	0.41
1:A:538:TYR:CE1	1:A:547:LEU:HD21	2.56	0.41
1:A:1075:ILE:HD13	1:A:1075:ILE:HG21	1.84	0.41
1:A:727:ASN:OD1	1:A:873:GLY:HA3	2.21	0.41
1:A:1138:PHE:HA	1:A:1140:TYR:CE1	2.56	0.41
1:A:738:PHE:HB2	1:A:743:ARG:NH1	2.36	0.40
1:A:759:LYS:HD3	1:A:820:ARG:O	2.21	0.40
1:A:1071:ILE:HG21	1:A:1157:LEU:CD2	2.51	0.40
1:A:1002:HIS:HB3	1:A:1008:GLN:HG2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	706/803 (88%)	691 (98%)	15 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	528/704 (75%)	527 (100%)	1 (0%)	93 96

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	979	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	850	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	708/803 (88%)	-0.07	19 (2%) 54 63	45, 70, 123, 173	0
2	B	8/8 (100%)	0.18	1 (12%) 3 4	54, 59, 93, 112	0
3	C	9/9 (100%)	0.49	1 (11%) 5 6	56, 60, 92, 128	0
All	All	725/820 (88%)	-0.06	21 (2%) 51 61	45, 70, 123, 173	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	964	ASP	5.3
1	A	614	TRP	3.9
2	B	1	DA	3.8
1	A	694	HIS	3.6
1	A	695	GLY	3.6
3	C	12	DA	3.6
1	A	626	ILE	3.4
1	A	645	ALA	3.3
1	A	1109	ALA	3.2
1	A	630	LYS	3.2
1	A	591	THR	2.9
1	A	1158	ILE	2.8
1	A	628	TYR	2.8
1	A	1106	TRP	2.5
1	A	1163	ALA	2.4
1	A	846	TYR	2.4
1	A	604	SER	2.3
1	A	613	GLU	2.2
1	A	629	TYR	2.1
1	A	859	PRO	2.1
1	A	605	PHE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	MN	A	1302	1/1	0.69	0.16	114,114,114,114	0
4	MN	A	1301	1/1	0.88	0.07	102,102,102,102	0

6.5 Other polymers [i](#)

There are no such residues in this entry.