



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 29, 2021 – 03:04 pm BST

PDB ID : 6ZG4
Title : Structure of M1-StaR-T4L in complex with HTL0009936 at 2.35Å
Authors : Rucktooa, P.; Cooke, R.M.
Deposited on : 2020-06-18
Resolution : 2.33 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2096 (2.36-2.32)
Clashscore	141614	2193 (2.36-2.32)
Ramachandran outliers	138981	2159 (2.36-2.32)
Sidechain outliers	138945	2160 (2.36-2.32)
RSRZ outliers	127900	2067 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	455	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	OLA	A	1206	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	OLA	A	1212	-	-	-	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3899 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Muscarinic acetylcholine receptor M1,Endolysin,Muscarinic acetylcholine receptor M1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	446	3556	2311	600	621	24	0	0	0

There are 49 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	20	MET	-	initiating methionine	UNP P11229
A	21	GLU	-	expression tag	UNP P11229
A	22	THR	-	expression tag	UNP P11229
A	23	VAL	-	expression tag	UNP P11229
A	24	GLU	-	expression tag	UNP P11229
A	25	MET	-	expression tag	UNP P11229
A	26	VAL	-	expression tag	UNP P11229
A	27	ALA	PHE	engineered mutation	UNP P11229
A	29	ALA	GLY	conflict	UNP P11229
A	30	THR	ILE	conflict	UNP P11229
A	31	VAL	THR	conflict	UNP P11229
A	32	ALA	THR	engineered mutation	UNP P11229
A	44	ILE	LEU	engineered mutation	UNP P11229
A	46	LEU	VAL	engineered mutation	UNP P11229
A	47	MET	LEU	conflict	UNP P11229
A	48	LEU	ILE	conflict	UNP P11229
A	50	ILE	PHE	conflict	UNP P11229
A	54	ARG	THR	conflict	UNP P11229
A	55	GLN	GLU	conflict	UNP P11229
A	57	GLN	LYS	conflict	UNP P11229
A	64	ALA	LEU	engineered mutation	UNP P11229
A	65	PHE	LEU	conflict	UNP P11229
A	76	ALA	THR	engineered mutation	UNP P11229
A	84	VAL	THR	engineered mutation	UNP P11229
A	86	ILE	LEU	conflict	UNP P11229
A	87	ILE	LEU	conflict	UNP P11229

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	95	ALA	THR	engineered mutation	UNP P11229
A	101	ALA	TRP	engineered mutation	UNP P11229
A	112	ALA	SER	engineered mutation	UNP P11229
A	143	LEU	ALA	engineered mutation	UNP P11229
A	196	THR	ALA	engineered mutation	UNP P11229
A	1012	GLY	ARG	conflict	UNP P00720
A	1054	THR	CYS	conflict	UNP P00720
A	1097	ALA	CYS	conflict	UNP P00720
A	1137	ARG	ILE	conflict	UNP P00720
A	362	ALA	LYS	engineered mutation	UNP P11229
A	364	LEU	ALA	engineered mutation	UNP P11229
A	411	ALA	SER	engineered mutation	UNP P11229
A	435	ALA	CYS	conflict	UNP P11229
A	439	HIS	-	expression tag	UNP P11229
A	440	HIS	-	expression tag	UNP P11229
A	441	HIS	-	expression tag	UNP P11229
A	442	HIS	-	expression tag	UNP P11229
A	443	HIS	-	expression tag	UNP P11229
A	444	HIS	-	expression tag	UNP P11229
A	445	HIS	-	expression tag	UNP P11229
A	446	HIS	-	expression tag	UNP P11229
A	447	HIS	-	expression tag	UNP P11229
A	448	HIS	-	expression tag	UNP P11229

- Molecule 2 is ethyl (4 {S})-4-[4-[(1-methylcyclobutyl)carbamoyl]piperidin-1-yl]azepane-1-carboxylate (three-letter code: QK8) (formula: C₂₀H₃₅N₃O₃) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	26	20	3	3	0	0

- Molecule 3 is OLEIC ACID (three-letter code: OLA) (formula: $C_{18}H_{34}O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			20	18	2		
3	A	1	Total	C	O	0	0
			20	18	2		
3	A	1	Total	C	O	0	0
			20	18	2		
3	A	1	Total	C	O	0	0
			20	18	2		
3	A	1	Total	C	O	0	0
			20	18	2		
3	A	1	Total	C	O	0	0
			20	18	2		
3	A	1	Total	C	O	0	0
			20	18	2		
3	A	1	Total	C	O	0	0
			20	18	2		

- Molecule 4 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 5 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	P	0	0
			5	4	1		
5	A	1	Total	O	P	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	P	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	72	Total	O	0	0
			72	72		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Muscarinic acetylcholine receptor M1,Endolysin,Muscarinic acetylcholine receptor M1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	62.47Å 65.45Å 154.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.98 – 2.33 49.98 – 2.33	Depositor EDS
% Data completeness (in resolution range)	74.0 (49.98-2.33) 74.0 (49.98-2.33)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.60 (at 2.34Å)	Xtrriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.204 , 0.234 0.216 , 0.250	Depositor DCC
R_{free} test set	1000 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	25.2	Xtrriage
Anisotropy	0.076	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.017 for k,h,-l	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3899	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, PO4, OLA, QK8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.46	0/3634	0.62	0/4942

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3556	0	3636	26	0
2	A	26	0	0	0	0
3	A	220	0	363	14	0
4	A	10	0	14	0	0
5	A	15	0	0	0	0
6	A	72	0	0	0	0
All	All	3899	0	4013	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 26 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:96:LEU:HB2	3:A:1213:OLA:H62	1.74	0.68
1:A:95:ALA:CB	3:A:1213:OLA:H22	2.35	0.57
1:A:1087:VAL:HG11	1:A:1118:LEU:HD22	1.87	0.55
1:A:114:MET:HE1	3:A:1205:OLA:H10	1.87	0.55
1:A:96:LEU:HB2	3:A:1213:OLA:C6	2.38	0.54

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	444/455 (98%)	436 (98%)	8 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	376/385 (98%)	368 (98%)	8 (2%)	53 65

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	439	HIS
1	A	431	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	394	CYS
1	A	1132	ASN
1	A	399	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1069	GLN
1	A	1132	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	OLA	A	1203	-	16,19,19	0.23	0	15,19,19	0.14	0
2	QK8	A	1201	-	26,28,28	2.07	6 (23%)	25,39,39	1.78	5 (20%)
3	OLA	A	1215	-	16,19,19	0.21	0	15,19,19	0.15	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	OLA	A	1212	-	16,19,19	0.21	0	15,19,19	0.15	0
3	OLA	A	1206	-	16,19,19	0.21	0	15,19,19	0.17	0
4	PGE	A	1208	-	9,9,9	0.21	0	8,8,8	0.10	0
3	OLA	A	1214	-	16,19,19	0.20	0	15,19,19	0.18	0
3	OLA	A	1207	-	16,19,19	0.23	0	15,19,19	0.26	0
3	OLA	A	1213	-	16,19,19	0.20	0	15,19,19	0.17	0
5	PO4	A	1216	-	4,4,4	2.49	1 (25%)	6,6,6	0.51	0
5	PO4	A	1209	-	4,4,4	2.47	1 (25%)	6,6,6	0.47	0
3	OLA	A	1205	-	16,19,19	0.22	0	15,19,19	0.22	0
3	OLA	A	1202	-	16,19,19	0.23	0	15,19,19	0.21	0
3	OLA	A	1204	-	16,19,19	0.25	0	15,19,19	0.16	0
3	OLA	A	1211	-	16,19,19	0.22	0	15,19,19	0.15	0
5	PO4	A	1210	-	4,4,4	2.49	1 (25%)	6,6,6	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OLA	A	1203	-	-	7/15/17/17	-
2	QK8	A	1201	-	-	4/19/49/49	0/3/3/3
3	OLA	A	1215	-	-	9/15/17/17	-
3	OLA	A	1212	-	-	8/15/17/17	-
3	OLA	A	1206	-	-	10/15/17/17	-
4	PGE	A	1208	-	-	6/7/7/7	-
3	OLA	A	1214	-	-	9/15/17/17	-
3	OLA	A	1207	-	-	9/15/17/17	-
3	OLA	A	1213	-	-	8/15/17/17	-
3	OLA	A	1205	-	-	9/15/17/17	-
3	OLA	A	1202	-	-	10/15/17/17	-
3	OLA	A	1204	-	-	8/15/17/17	-
3	OLA	A	1211	-	-	12/15/17/17	-

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1201	QK8	C5-N3	5.23	1.57	1.47
2	A	1201	QK8	C1-N3	4.98	1.56	1.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1201	QK8	C16-N2	4.26	1.42	1.35
5	A	1210	PO4	P-O1	4.14	1.60	1.50
5	A	1216	PO4	P-O1	4.12	1.60	1.50

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1201	QK8	C17-O3-C16	5.61	124.00	115.59
2	A	1201	QK8	O1-C6-C3	-2.94	118.29	122.12
2	A	1201	QK8	C5-N3-C1	2.60	113.83	109.08
2	A	1201	QK8	C14-C13-N2	2.48	120.48	113.93
2	A	1201	QK8	C4-C3-C2	2.19	114.60	109.97

There are no chirality outliers.

5 of 109 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1201	QK8	C11-C10-N3-C1
3	A	1202	OLA	C1-C2-C3-C4
3	A	1203	OLA	C1-C2-C3-C4
3	A	1203	OLA	C11-C10-C9-C8
3	A	1205	OLA	C1-C2-C3-C4


There are no ring outliers.




5 monomers are involved in 14 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1215	OLA	2	0
3	A	1207	OLA	1	0
3	A	1213	OLA	8	0
3	A	1205	OLA	2	0
3	A	1202	OLA	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	446/455 (98%)	0.60	45 (10%) 7 11	12, 30, 60, 82	0

The worst 5 of 45 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	437	TRP	7.2
1	A	438	ASP	6.2
1	A	354	THR	6.2
1	A	1054	THR	5.6
1	A	1057	VAL	4.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	OLA	A	1212	20/20	0.57	0.51	67,75,82,82	0
3	OLA	A	1211	20/20	0.62	0.34	73,74,77,77	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	OLA	A	1207	20/20	0.63	0.38	68,70,73,73	0
5	PO4	A	1216	5/5	0.64	0.28	117,117,118,118	0
3	OLA	A	1206	20/20	0.65	0.44	54,58,63,63	0
3	OLA	A	1215	20/20	0.72	0.29	42,51,59,59	0
4	PGE	A	1208	10/10	0.74	0.20	46,50,53,54	0
3	OLA	A	1213	20/20	0.75	0.30	51,57,59,59	0
3	OLA	A	1203	20/20	0.77	0.27	33,39,59,60	0
3	OLA	A	1205	20/20	0.79	0.28	44,50,60,60	0
3	OLA	A	1214	20/20	0.80	0.20	47,50,53,54	0
3	OLA	A	1204	20/20	0.82	0.19	33,40,57,60	0
3	OLA	A	1202	20/20	0.87	0.27	27,31,42,42	0
5	PO4	A	1210	5/5	0.92	0.26	79,80,82,82	0
5	PO4	A	1209	5/5	0.93	0.18	68,68,69,70	0
2	QK8	A	1201	26/26	0.96	0.15	13,15,29,30	0



The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.








Electron density around OLA A 1207:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around OLA A 1206:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLA A 1215:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around OLA A 1213:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLA A 1203:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around OLA A 1205:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)








Electron density around OLA A 1204:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OLA A 1202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around QK8 A 1201:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.