

# Full wwPDB X-ray Structure Validation Report (i)

#### Jun 12, 2024 – 08:48 PM EDT

PDB ID	:	3ZHQ
Title	:	Crystal structure of the H747A mutant of the SucA domain of Mycobacterium
		smegmatis KGD
Authors	:	Wagner, T.; Barilone, N.; Bellinzoni, M.; Alzari, P.M.
Deposited on	:	2012-12-24
Resolution	:	2.50  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	•	4.02b-467
Morry	÷	20002.2  0 CCD $ac 542  b a (2002)$
Mogui	•	2022.3.0, CSD as $3430e(2022)$
Xtriage (Phenix)	:	1.20.1
$\mathrm{EDS}$	:	2.36.2
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
	$(\# { m Entries})$	$(\# { m Entries},  { m resolution}  { m range}({ m \AA}))$
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	$5231 \ (2.50-2.50)$
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	868	2% <b>88%</b>	5%	7%
1	В	868	4% 87%	6%	7%
1	С	868	4% 88%	5%	6%
1	D	868	4%	•	7%



#### 3ZHQ

# 2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 25781 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called MULTIFUNCTIONAL 2-OXOGLUTARATE METABOLISM ENZYME.

Mol	Chain	Residues		Α	toms			ZeroOcc	AltConf	Trace
1	Λ	<b>Q</b> 11	Total	С	Ν	Ο	S	0	1	0
1	Л	011	6283	3959	1108	1194	22	0		
1	В	808	Total	С	Ν	Ο	S	0	0	0
1	I D	000	6208	3913	1103	1168	24			
1	C	812	Total	С	Ν	Ο	S	0	0	0
1			6301	3971	1110	1197	23			
1	1 D	810	Total	С	Ν	Ο	S	0	0	0
1			6234	3933	1097	1180	24			0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	360	GLY	-	expression tag	UNP A0R2B1
А	747	ALA	HIS	engineered mutation	UNP A0R2B1
В	360	GLY	-	expression tag	UNP A0R2B1
В	747	ALA	HIS	engineered mutation	UNP A0R2B1
С	360	GLY	-	expression tag	UNP A0R2B1
С	747	ALA	HIS	engineered mutation	UNP A0R2B1
D	360	GLY	-	expression tag	UNP A0R2B1
D	747	ALA	HIS	engineered mutation	UNP A0R2B1

• Molecule 2 is THIAMINE DIPHOSPHATE (three-letter code: TPP) (formula:  $C_{12}H_{19}N_4O_7P_2S$ ).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf		
0	Λ	1	Total	С	Ν	0	Р	S	0	0	
	A		26	12	4	7	2	1	0	0	
9	В	1	Total	С	Ν	0	Р	S	0	0	
	D	1	26	12	4	7	2	1	0		
0	С	1	Total	С	Ν	Ο	Р	$\mathbf{S}$	0	0	
	U		26	12	4	7	2	1	0	0	
	1	Total	С	Ν	0	Р	S	0	0		
	D	1	26	12	4	7	2	1	0	0	

• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total Mg 1 1	0	0
3	В	1	Total Mg 1 1	0	0
3	С	1	Total Mg 1 1	0	0
3	D	1	Total Mg 1 1	0	0

• Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	Total Ca 1 1	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	В	1	Total Ca 1 1	0	0
4	С	1	Total Ca 1 1	0	0
4	D	1	Total Ca 1 1	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	197	Total O 197 197	0	0
5	В	154	Total O 154 154	0	0
5	С	165	Total O 165 165	0	0
5	D	127	Total O 127 127	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: MULTIFUNCTIONAL 2-OXOGLUTARATE METABOLISM ENZYME



• Molecule 1: MULTIFUNCTIONAL 2-OXOGLUTARATE METABOLISM ENZYME







# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants	80.67Å 83.74Å 159.58Å	Deperitor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$99.86^{\circ}$ $98.95^{\circ}$ $100.44^{\circ}$	Depositor
$\mathbf{P}_{\text{assolution}}(\hat{\mathbf{A}})$	48.97 - 2.50	Depositor
Resolution (A)	48.97 - 2.50	EDS
% Data completeness	96.8 (48.97-2.50)	Depositor
(in resolution range)	96.7(48.97 - 2.50)	EDS
R <sub>merge</sub>	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.28 (at 2.51 \text{\AA})$	Xtriage
Refinement program	BUSTER 2.11.2	Depositor
P. P.	0.194 , $0.223$	Depositor
$\Pi, \Pi_{free}$	0.203 , $0.238$	DCC
$R_{free}$ test set	6718 reflections $(5.05%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	36.9	Xtriage
Anisotropy	0.401	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.32, 54.4	EDS
L-test for $twinning^2$	$< L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	0.012 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	25781	wwPDB-VP
Average B, all atoms $(Å^2)$	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 15.98% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, TPP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond lengths		Bond angles	
	Unam	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.51	0/6411	0.65	0/8699
1	В	0.51	0/6335	0.65	0/8600
1	С	0.50	0/6427	0.65	0/8712
1	D	0.50	0/6359	0.65	0/8625
All	All	0.50	0/25532	0.65	0/34636

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	6283	0	6051	23	0
1	В	6208	0	5963	27	0
1	С	6301	0	6108	27	0
1	D	6234	0	6017	23	0
2	А	26	0	16	2	0
2	В	26	0	16	2	0
2	С	26	0	16	2	0
2	D	26	0	16	2	0
3	A	1	0	0	0	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	В	1	0	0	0	0
3	С	1	0	0	0	0
3	D	1	0	0	0	0
4	А	1	0	0	0	0
4	В	1	0	0	0	0
4	С	1	0	0	0	0
4	D	1	0	0	0	0
5	А	197	0	0	2	0
5	В	154	0	0	1	0
5	С	165	0	0	0	0
5	D	127	0	0	4	0
All	All	25781	0	24203	99	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (99) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
1:C:705:PRO:HG2	1:C:735:VAL:HG13	1.69	0.73
1:C:373:LEU:CD1	1:C:436:LEU:HD21	2.20	0.71
1:C:692:SER:HB2	1:C:697:ASP:OD2	1.93	0.67
1:C:373:LEU:HD11	1:C:436:LEU:CD2	2.27	0.65
1:A:692:SER:HB2	1:A:697:ASP:OD2	1.98	0.63
1:C:373:LEU:HD11	1:C:436:LEU:HD21	1.82	0.62
1:C:373:LEU:CD1	1:C:436:LEU:CD2	2.78	0.61
1:C:373:LEU:HD12	1:C:436:LEU:HD21	1.84	0.57
1:A:530:LEU:HD22	1:A:636:SER:HA	1.88	0.56
1:A:901:GLN:OE1	2:B:2001:TPP:H6'	2.07	0.55
1:D:1174:GLN:OE1	1:D:1206:PRO:HA	2.07	0.55
1:B:542:ARG:HD3	1:B:578:TYR:HA	1.90	0.53
1:C:1112:LEU:HD12	1:C:1136:ALA:HB3	1.91	0.53
1:D:1069:PRO:HB3	1:D:1072:MET:HB3	1.91	0.53
1:A:442:ALA:HB1	1:A:467:VAL:HG12	1.91	0.52
1:B:603:PRO:HD3	1:B:991:PHE:CZ	2.45	0.52
1:C:898:LEU:O	1:C:945:VAL:HA	2.09	0.52
1:A:603:PRO:HD3	1:A:991:PHE:CZ	2.45	0.51
1:D:1112:LEU:HD12	1:D:1136:ALA:HB3	1.93	0.51
1:B:1112:LEU:HD12	1:B:1136:ALA:HB3	1.92	0.51
1:C:901:GLN:OE1	2:D:2001:TPP:H6'	2.11	0.51
1:A:1112:LEU:HD12	1:A:1136:ALA:HB3	1.92	0.50



		Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
1:B:618:ARG:HH21	1:B:629:GLU:HG2	1.77	0.49
1:C:441:ASP:HA	1:C:445:ARG:HG3	1.94	0.49
1:D:639:PRO:HB2	1:D:671:THR:HG23	1.94	0.49
1:B:530:LEU:HD22	1:B:636:SER:HA	1.93	0.49
1:C:1148:ARG:HG3	1:C:1187:ILE:HD12	1.94	0.49
2:C:2001:TPP:H6'	1:D:901:GLN:OE1	2.12	0.49
1:C:1112:LEU:HD21	1:C:1155:LEU:HD22	1.94	0.49
1:D:538:PRO:HB2	5:D:3019:HOH:O	2.13	0.49
1:D:493:GLU:O	1:D:497:GLN:HG2	2.13	0.48
1:B:536:ALA:HB3	1:B:613:LEU:HD22	1.94	0.48
1:D:981:VAL:HG22	1:D:988:ILE:HD11	1.95	0.48
1:A:1086:THR:O	1:C:1157:ARG:NH2	2.41	0.48
1:B:452:THR:HA	1:B:460:GLN:NE2	2.29	0.48
1:D:1147:PRO:O	1:D:1151:LEU:HB2	2.13	0.48
1:A:981:VAL:HG22	1:A:988:ILE:HD11	1.96	0.47
1:C:981:VAL:HG22	1:C:988:ILE:HD11	1.96	0.47
1:A:878:TRP:HB3	1:A:1073:LEU:HD23	1.95	0.47
1:B:510:GLY:O	1:B:741:CYS:HB2	2.15	0.47
1:C:603:PRO:HD3	1:C:991:PHE:CZ	2.51	0.46
1:A:1112:LEU:HD21	1:A:1155:LEU:HD22	1.98	0.46
1:D:986:SER:O	1:D:990:GLU:HB2	2.16	0.46
1:C:1069:PRO:HB2	1:C:1072:MET:HB3	1.97	0.46
1:C:1177:TRP:CD1	1:C:1197:ARG:HD3	2.51	0.46
1:B:908:PHE:CZ	1:B:1070:LYS:HG2	2.51	0.46
1:B:981:VAL:HG22	1:B:988:ILE:HD11	1.96	0.46
1:A:538:PRO:HD2	5:A:3028:HOH:O	2.15	0.46
1:B:1177:TRP:CD1	1:B:1197:ARG:HD3	2.51	0.45
1:C:1069:PRO:CB	1:C:1072:MET:HB3	2.47	0.45
1:C:1174:GLN:OE1	1:C:1206:PRO:HA	2.17	0.45
2:C:2001:TPP:HM41	1:D:950:LEU:HD22	1.98	0.45
1:B:898:LEU:O	1:B:945:VAL:HA	2.17	0.45
1:C:510:GLY:O	1:C:741:CYS:HB2	2.16	0.45
1:A:692:SER:HB3	1:B:701:MET:O	2.16	0.45
1:A:1083:ARG:HA	1:A:1086:THR:OG1	2.16	0.44
1:D:530:LEU:HD22	1:D:636:SER:HA	1.98	0.44
1:D:908:PHE:CZ	1:D:1070:LYS:HG2	2.53	0.44
1:D:1177:TRP:CD1	1:D:1197:ARG:HD3	2.52	0.44
1:A:870:MET:HB3	1:A:876:ILE:HG12	2.00	0.44
1:B:538:PRO:HB2	5:B:3018:HOH:O	2.18	0.44
1:A:1177:TRP:CD1	1:A:1197:ARG:HD3	2.53	0.44
1:D:596:GLU:HG3	5:D:3029:HOH:O	2.17	0.44



A 4 amo 1	A4	Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
1:B:692:SER:HB2	1:B:697:ASP:OD2	2.18	0.44
1:D:878:TRP:HB3	1:D:1073:LEU:HD23	2.00	0.44
1:C:554:TYR:HD2	1:C:806:GLU:OE1	2.01	0.43
1:D:1069:PRO:CB	1:D:1072:MET:HB3	2.48	0.43
2:A:2001:TPP:H61	2:A:2001:TPP:HM41	1.93	0.43
1:B:878:TRP:HB3	1:B:1073:LEU:HD23	2.01	0.43
1:B:1112:LEU:HD21	1:B:1155:LEU:HD22	2.00	0.43
1:C:682:PHE:HD1	1:C:683:THR:HG23	1.83	0.43
1:B:447:VAL:HG22	1:B:709:VAL:HG23	2.00	0.43
1:A:1016:GLN:NE2	5:A:3143:HOH:O	2.22	0.43
1:D:656:GLU:HB3	1:D:954:ALA:HB2	2.01	0.43
1:B:1174:GLN:OE1	1:B:1206:PRO:HA	2.18	0.42
1:A:656:GLU:HB3	1:A:954:ALA:HB2	2.01	0.42
1:B:656:GLU:HB3	1:B:954:ALA:HB2	2.02	0.42
1:A:696:THR:HG21	1:A:738:ASP:HB2	2.01	0.42
2:A:2001:TPP:H6'	1:B:901:GLN:OE1	2.18	0.42
1:B:543:LEU:HA	1:B:546:LEU:HD12	2.01	0.42
2:B:2001:TPP:H61	2:B:2001:TPP:HM41	1.93	0.42
1:A:778:LEU:HB3	1:A:784:ILE:HG12	2.01	0.41
1:D:1021:THR:HG21	1:D:1207:SER:HB3	2.02	0.41
1:B:536:ALA:CB	1:B:613:LEU:HD22	2.50	0.41
1:A:1155:LEU:HD11	1:A:1192:PHE:CZ	2.55	0.41
1:B:415:LEU:HA	1:B:432:LEU:HB3	2.02	0.41
1:C:656:GLU:HB3	1:C:954:ALA:HB2	2.02	0.41
1:D:596:GLU:CG	5:D:3029:HOH:O	2.68	0.41
1:D:769:GLY:HA3	5:D:3009:HOH:O	2.21	0.41
1:B:1021:THR:HG21	1:B:1207:SER:HB3	2.03	0.41
1:D:1112:LEU:HD21	1:D:1155:LEU:HD22	2.01	0.41
1:A:510:GLY:O	1:A:741:CYS:HB2	2.21	0.41
1:B:588:GLN:OE1	1:B:593:ASN:HB2	2.21	0.41
1:C:603:PRO:HG3	1:C:990:GLU:HB3	2.03	0.40
1:C:1021:THR:HG21	1:C:1207:SER:HB3	2.03	0.40
1:A:1115:THR:O	1:A:1139:ARG:HA	2.21	0.40
1:C:950:LEU:HD22	2:D:2001:TPP:HM41	2.02	0.40
1:A:1019:ASP:CB	1:B:604:SER:HB2	2.51	0.40
1:D:1109:VAL:HG21	1:D:1136:ALA:HB2	2.03	0.40

There are no symmetry-related clashes.



## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percent	iles
1	А	802/868~(92%)	779~(97%)	20 (2%)	3~(0%)	34 5	54
1	В	798/868~(92%)	779~(98%)	18 (2%)	1 (0%)	51 7	73
1	С	802/868~(92%)	779~(97%)	21 (3%)	2(0%)	47 6	38
1	D	800/868~(92%)	777~(97%)	22 (3%)	1 (0%)	51 7	73
All	All	3202/3472~(92%)	3114 (97%)	81 (2%)	7 (0%)	47 6	38

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	С	605	HIS
1	А	605	HIS
1	А	636	SER
1	А	682	PHE
1	В	682	PHE
1	С	682	PHE
1	D	682	PHE

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percer	ntiles
1	А	645/725~(89%)	637~(99%)	8 (1%)	71	88
1	В	629/725~(87%)	622~(99%)	7 (1%)	73	89
1	С	651/725~(90%)	644 (99%)	7 (1%)	73	89



COULU	Continuated from previous page							
Mol	Chain	Analysed	Rotameric	Outliers	P	<b>e</b> rce	ntile	es
1	D	637/725~(88%)	634 (100%)	3~(0%)		88	96	
All	All	2562/2900~(88%)	2537 (99%)	25 (1%)		76	90	

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	373	LEU
1	А	793	LEU
1	А	950	LEU
1	А	953	PHE
1	А	979	ASP
1	А	1100	THR
1	А	1114	LEU
1	А	1210	SER
1	В	432	LEU
1	В	616	LEU
1	В	793	LEU
1	В	950	LEU
1	В	953	PHE
1	В	979	ASP
1	В	1187	ILE
1	С	373	LEU
1	С	793	LEU
1	С	930	LEU
1	С	950	LEU
1	С	953	PHE
1	С	979	ASP
1	С	1083	ARG
1	D	467	VAL
1	D	950	LEU
1	D	953	PHE

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such side chains are listed below:

Mol	Chain	Res	Type
1	С	464	GLN
1	D	464	GLN



#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Tune	Tuno	Chain	Dog	Tink	Bond lengths			Bond angles		
	туре	Unain	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	TPP	C	2001	3	23,27,27	2.10	5 (21%)	30,40,40	1.82	7 (23%)
2	TPP	В	2001	3	23,27,27	2.07	3 (13%)	30,40,40	1.74	7 (23%)
2	TPP	А	2001	3	23,27,27	1.64	3 (13%)	30,40,40	1.80	8 (26%)
2	TPP	D	2001	3	23,27,27	2.00	3 (13%)	30,40,40	1.70	7 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TPP	С	2001	3	-	5/16/17/17	0/2/2/2
2	TPP	В	2001	3	-	3/16/17/17	0/2/2/2
2	TPP	А	2001	3	-	3/16/17/17	0/2/2/2
2	TPP	D	2001	3	-	1/16/17/17	0/2/2/2



3ZHQ
------

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	2001	TPP	C4-N3	-7.49	1.33	1.39
2	С	2001	TPP	C4-N3	-7.38	1.33	1.39
2	В	2001	TPP	C4-N3	-7.34	1.33	1.39
2	А	2001	TPP	C4-N3	-5.16	1.35	1.39
2	В	2001	TPP	PB-O1B	3.82	1.62	1.50
2	D	2001	TPP	PB-O1B	3.74	1.62	1.50
2	С	2001	TPP	PB-O1B	3.15	1.60	1.50
2	С	2001	TPP	C2-N3	-3.10	1.29	1.36
2	В	2001	TPP	C2-N3	-2.90	1.29	1.36
2	А	2001	TPP	PB-O1B	2.81	1.59	1.50
2	А	2001	TPP	C2-N3	-2.67	1.30	1.36
2	D	2001	TPP	C2-N3	-2.67	1.30	1.36
2	С	2001	TPP	PA-O3A	2.38	1.62	1.59
2	С	2001	TPP	C6-C5	2.17	1.52	1.51

All (14) bond length outliers are listed below:

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	А	2001	TPP	CM4-C4-C5	-4.46	117.84	127.60
2	В	2001	TPP	CM4-C4-C5	-4.31	118.19	127.60
2	А	2001	TPP	C5-C4-N3	4.24	116.06	107.57
2	С	2001	TPP	CM4-C4-C5	-4.12	118.59	127.60
2	D	2001	TPP	CM4-C4-C5	-3.88	119.11	127.60
2	В	2001	TPP	C5-C4-N3	3.82	115.22	107.57
2	С	2001	TPP	C5-C4-N3	3.77	115.11	107.57
2	D	2001	TPP	C5-C4-N3	3.52	114.62	107.57
2	С	2001	TPP	C6'-N1'-C2'	3.40	121.66	116.07
2	С	2001	TPP	C7'-N3-C2	-3.35	119.30	125.35
2	А	2001	TPP	C7'-N3-C2	-3.30	119.39	125.35
2	В	2001	TPP	CM4-C4-N3	3.15	126.55	122.53
2	D	2001	TPP	C7'-N3-C2	-3.15	119.66	125.35
2	D	2001	TPP	C6'-N1'-C2'	3.12	121.20	116.07
2	В	2001	TPP	C6'-N1'-C2'	3.03	121.05	116.07
2	А	2001	TPP	CM2-C2'-N1'	2.86	120.24	117.20
2	С	2001	TPP	CM4-C4-N3	2.85	126.17	122.53
2	D	2001	TPP	CM4-C4-N3	2.83	126.14	122.53
2	С	2001	TPP	N1'-C2'-N3'	-2.80	120.87	125.53
2	В	2001	TPP	C7'-N3-C2	-2.69	120.49	125.35
2	А	2001	TPP	CM4-C4-N3	2.67	125.93	122.53
2	D	2001	TPP	C5'-C6'-N1'	-2.64	119.53	123.83
2	А	2001	TPP	C6'-N1'-C2'	2.56	120.27	116.07
2	А	2001	TPP	N1'-C2'-N3'	-2.50	121.37	125.53



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	С	2001	TPP	CM2-C2'-N1'	2.33	119.68	117.20
2	В	2001	TPP	N1'-C2'-N3'	-2.32	121.66	125.53
2	В	2001	TPP	C5'-C6'-N1'	-2.27	120.14	123.83
2	А	2001	TPP	C2'-N3'-C4'	2.22	121.51	118.10
2	D	2001	TPP	N1'-C2'-N3'	-2.11	122.01	125.53

There are no chirality outliers.

Mol	Chain	Res	Type	Atoms
2	А	2001	TPP	C7-O7-PA-O1A
2	В	2001	TPP	C7-O7-PA-O1A
2	С	2001	TPP	C7-O7-PA-O1A
2	D	2001	TPP	C7-O7-PA-O1A
2	А	2001	TPP	C4-C5-C6-C7
2	В	2001	TPP	C4-C5-C6-C7
2	С	2001	TPP	C5-C6-C7-O7
2	С	2001	TPP	PA-O3A-PB-O2B
2	С	2001	TPP	PA-O3A-PB-O3B
2	С	2001	TPP	C7-O7-PA-O3A
2	А	2001	TPP	PA-O3A-PB-O3B
2	В	2001	TPP	C5-C6-C7-O7

All (12) torsion outliers are listed below:

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	С	2001	TPP	2	0
2	В	2001	TPP	2	0
2	А	2001	TPP	2	0
2	D	2001	TPP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.



The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers (i)

There are no such residues in this entry.



## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	< <b>RSRZ</b> >	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	811/868~(93%)	0.01	20 (2%) 57 61	22, 38, 68, 97	0
1	В	808/868~(93%)	0.11	38 (4%) 31 33	20, 40, 75, 103	0
1	С	812/868~(93%)	0.14	32 (3%) 39 42	22, 42, 74, 98	0
1	D	810/868~(93%)	0.18	38 (4%) 31 33	22, 42, 75, 114	0
All	All	3241/3472~(93%)	0.11	128 (3%) 39 42	20, 41, 73, 114	0

All (128) RSRZ outliers are listed below:

Mol	Chain	$\mathbf{Res}$	Type	RSRZ
1	А	368	ALA	5.9
1	В	472	ASP	5.8
1	С	779	ILE	5.4
1	В	810	LEU	5.4
1	D	472	ASP	5.2
1	D	368	ALA	5.1
1	D	471	HIS	4.9
1	В	368	ALA	4.9
1	А	394	ASN	4.4
1	D	394	ASN	4.3
1	D	473	LYS	4.2
1	В	471	HIS	3.9
1	А	469	THR	3.8
1	С	939	THR	3.6
1	В	470	LYS	3.6
1	D	501	VAL	3.5
1	D	367	ASN	3.5
1	D	810	LEU	3.5
1	В	808	ARG	3.5
1	В	793	LEU	3.5
1	D	811	GLU	3.5



3ZHQ
------

Mol	Chain	Res	Type	RSRZ
1	В	775	THR	3.4
1	D	937	THR	3.4
1	С	394	ASN	3.3
1	А	415	LEU	3.3
1	В	779	ILE	3.3
1	С	421	VAL	3.2
1	D	470	LYS	3.2
1	D	370	VAL	3.2
1	С	792	ALA	3.2
1	А	779	ILE	3.2
1	D	813	HIS	3.2
1	D	371	ILE	3.2
1	В	797	GLN	3.2
1	D	797	GLN	3.2
1	С	785	SER	3.1
1	В	786	MET	3.1
1	В	501	VAL	3.1
1	С	938	PRO	3.1
1	В	805	ASN	3.0
1	В	858	PRO	3.0
1	А	471	HIS	3.0
1	А	810	LEU	3.0
1	D	369	ARG	3.0
1	В	809	GLU	2.9
1	В	1102	GLY	2.9
1	D	934	PRO	2.9
1	D	784	ILE	2.9
1	D	786	MET	2.9
1	С	937	THR	2.8
1	D	935	ASP	2.8
1	С	786	MET	2.8
1	В	473	LYS	2.8
1	С	368	ALA	2.7
1	D	805	ASN	2.7
1	В	367	ASN	2.7
1	В	559	SER	2.7
1	D	415	LEU	2.7
1	D	953	PHE	2.7
1	С	420	LYS	2.6
1	D	632	ASP	2.6
1	D	420	LYS	2.6
1	В	419	PHE	2.6



3ZHQ
------

Mol	Chain	Res	Type	RSRZ	
1	С	561	PHE	2.6	
1	С	805 ASN		2.6	
1	С	931	ALA	2.6	
1	А	1224 THR		2.5	
1	D	419	PHE	2.5	
1	А	413	TRP	2.5	
1	В	413	TRP	2.5	
1	С	477	ALA	2.5	
1	D	416	ASP	2.5	
1	С	652	GLY	2.5	
1	D	938	PRO	2.5	
1	А	472	ASP	2.5	
1	С	436	LEU	2.4	
1	C	371	ILE	2.4	
1	В	811	GLU	2.4	
1	В	394	ASN	2.4	
1	В	647	ALA	2.4	
1	В	794	ARG	2.4	
1	С	793	LEU	2.4	
1	В	802	ARG	2.3	
1	В	556	GLN	2.3	
1	С	813	HIS	2.3	
1	D	803	VAL	2.3	
1	В	801	GLU	2.3	
1	В	803	VAL	2.3	
1	В	807	VAL	2.3	
1	С	953	PHE	2.3	
1	В	469	THR	2.3	
1	D	436	LEU	2.3	
1	С	702	ILE	2.2	
1	С	1210	SER	2.2	
1	В	777	ALA	2.2	
1	С	784	ILE	2.2	
1	С	775	THR	2.2	
1	С	476	VAL	2.2	
1	А	395	THR	2.2	
1	А	953	PHE	2.2	
1	D	500	TYR	2.2	
1	С	778	LEU	2.2	
1	В	953	PHE	2.1	
1	С	795	ASP	2.1	
1	С	954	ALA	2.1	



Mol	Chain	Res	Type	RSRZ
1	D	796	TYR	2.1
1	С	419 PHE		2.1
1	В	417	ARG	2.1
1	D	936	GLY	2.1
1	D	395	THR	2.1
1	А	419	PHE	2.1
1	В	795	ASP	2.1
1	А	785	SER	2.1
1	В	432	LEU	2.1
1	D	804	PHE	2.1
1	А	501	VAL	2.1
1	А	1213	VAL	2.1
1	В	415	LEU	2.1
1	D	791	ASP	2.0
1	D	469	THR	2.0
1	D	1216	VAL	2.0
1	А	397	PHE	2.0
1	А	937	THR	2.0
1	С	395	THR	2.0
1	А	933	ASN	2.0
1	В	785	SER	2.0
1	А	935	ASP	2.0
1	С	986	SER	2.0

Continued from previous page...

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Continuea from previous page								
Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(A^2)$	Q < 0.9
Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(A^2)$	$\mathbf{Q} < 0.9$
4	CA	С	2003	1/1	0.95	0.04	42,42,42,42	0
3	MG	А	2002	1/1	0.96	0.15	20,20,20,20	0
2	TPP	А	2001	26/26	0.97	0.15	16,26,33,34	0
3	MG	D	2002	1/1	0.97	0.12	23,23,23,23	0
2	TPP	D	2001	26/26	0.97	0.18	23,33,36,38	0
2	TPP	В	2001	26/26	0.98	0.16	20,31,37,38	0
2	TPP	С	2001	26/26	0.98	0.15	23,34,39,40	0
3	MG	В	2002	1/1	0.99	0.08	18,18,18,18	0
4	CA	А	2003	1/1	0.99	0.09	32,32,32,32	0
4	CA	В	2003	1/1	0.99	0.06	38,38,38,38	0
3	MG	С	2002	1/1	0.99	0.14	$15,\!15,\!15,\!15$	0
4	CA	D	2003	1/1	0.99	0.06	40,40,40,40	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.











## 6.5 Other polymers (i)

There are no such residues in this entry.

