



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 22, 2024 – 08:17 PM EDT

PDB ID : 4ZHS  
Title : Crystal Structure of Aspartate Semialdehyde Dehydrogenase from *Trichophyton rubrum*  
Authors : Li, Q.; Cui, S.  
Deposited on : 2015-04-27  
Resolution : 2.60 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	379	 4% 77% 16% • 7%
1	B	379	 4% 74% 17% • 8%
1	C	379	 6% 73% 20% • 6%
1	D	379	 4% 74% 19% • 6%
1	E	379	 5% 72% 19% • 8%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	379	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	F	401	-	-	X	-

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 32241 atoms, of which 15998 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aspartate Semialdehyde Dehydrogenase.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S	Se			
1	D	356	Total 5365	C 1681	H 2697	N 464	O 504	S 10	Se 9	0	0	0
1	C	355	Total 5352	C 1677	H 2692	N 463	O 501	S 10	Se 9	0	0	0
1	B	349	Total 5278	C 1656	H 2653	N 459	O 491	S 10	Se 9	0	0	0
1	E	348	Total 5253	C 1650	H 2641	N 454	O 489	S 10	Se 9	0	0	0
1	A	354	Total 5341	C 1674	H 2687	N 462	O 499	S 10	Se 9	0	0	0
1	F	346	Total 5221	C 1637	H 2628	N 452	O 485	S 10	Se 9	0	0	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	D	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	E	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	F	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	81	Total	O	0	0
			81	81		
3	C	66	Total	O	0	0
			66	66		
3	B	67	Total	O	0	0
			67	67		
3	E	66	Total	O	0	0
			66	66		
3	A	84	Total	O	0	0
			84	84		
3	F	37	Total	O	0	0
			37	37		








● Molecule 1: Aspartate Semialdehyde Dehydrogenase




● Molecule 1: Aspartate Semialdehyde Dehydrogenase



● Molecule 1: Aspartate Semialdehyde Dehydrogenase







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	157.50Å 157.50Å 188.40Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	27.62 – 2.60 27.62 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.6 (27.62-2.60) 99.7 (27.62-2.60)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.90 (at 2.61Å)	Xtrriage
Refinement program	PHENIX 1.7.3_928	Depositor
R, $R_{free}$	0.214 , 0.257 0.225 , 0.266	Depositor DCC
$R_{free}$ test set	4138 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	48.2	Xtrriage
Anisotropy	0.131	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 40.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.035 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	32241	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	61.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 34.72 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.5282e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	0/2695	0.54	0/3640
1	B	0.37	0/2663	0.52	0/3592
1	C	0.38	0/2701	0.54	0/3648
1	D	0.39	0/2709	0.55	0/3659
1	E	0.35	0/2651	0.52	0/3580
1	F	0.34	0/2630	0.53	0/3549
All	All	0.37	0/16049	0.53	0/21668

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	171	GLY	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2654	2687	2681	54	0
1	B	2625	2653	2647	54	0
1	C	2660	2692	2686	62	0
1	D	2668	2697	2690	78	0
1	E	2612	2641	2636	70	1
1	F	2593	2628	2622	86	0
2	A	5	0	0	1	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	5	0	0	0	0
2	E	5	0	0	0	0
2	F	5	0	0	2	0
3	A	84	0	0	15	0
3	B	67	0	0	15	0
3	C	66	0	0	14	0
3	D	81	0	0	9	0
3	E	66	0	0	17	1
3	F	37	0	0	15	0
All	All	16243	15998	15962	397	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 397 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:55:LYS:NZ	1:F:196:ASP:OD1	1.89	1.04
1:A:18:VAL:N	3:A:501:HOH:O	1.90	1.03
1:B:55:LYS:HE2	3:B:501:HOH:O	1.63	0.99
1:E:38:GLY:O	3:E:501:HOH:O	1.85	0.94
1:E:240:VAL:O	3:E:502:HOH:O	1.90	0.88

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:323:ARG:HH21	3:E:502:HOH:O[4_465]	1.53	0.07

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	350/379 (92%)	327 (93%)	20 (6%)	3 (1%)	17	35
1	B	339/379 (89%)	323 (95%)	14 (4%)	2 (1%)	25	47
1	C	351/379 (93%)	334 (95%)	14 (4%)	3 (1%)	17	35
1	D	352/379 (93%)	333 (95%)	15 (4%)	4 (1%)	14	30
1	E	341/379 (90%)	325 (95%)	13 (4%)	3 (1%)	17	35
1	F	338/379 (89%)	310 (92%)	23 (7%)	5 (2%)	10	21
All	All	2071/2274 (91%)	1952 (94%)	99 (5%)	20 (1%)	15	32

5 of 20 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	189	TYR
1	C	189	TYR
1	E	189	TYR
1	A	189	TYR
1	F	189	TYR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	292/303 (96%)	290 (99%)	2 (1%)	84	94
1	B	288/303 (95%)	282 (98%)	6 (2%)	53	77
1	C	293/303 (97%)	284 (97%)	9 (3%)	40	66

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	294/303 (97%)	290 (99%)	4 (1%)	67	85
1	E	287/303 (95%)	283 (99%)	4 (1%)	67	85
1	F	286/303 (94%)	280 (98%)	6 (2%)	53	77
All	All	1740/1818 (96%)	1709 (98%)	31 (2%)	59	80

5 of 31 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	47	LYS
1	F	78	SER
1	B	213	SER
1	F	198	PHE
1	A	153	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	56	GLN
1	B	109	ASN
1	B	151	ASN
1	F	109	ASN
1	F	339	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	F	401	-	4,4,4	0.16	0	6,6,6	0.19	0
2	SO4	D	401	-	4,4,4	0.21	0	6,6,6	0.36	0
2	SO4	C	401	-	4,4,4	0.16	0	6,6,6	0.31	0
2	SO4	A	401	-	4,4,4	0.13	0	6,6,6	0.22	0
2	SO4	E	401	-	4,4,4	0.14	0	6,6,6	0.25	0
2	SO4	B	401	-	4,4,4	0.22	0	6,6,6	0.24	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	401	SO4	2	0
2	A	401	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	345/379 (91%)	0.15	16 (4%) 32 26	25, 45, 86, 115	0
1	B	340/379 (89%)	0.16	16 (4%) 31 25	25, 48, 87, 117	1 (0%)
1	C	346/379 (91%)	0.17	21 (6%) 21 16	24, 48, 90, 121	0
1	D	347/379 (91%)	0.15	17 (4%) 29 23	24, 45, 93, 117	0
1	E	339/379 (89%)	0.23	19 (5%) 24 19	26, 53, 91, 123	0
1	F	337/379 (88%)	0.46	26 (7%) 13 10	36, 62, 92, 118	0
All	All	2054/2274 (90%)	0.22	115 (5%) 24 19	24, 52, 91, 123	1 (0%)

The worst 5 of 115 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	189	TYR	7.4
1	D	189	TYR	6.7
1	D	50	ASP	5.8
1	A	53	ARG	5.8
1	F	67	LEU	5.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	E	401	5/5	0.92	0.13	53,60,69,70	0
2	SO4	C	401	5/5	0.93	0.12	43,59,64,68	0
2	SO4	A	401	5/5	0.95	0.12	52,56,61,62	0
2	SO4	F	401	5/5	0.96	0.14	64,67,74,84	0
2	SO4	B	401	5/5	0.97	0.09	47,53,59,64	0
2	SO4	D	401	5/5	0.97	0.13	46,49,53,65	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.