



# wwPDB EM Validation Summary Report ⓘ

Jun 18, 2024 – 04:24 PM JST

PDB ID : 8ZJ2  
EMDB ID : EMD-60136  
Title : Cryo-EM structure of the RhoG/DOCK5/ELMO1/Rac1 complex  
Authors : Kukimoto-Niino, M.; Katsura, K.; Ishizuka-Katsura, Y.; Mishima-Tsumagari, C.; Yonemochi, M.; Inoue, M.; Nakagawa, R.; Kaushik, R.; Zhang, K.Y.J.; Shirouzu, M.  
Deposited on : 2024-05-14  
Resolution : 4.66 Å (reported)  
Based on initial models : 6IE1, 7Y4A, 7DPA

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev92  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
MolProbity : 4.02b-467  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 4.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	1648	 31% 68%
1	F	1648	 18% 38% 61%
2	C	184	 33% 63%
2	G	184	 32% 64%
3	A	733	 48% 36% 62%
3	E	733	 9% 18% 73%
4	D	203	 21% 33% 55% 11%

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 38587 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Deducator of cytokinesis protein 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	B	1642	Total	C	N	O	S	0	0
			13436	8618	2264	2484	70		
1	F	1642	Total	C	N	O	S	0	0
			13436	8618	2264	2484	70		

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-5	GLY	-	expression tag	UNP Q9H7D0
B	-4	GLY	-	expression tag	UNP Q9H7D0
B	-3	SER	-	expression tag	UNP Q9H7D0
B	-2	GLY	-	expression tag	UNP Q9H7D0
B	-1	GLY	-	expression tag	UNP Q9H7D0
B	0	SER	-	expression tag	UNP Q9H7D0
B	1285	ARG	LYS	variant	UNP Q9H7D0
F	-5	GLY	-	expression tag	UNP Q9H7D0
F	-4	GLY	-	expression tag	UNP Q9H7D0
F	-3	SER	-	expression tag	UNP Q9H7D0
F	-2	GLY	-	expression tag	UNP Q9H7D0
F	-1	GLY	-	expression tag	UNP Q9H7D0
F	0	SER	-	expression tag	UNP Q9H7D0
F	1285	ARG	LYS	variant	UNP Q9H7D0

- Molecule 2 is a protein called Ras-related C3 botulinum toxin substrate 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	C	177	Total	C	N	O	S	0	0
			1385	890	228	259	8		
2	G	177	Total	C	N	O	S	0	0
			1385	890	228	259	8		

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-6	GLY	-	expression tag	UNP P63000
C	-5	SER	-	expression tag	UNP P63000
C	-4	SER	-	expression tag	UNP P63000
C	-3	GLY	-	expression tag	UNP P63000
C	-2	SER	-	expression tag	UNP P63000
C	-1	SER	-	expression tag	UNP P63000
C	0	GLY	-	expression tag	UNP P63000
C	15	ALA	GLY	engineered mutation	UNP P63000
G	-6	GLY	-	expression tag	UNP P63000
G	-5	SER	-	expression tag	UNP P63000
G	-4	SER	-	expression tag	UNP P63000
G	-3	GLY	-	expression tag	UNP P63000
G	-2	SER	-	expression tag	UNP P63000
G	-1	SER	-	expression tag	UNP P63000
G	0	GLY	-	expression tag	UNP P63000
G	15	ALA	GLY	engineered mutation	UNP P63000

- Molecule 3 is a protein called Engulfment and cell motility protein 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	E	199	1617	1023	279	305	10	0	0
3	A	727	5879	3721	1009	1108	41	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	-5	GLY	-	expression tag	UNP Q92556
E	-4	GLY	-	expression tag	UNP Q92556
E	-3	SER	-	expression tag	UNP Q92556
E	-2	GLY	-	expression tag	UNP Q92556
E	-1	GLY	-	expression tag	UNP Q92556
E	0	SER	-	expression tag	UNP Q92556
A	-5	GLY	-	expression tag	UNP Q92556
A	-4	GLY	-	expression tag	UNP Q92556
A	-3	SER	-	expression tag	UNP Q92556
A	-2	GLY	-	expression tag	UNP Q92556
A	-1	GLY	-	expression tag	UNP Q92556
A	0	SER	-	expression tag	UNP Q92556

- Molecule 4 is a protein called Rho-related GTP-binding protein RhoG.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	D	181	1416	897	248	263	8	0	0


There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-6	GLY	-	expression tag	UNP P84095
D	-5	SER	-	expression tag	UNP P84095
D	-4	SER	-	expression tag	UNP P84095
D	-3	GLY	-	expression tag	UNP P84095
D	-2	SER	-	expression tag	UNP P84095
D	-1	SER	-	expression tag	UNP P84095
D	0	GLY	-	expression tag	UNP P84095
D	61	LEU	GLN	engineered mutation	UNP P84095
D	185	SER	-	expression tag	UNP P84095
D	186	GLY	-	expression tag	UNP P84095
D	187	PRO	-	expression tag	UNP P84095
D	188	SER	-	expression tag	UNP P84095
D	189	SER	-	expression tag	UNP P84095
D	190	GLY	-	expression tag	UNP P84095
D	191	GLU	-	expression tag	UNP P84095
D	192	ASN	-	expression tag	UNP P84095
D	193	LEU	-	expression tag	UNP P84095
D	194	TYR	-	expression tag	UNP P84095
D	195	PHE	-	expression tag	UNP P84095
D	196	GLN	-	expression tag	UNP P84095

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
5	D	1	Total	Mg	0
			1	1	

- Molecule 6 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>14</sub>P<sub>3</sub>) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
6	D	1	32	10	5	14	3	0

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Deducator of cytokinesis protein 5











● Molecule 1: Dedicator of cytokinesis protein 5



L1599	V1533	R1463	S1321	I1241	L1175	L1110	S1037	H968	Q904	S837	Y774
L1600	Q1534	Y1464	K1322	R1242	L1176	E1111	F1038	Y969	L905	V838	L775
E1602	S1466	R1466	L1324	Y1243	E1178	L1113	E1039	S970	L906	L839	L776
I1604	A1537	P1467	S1395	Y1245	R1181	T1114	L1041	H971	N908	C841	G779
R1605	W1538	F1468	E1326	R1248	K1182	T1115	Q1042	Y972	I909	K842	K782
I1606	D1539	R1469	T1327	D1249	H1183	E1117	W1043	S974	L910	F843	D783
H1607	R1540	K1470	E1328	L1250	H1184	W1118	M1044	T975	E911	Q845	F784
G1608	S1541	K1473	V1332	R1252	Y1185	L1119	F1047	F976	L913	Q846	E785
E1609	Q1401	H1478	E1259	R1252	Y1186	L1120	Y1046	R979	D914	S846	D786
K1610	Q1405	E1478	Y1335	S1252	S1187	R1121	H1043	I982	R915	E786	F787
L1611	A1405	E1478	E1336	E1259	S1188	K1122	K916	L851	F888	F787	F788
E1613	E1406	T1481	E1336	Y1262	E1191	A1123	L1064	D917	L852	L788	F789
L1614	M1482	M1482	G1337	Y1262	V1192	T1124	T1065	V918	V918	S789	S790
K1615	L1483	L1338	L1337	Y1262	W1192	L1125	H1056	R854	R854	N789	N790
K1616	I1484	G1339	G1339	Q1272	F1193	L1126	E1057	T921	T921	I791	I791
P1617	E1485	M1340	M1340	W1273	A1194	I1127	S1058	A922	A922	Q792	Q792
L1618	R1486	L1341	L1341	S1274	L1195	F1128	L1069	V923	V923	W793	W793
H1619	E1416	L1342	L1342	D1275	L1196	F1129	Q1060	H924	H924	L794	L794
E1620	D1417	K1343	K1343	K1276	V1197	D1130	L1061	I925	I925	C859	C859
R1621	K1418	K1344	K1344	P1277	S1198	M1131	E1062	Q826	Q826	M860	M860
L1622	T1490	R1345	R1345	C1278	S1199	M1132	T1063	L927	L927	T861	T861
S1623	T1491	A1346	A1346	V1279	L1200	Q1133	F1064	I928	I928	K862	K862
F1626	S1421	S1347	P1280	P1280	L1201	E1135	S1065	I997	I997	M799	M799
E1628	Q1424	H1251	H1251	H1251	L1204	E1136	Q1066	Y1002	Y1002	M800	M800
K1630	Y1425	L1282	L1282	L1282	L1205	M1137	A1067	R831	R831	L801	L801
E1631	M1426	L1283	L1283	Q1284	D1206	F1138	K1068	D1005	D1005	R804	R804
K1632	F1429	Q1284	Q1284	S1139	Y1207	S1139	R1069	W1006	W1006	P805	P805
L1633	M1435	S1287	S1287	T1209	R1208	G1140	I1072	V1007	V1007	L806	L806
E1634	Q1369	Y1288	Y1288	I1210	I1210	H1145	G1077	M1009	M1009	E807	E807
K1635	P1361	Y1289	Y1289	M1211	M1211	M1146	D1078	M1010	M1010	E808	E808
H1636	Y1362	V1290	V1290	Q1213	Q1213	F1147	M1079	T1012	T1012	A809	A809
Y1637	V1366	Q1293	Q1293	S1216	S1216	E1148	R1080	Q1013	Q1013	V810	V810
G1638	G1366	E1294	E1294	K1217	K1217	M1149	K1081	M1014	M1014	K811	K811
L1642	G1366	Q1302	Q1302	E1218	E1218	E1150	E1082	R1015	R1015	I812	I812
	R1375	I1304	I1304	R1220	R1220	I1152	I1083	F1017	F1017	K813	K813
	R1377	L1305	L1305	M1219	M1219	I1152	R1086	L1018	L1018	G814	G814
	M1378	S1306	S1306	R1221	R1221	D1156	I1087	R1019	R1019	L817	L817
	K1379	Y1307	Y1307	S1222	S1222	E1157	D1089	R1020	R1020	K818	K818
	I1380	K1310	K1310	C1223	C1223	E1158	M1090	H1021	H1021	Y819	Y819
	F1381	G1311	G1311	T1224	T1224	V1159	Y1091	I1022	I1022	L820	L820
	R1384	E1316	E1316	V1225	V1225	G1162	Y1092	F1024	F1024	P821	P821
	K1386	K1317	K1317	M1229	M1229	G1163	H1097	A1025	A1025	S822	S822
	E1387	I1318	I1318	F1230	F1230	G1164	K1098	E1026	E1026	I823	I823
	E1388	K1319	K1319	Y1231	Y1231	D1165	I1099	V1027	V1027	I824	I824
	E1389	L1320	L1320	K1232	K1232	Q1167	I1100	L1028	L1028	N825	N825
				E1233	E1233	Y1168	F1101	T1029	T1029	D826	D826
				K1234	K1234	Y1169	I1102	M957	M957	V827	V827
				R1236	R1236	W1170	M1105	L1890	L1890	K828	K828
				E1237	E1237	L1171	P1108	M957	M957	L829	L829
				I1318	I1318	L1172	I1109	L1891	L1891	V830	V830
				K1319	K1319	E1173	Q1035	K896	K896	D832	D832
				L1320	L1320	K1174	I1109	D898	D898	P833	P833
								M964	M964	Y834	Y834
										E835	E835
										L836	L836

● Molecule 2: Ras-related C3 botulinum toxin substrate 1







● Molecule 2: Ras-related C3 botulinum toxin substrate 1





● Molecule 3: Engulfment and cell motility protein 1






• Molecule 3: Engulfment and cell motility protein 1





● Molecule 4: Rho-related GTP-binding protein RhoG



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	169096	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	50	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	64000	Depositor
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.062	Depositor
Minimum map value	-0.021	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.01	Depositor
Map size (Å)	452.2, 452.2, 452.2	wwPDB
Map dimensions	340, 340, 340	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.33, 1.33, 1.33	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	B	0.41	0/13722	0.58	1/18514 (0.0%)
1	F	0.34	0/13722	0.55	2/18514 (0.0%)
2	C	0.36	0/1415	0.55	0/1924
2	G	0.34	0/1415	0.54	0/1924
3	A	0.30	0/5992	0.55	0/8086
3	E	0.30	0/1650	0.56	0/2230
4	D	0.30	0/1449	0.51	0/1977
All	All	0.36	0/39365	0.56	3/53169 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	676	LEU	CA-CB-CG	6.09	129.30	115.30
1	B	992	MET	CA-CB-CG	-5.47	104.00	113.30
1	F	96	ARG	CB-CG-CD	5.10	124.86	111.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	110	LYS	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	13436	0	13516	1324	0
1	F	13436	0	13516	1068	0
2	C	1385	0	1407	117	0
2	G	1385	0	1407	139	0
3	A	5879	0	5902	479	0
3	E	1617	0	1625	159	0
4	D	1416	0	1413	123	0
5	D	1	0	0	0	0
6	D	32	0	12	4	0
All	All	38587	0	38798	3295	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 43.

The worst 5 of 3295 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:1028:LEU:O	1:F:1032:PHE:HB2	1.63	0.98
1:F:1217:LYS:HA	1:F:1220:ARG:HE	1.31	0.94
3:A:302:PHE:HB3	3:A:431:GLY:H	1.31	0.94
1:B:1462:PHE:HB2	1:B:1489:TYR:HB2	1.46	0.94
1:F:740:TYR:HA	1:F:749:LYS:HD3	1.47	0.94

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM



entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	1640/1648 (100%)	1416 (86%)	224 (14%)	0	100	100
1	F	1640/1648 (100%)	1466 (89%)	174 (11%)	0	100	100
2	C	175/184 (95%)	156 (89%)	19 (11%)	0	100	100
2	G	175/184 (95%)	157 (90%)	18 (10%)	0	100	100
3	A	725/733 (99%)	657 (91%)	68 (9%)	0	100	100
3	E	197/733 (27%)	162 (82%)	35 (18%)	0	100	100
4	D	179/203 (88%)	161 (90%)	18 (10%)	0	100	100
All	All	4731/5333 (89%)	4175 (88%)	556 (12%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	1495/1497 (100%)	1490 (100%)	5 (0%)	92	95
1	F	1495/1497 (100%)	1489 (100%)	6 (0%)	91	94
2	C	153/157 (98%)	153 (100%)	0	100	100
2	G	153/157 (98%)	151 (99%)	2 (1%)	69	82
3	A	662/664 (100%)	659 (100%)	3 (0%)	88	93
3	E	184/664 (28%)	183 (100%)	1 (0%)	88	93
4	D	157/174 (90%)	155 (99%)	2 (1%)	69	82
All	All	4299/4810 (89%)	4280 (100%)	19 (0%)	91	94

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	218	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
4	D	66	ARG
4	D	147	LYS
3	A	516	LYS
1	F	743	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 53 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	793	GLN
1	F	1526	ASN
3	A	703	ASN
1	F	799	ASN
1	F	1035	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	GTP	D	202	4,5	26,34,34	1.11	2 (7%)	32,54,54	1.70	7 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GTP	D	202	4,5	-	2/18/38/38	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	D	202	GTP	C5-C6	-3.95	1.39	1.47
6	D	202	GTP	C2-N3	2.15	1.38	1.33

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	D	202	GTP	PA-O3A-PB	-5.03	115.55	132.83
6	D	202	GTP	PB-O3B-PG	-3.40	121.15	132.83
6	D	202	GTP	C5-C6-N1	3.20	119.61	113.95
6	D	202	GTP	C3'-C2'-C1'	2.95	105.42	100.98
6	D	202	GTP	C8-N7-C5	2.88	108.48	102.99

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	D	202	GTP	PB-O3B-PG-O3G
6	D	202	GTP	PB-O3B-PG-O1G


There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	D	202	GTP	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-60136. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 170




Y Index: 170




Z Index: 170


### 6.2.2 Raw map



X Index: 170



Y Index: 170




Z Index: 170

The images above show central slices of the map in three orthogonal directions.




## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 118




Y Index: 169




Z Index: 149


### 6.3.2 Raw map



X Index: 104



Y Index: 170




Z Index: 149


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X



Y



Z

### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.




## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.01. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 551  $\text{nm}^3$ ; this corresponds to an approximate mass of 498 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum [i](#)




\*Reported resolution corresponds to spatial frequency of 0.215 Å<sup>-1</sup>

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.215 Å<sup>-1</sup>

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.66	-	-
Author-provided FSC curve	4.65	7.27	5.03
Unmasked-calculated*	6.99	8.40	7.16

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 6.99 differs from the reported value 4.66 by more than 10 %

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-60136 and PDB model 8ZJ2. Per-residue inclusion information can be found in section 3 on page 7.

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.01 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.01).



















## 9.4 Atom inclusion [i](#)



At the recommended contour level, 83% of all backbone atoms, 80% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.01) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8010	 0.2310
A	 0.4950	 0.1540
B	 0.9400	 0.2700
C	 0.9660	 0.3030
D	 0.6550	 0.1510
E	 0.8810	 0.2070
F	 0.7700	 0.2260
G	 0.9550	 0.2650

