



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 25, 2022 – 08:31 pm BST

PDB ID : 7ZKV  
Title : F231A variant of the CODH/ACS complex of *C. hydrogenoformans*  
Authors : Ruickoldt, J.; Jeoung, J.-H.; Basak, Y.; Domnik, L.; Dobbek, H.  
Deposited on : 2022-04-13  
Resolution : 2.07 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.31.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2684 (2.08-2.04)
Clashscore	141614	2801 (2.08-2.04)
Ramachandran outliers	138981	2768 (2.08-2.04)
Sidechain outliers	138945	2768 (2.08-2.04)
RSRZ outliers	127900	2646 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	669	 93% 7%
2	B	730	 94% 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	OH	A	705	-	-	X	-
6	MRD	A	706	-	-	X	-
8	ACT	B	804	-	-	X	-

## 2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 11973 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Carbon monoxide dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	669	5141	3258	887	961	35	0	4	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	17	ASP	GLU	conflict	UNP A0A1L8D0M5
A	29	ILE	THR	conflict	UNP A0A1L8D0M5
A	73	GLN	MET	conflict	UNP A0A1L8D0M5
A	120	ALA	THR	conflict	UNP A0A1L8D0M5
A	153	THR	ILE	conflict	UNP A0A1L8D0M5
A	159	MET	LEU	conflict	UNP A0A1L8D0M5
A	199	GLU	ASP	conflict	UNP A0A1L8D0M5
A	205	SER	ALA	conflict	UNP A0A1L8D0M5
A	220	ILE	MET	conflict	UNP A0A1L8D0M5
A	231	ALA	PHE	engineered mutation	UNP A0A1L8D0M5
A	389	ILE	VAL	conflict	UNP A0A1L8D0M5
A	393	LEU	PHE	conflict	UNP A0A1L8D0M5
A	494	THR	ALA	conflict	UNP A0A1L8D0M5
A	602	THR	SER	conflict	UNP A0A1L8D0M5


- Molecule 2 is a protein called CO-methylating acetyl-CoA synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	730	5809	3727	974	1079	29	8	6	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	733	ARG	-	expression tag	UNP Q3ACS4
B	734	SER	-	expression tag	UNP Q3ACS4

- Molecule 3 is Fe(3)-Ni(1)-S(4) cluster (three-letter code: RQM) (formula:  $\text{Fe}_4\text{NiS}_4$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	Fe	Ni	S		
3	A	1	9	4	1	4	0	0

- Molecule 4 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula:  $\text{Fe}_4\text{S}_4$ ) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Fe S		
4	A	1	4	2 2	0	0

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	Fe	S		
4	A	1	8	4	4	0	0
4	B	1	8	4	4	0	0

- Molecule 5 is HYDROXIDE ION (three-letter code: OH) (formula: HO).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	1	1	1	0	0
5	A	1	1	1	0	0

- Molecule 6 is (4R)-2-METHYLPENTANE-2,4-DIOL (three-letter code: MRD) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 8 6 2	0	0
6	A	1	Total C O 8 6 2	0	0

- Molecule 7 is NICKEL (II) ION (three-letter code: NI) (formula: Ni) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	2	Total Ni 2 2	0	0

- Molecule 8 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	B	1	Total C O 4 2 2	0	0

- Molecule 9 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	B	1	Total Na 1 1	0	0

- Molecule 10 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	403	Total O 403 403	0	0
10	B	566	Total O 566 566	0	0




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Carbon monoxide dehydrogenase



- Molecule 2: CO-methylating acetyl-CoA synthase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	142.64Å 142.64Å 291.31Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	42.38 – 2.07 48.55 – 2.07	Depositor EDS
% Data completeness (in resolution range)	99.7 (42.38-2.07) 99.7 (48.55-2.07)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.54 (at 2.07Å)	Xtrriage
Refinement program	PHENIX 1.11.1_2575	Depositor
R, $R_{free}$	0.173 , 0.214 0.173 , 0.216	Depositor DCC
$R_{free}$ test set	1124 reflections (1.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.0	Xtrriage
Anisotropy	0.323	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	11973	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.26% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: RQM, NA, ACT, MRD, SF4, OH, NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.45	1/5252 (0.0%)	0.60	1/7109 (0.0%)
2	B	0.45	0/5961	0.63	4/8068 (0.0%)
All	All	0.45	1/11213 (0.0%)	0.62	5/15177 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	1
All	All	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	467	CYS	CB-SG	6.01	1.92	1.82

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	224	TYR	CB-CG-CD2	-7.59	116.44	121.00
2	B	262	ASP	CB-CG-OD1	5.99	123.69	118.30
2	B	224	TYR	CB-CG-CD1	5.48	124.29	121.00
1	A	590	TRP	CA-CB-CG	5.23	123.64	113.70
2	B	601	PHE	CB-CG-CD2	-5.05	117.26	120.80

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	466	GLY	Peptide
2	B	601	PHE	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5141	0	5165	37	0
2	B	5809	0	5795	33	0
3	A	9	0	0	0	0
4	A	12	0	0	0	0
4	B	8	0	0	0	0
5	A	2	0	0	2	0
6	A	16	0	28	14	0
7	B	2	0	0	0	0
8	B	4	0	3	2	0
9	B	1	0	0	0	0
10	A	403	0	0	5	3
10	B	566	0	0	12	1
All	All	11973	0	10991	72	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:167:GLU:OE1	10:A:801:HOH:O	1.82	0.96
1:A:616:THR:HG22	6:A:706:MRD:H3C1	1.53	0.90
2:B:734:SER:O	10:B:901:HOH:O	1.92	0.87
2:B:306:GLN:NE2	10:B:906:HOH:O	2.13	0.81
2:B:380:GLU:OE2	10:B:903:HOH:O	2.02	0.76
1:A:546[B]:CYS:SG	5:A:705:OH:O	2.44	0.76
1:A:163:LYS:O	1:A:167:GLU:HG3	1.86	0.75
2:B:368:LYS:NZ	10:B:902:HOH:O	1.94	0.75
1:A:234:LEU:HD21	6:A:706:MRD:HMC3	1.69	0.74
2:B:362:ASP:OD2	10:B:904:HOH:O	2.07	0.73

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:530:LEU:HD13	2:B:601:PHE:HB3	1.69	0.73
2:B:144:VAL:HG22	2:B:224:TYR:CE1	2.26	0.70
2:B:107:GLU:OE1	10:B:905:HOH:O	2.09	0.69
1:A:157:ASP:OD1	1:A:157:ASP:N	2.23	0.69
1:A:624:TYR:HE2	6:A:707:MRD:H5C3	1.58	0.69
1:A:612:PHE:O	1:A:616:THR:HG23	1.94	0.67
2:B:144:VAL:HG22	2:B:224:TYR:CD1	2.29	0.67
1:A:234:LEU:HD21	6:A:706:MRD:CM	2.29	0.63
2:B:5:ILE:N	10:B:911:HOH:O	2.31	0.62
2:B:688:ARG:NH1	2:B:691:GLU:OE2	2.32	0.62
2:B:260:GLU:OE1	10:B:907:HOH:O	2.17	0.59
1:A:121:HIS:NE2	1:A:125:GLU:OE2	2.36	0.58
2:B:173:LYS:NZ	2:B:177:ASP:OD2	2.36	0.58
1:A:383:TYR:OH	10:A:802:HOH:O	2.16	0.57
1:A:230:THR:CG2	6:A:706:MRD:HMC1	2.36	0.56
2:B:338:LYS:NZ	10:B:918:HOH:O	2.38	0.56
2:B:139:ILE:HG21	2:B:224:TYR:HE2	1.72	0.54
2:B:139:ILE:HG21	2:B:224:TYR:CE2	2.43	0.54
1:A:616:THR:HA	6:A:706:MRD:O2	2.09	0.53
1:A:101:VAL:HG11	6:A:706:MRD:H5C1	1.90	0.53
1:A:85:LYS:NZ	10:A:808:HOH:O	2.43	0.52
1:A:624:TYR:CE2	6:A:707:MRD:H5C3	2.42	0.52
2:B:232:PHE:CG	8:B:804:ACT:H3	2.45	0.52
1:A:141:LYS:NZ	1:A:252:ASP:OD2	2.43	0.51
2:B:64:PHE:CE2	2:B:75:GLU:HG2	2.45	0.51
2:B:287:GLU:OE2	10:B:908:HOH:O	2.19	0.51
2:B:144:VAL:HA	2:B:224:TYR:CZ	2.46	0.51
1:A:247:ALA:HB3	1:A:547:VAL:HG21	1.93	0.51
2:B:592:GLU:HG2	2:B:734:SER:HB3	1.93	0.50
1:A:546[A]:CYS:HB2	5:A:705:OH:O	2.12	0.49
2:B:601:PHE:CD1	2:B:601:PHE:N	2.80	0.49
1:A:391:ASN:HB2	1:A:394:GLU:CD	2.33	0.48
1:A:440:VAL:HG22	1:A:530:LYS:HG3	1.94	0.48
1:A:616:THR:CG2	6:A:706:MRD:H3C1	2.36	0.48
2:B:38:TYR:CZ	2:B:42:LEU:HD11	2.49	0.48
1:A:391:ASN:O	1:A:394:GLU:HG2	2.14	0.47
6:A:707:MRD:H5C1	10:A:805:HOH:O	2.13	0.47
1:A:230:THR:HG23	6:A:706:MRD:H3C2	1.97	0.46
2:B:364:ILE:HG23	2:B:465:ARG:HD2	1.98	0.45
1:A:259:LYS:HA	1:A:259:LYS:HD3	1.81	0.45
6:A:707:MRD:C5	10:A:805:HOH:O	2.65	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:482:LEU:HD22	1:A:502:PHE:CD1	2.52	0.45
1:A:231:ALA:HB2	6:A:707:MRD:HMC1	1.99	0.45
1:A:143:ARG:NH1	1:A:156:LYS:O	2.50	0.45
2:B:232:PHE:CD1	8:B:804:ACT:H3	2.52	0.45
1:A:314:MET:CE	1:A:336:SER:HB2	2.47	0.44
2:B:144:VAL:HA	2:B:224:TYR:OH	2.17	0.44
1:A:112:HIS:ND1	1:A:240:ASP:OD1	2.47	0.44
1:A:302:LYS:HA	1:A:306:ALA:O	2.19	0.43
2:B:376:ILE:HD11	2:B:442:ALA:O	2.19	0.42
1:A:230:THR:HG23	6:A:706:MRD:HMC1	2.01	0.42
2:B:60:ASP:OD2	10:B:910:HOH:O	2.21	0.42
1:A:278:ALA:HA	1:A:312:VAL:O	2.20	0.42
2:B:556:LYS:O	2:B:557:GLU:HG2	2.19	0.42
1:A:160:THR:O	1:A:164:GLU:HG3	2.19	0.41
2:B:170:LYS:HA	2:B:170:LYS:HD3	1.90	0.41
1:A:39:LYS:NZ	1:A:43:ASP:OD2	2.45	0.41
1:A:280:HIS:O	1:A:350:VAL:HA	2.21	0.41
1:A:391:ASN:HB2	1:A:394:GLU:OE2	2.21	0.40
2:B:144:VAL:HG22	2:B:224:TYR:CZ	2.54	0.40
2:B:159:GLU:O	2:B:253:ALA:HA	2.21	0.40
2:B:173:LYS:HE2	10:B:963:HOH:O	2.21	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
10:A:1044:HOH:O	10:B:1364:HOH:O[6_455]	2.08	0.12
10:A:862:HOH:O	10:A:862:HOH:O[10_444]	2.11	0.09
10:A:862:HOH:O	10:A:1122:HOH:O[10_444]	2.17	0.03

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	671/669 (100%)	642 (96%)	27 (4%)	2 (0%)	41	32
2	B	734/730 (100%)	720 (98%)	11 (2%)	3 (0%)	34	25
All	All	1405/1399 (100%)	1362 (97%)	38 (3%)	5 (0%)	34	25

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	266	ASN
2	B	599	GLY
1	A	353	GLN
2	B	190	ASP
2	B	661	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	551/547 (101%)	549 (100%)	2 (0%)	91	91
2	B	618/612 (101%)	612 (99%)	6 (1%)	76	75
All	All	1169/1159 (101%)	1161 (99%)	8 (1%)	86	84

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ARG
1	A	184	TRP
2	B	224	TYR
2	B	262	ASP
2	B	401[A]	ASP
2	B	401[B]	ASP
2	B	486	LYS
2	B	601	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	147	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 2 are modelled with single atom and 3 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	RQM	A	701	1,5	0,12,12	-	-	-		
4	SF4	B	802	2	0,12,12	-	-	-		
4	SF4	A	703	1	0,12,12	-	-	-		
6	MRD	A	706	-	7,7,7	0.42	0	9,10,10	1.31	1 (11%)
4	SF4	A	702	1	0,4,12	-	-	-		
8	ACT	B	804	7	3,3,3	0.70	0	3,3,3	1.17	0
6	MRD	A	707	-	7,7,7	0.61	0	9,10,10	2.73	4 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	RQM	A	701	1,5	-	-	0/4/4/4
4	SF4	B	802	2	-	-	0/6/5/5
4	SF4	A	703	1	-	-	0/6/5/5
4	SF4	A	702	1	-	-	0/1/1/5
6	MRD	A	706	-	-	2/5/5/5	-
6	MRD	A	707	-	-	2/5/5/5	-

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	707	MRD	O2-C2-C3	-4.77	91.89	109.80
6	A	707	MRD	O2-C2-C1	4.65	123.00	108.08
6	A	707	MRD	CM-C2-C3	3.36	125.61	109.96
6	A	706	MRD	O2-C2-CM	-2.46	100.19	108.08
6	A	707	MRD	C5-C4-C3	2.38	122.90	111.69

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	707	MRD	C2-C3-C4-C5
6	A	706	MRD	O2-C2-C3-C4
6	A	706	MRD	C2-C3-C4-O4
6	A	707	MRD	C2-C3-C4-O4


There are no ring outliers.


3 monomers are involved in 16 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	706	MRD	9	0
8	B	804	ACT	2	0
6	A	707	MRD	5	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	669/669 (100%)	-0.30	8 (1%) 79 80	22, 37, 62, 85	0
2	B	730/730 (100%)	-0.57	5 (0%) 87 88	23, 35, 56, 82	0
All	All	1399/1399 (100%)	-0.44	13 (0%) 84 85	22, 36, 59, 85	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	575	TYR	3.2
2	B	363	GLU	3.1
1	A	412	GLY	3.1
2	B	5	ILE	2.7
1	A	375[A]	ASN	2.7
2	B	580	ARG	2.5
1	A	305	GLY	2.5
1	A	306	ALA	2.4
1	A	415	VAL	2.4
1	A	367	HIS	2.2
1	A	413	LYS	2.2
1	A	411	VAL	2.2
2	B	224	TYR	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	MRD	A	706	8/8	0.86	0.33	48,58,63,68	0
6	MRD	A	707	8/8	0.89	0.29	26,42,48,53	0
5	OH	A	705	1/1	0.91	0.28	43,43,43,43	0
5	OH	A	704	1/1	0.97	0.21	45,45,45,45	0
3	RQM	A	701	9/9	0.97	0.08	31,41,50,63	8
9	NA	B	805	1/1	0.97	0.04	31,31,31,31	0
4	SF4	B	802	8/8	0.99	0.05	28,30,32,33	0
4	SF4	A	702	4/8	0.99	0.06	29,30,30,31	0
8	ACT	B	804	4/4	0.99	0.12	30,32,33,34	0
4	SF4	A	703	8/8	0.99	0.08	26,29,29,30	0
7	NI	B	801	1/1	1.00	0.06	29,29,29,29	0
7	NI	B	803	1/1	1.00	0.04	35,35,35,35	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around RQM A 701:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around SF4 B 802:**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around SF4 A 702:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around SF4 A 703:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.