



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 14, 2023 – 12:30 PM JST

PDB ID : 5ZLR
Title : Structure of NeuC
Authors : Ko, T.P.; Hsieh, T.J.; Yang, C.S.; Chen, Y.
Deposited on : 2018-03-29
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	380	 22% 75% 21% . .
1	B	380	 2% 89% 7% .

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6367 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called GDP/UDP-N,N'-diacetylbaicillosamine 2-epimerase (Hydrolyzing).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	370	2905	1860	484	550	11	0	0	0
1	B	366	2879	1842	480	546	11	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	379	LEU	-	expression tag	UNP A0A154EJU5
A	380	GLU	-	expression tag	UNP A0A154EJU5
B	379	LEU	-	expression tag	UNP A0A154EJU5
B	380	GLU	-	expression tag	UNP A0A154EJU5

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is LITHIUM ION (three-letter code: LI) (formula: Li).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Li	0	0
			1	1		
3	B	1	Total	Li	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	252	Total	O	0	0
			252	252		
4	B	319	Total	O	0	0
			319	319		

4 Data and refinement statistics i

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	86.53Å 148.71Å 125.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.95 – 2.00 28.98 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (28.95-2.00) 99.8 (28.98-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.24 (at 2.00Å)	Xtrriage
Refinement program	PHENIX 1.10.1_2155	Depositor
R, R_{free}	0.166 , 0.213 0.166 , 0.213	Depositor DCC
R_{free} test set	2746 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	32.8	Xtrriage
Anisotropy	0.035	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 51.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.014 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.019 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6367	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, LI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/2959	0.62	3/4005 (0.1%)
1	B	0.46	0/2931	0.57	0/3966
All	All	0.46	0/5890	0.60	3/7971 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	93	MET	C-N-CA	-10.00	96.70	121.70
1	A	207	ILE	CG1-CB-CG2	-7.52	94.86	111.40
1	A	314	LEU	CB-CG-CD1	-5.06	102.40	111.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	335	LYS	Peptide
1	A	338	ASN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2905	0	2936	73	0
1	B	2879	0	2907	23	0
2	A	5	0	0	0	0
2	B	5	0	0	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	252	0	0	11	0
4	B	319	0	0	7	0
All	All	6367	0	5843	96	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (96) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:207:ILE:HG22	1:A:286:VAL:HG23	1.49	0.93
1:B:11:ARG:HD2	1:B:269:SER:OG	1.70	0.91
1:A:286:VAL:HG12	1:A:304:VAL:HB	1.53	0.90
1:A:227:LEU:HD23	1:A:333:LEU:HD21	1.58	0.85
1:A:354:GLY:O	4:A:502:HOH:O	1.99	0.81
1:A:90:LEU:O	4:A:501:HOH:O	1.99	0.80
1:A:207:ILE:HG22	1:A:286:VAL:CG2	2.12	0.80
1:A:207:ILE:HD12	1:A:237:PHE:CD2	2.17	0.79
1:B:11:ARG:HG3	4:B:715:HOH:O	1.85	0.77
1:B:210:HIS:HD2	1:B:241:ASN:H	1.34	0.75
1:A:273:GLN:NE2	4:A:504:HOH:O	2.20	0.75
1:A:237:PHE:HE2	4:A:505:HOH:O	1.72	0.73
1:A:306:ILE:HD11	1:A:324:LEU:HA	1.72	0.70
1:A:211:PRO:HD3	1:A:240:PRO:HB2	1.74	0.70
1:A:190:SER:OG	4:A:503:HOH:O	2.09	0.69
1:B:11:ARG:NH2	1:B:12:ALA:HB2	2.07	0.69
1:B:316:CYS:SG	4:B:739:HOH:O	2.50	0.69
1:A:230:ILE:CD1	1:A:334:GLN:HG2	2.23	0.69
1:A:237:PHE:CD1	1:A:238:SER:N	2.61	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:239:TYR:CD1	1:B:268:LYS:HA	2.29	0.68
1:A:223:LEU:HD12	1:A:224:PHE:CD1	2.28	0.68
1:A:208:THR:HA	1:A:237:PHE:CZ	2.29	0.67
1:A:93:MET:O	4:A:501:HOH:O	2.12	0.67
1:A:230:ILE:HG13	1:A:333:LEU:HD22	1.78	0.66
1:A:94:GLN:NE2	4:A:501:HOH:O	2.28	0.66
1:B:272:ILE:N	4:B:504:HOH:O	2.28	0.65
1:A:206:LEU:HD23	1:A:279:LEU:HD23	1.80	0.64
1:A:330:VAL:HA	1:A:333:LEU:HD13	1.80	0.64
1:B:210:HIS:CD2	1:B:241:ASN:H	2.16	0.63
2:B:401:SO4:O4	4:B:501:HOH:O	2.13	0.62
1:A:223:LEU:HD12	1:A:224:PHE:CE1	2.35	0.62
1:A:50:LYS:O	1:A:54:LYS:HE2	2.00	0.61
1:A:257:LYS:HD2	1:A:266:LEU:HD22	1.82	0.61
1:B:88:ASP:HB3	1:B:92:ARG:HH12	1.67	0.59
1:B:269:SER:N	4:B:508:HOH:O	2.35	0.59
1:A:207:ILE:CD1	1:A:237:PHE:CD2	2.86	0.58
1:A:287:GLY:C	1:A:306:ILE:HG22	2.25	0.57
1:A:235:PHE:HB2	1:A:264:VAL:HG22	1.87	0.56
1:A:305:ASN:ND2	4:A:510:HOH:O	2.39	0.56
1:A:279:LEU:HD12	1:A:299:LEU:HD12	1.89	0.55
1:A:329:ILE:HG22	1:A:333:LEU:HD11	1.87	0.55
1:A:268:LYS:HD3	1:A:268:LYS:H	1.73	0.54
1:A:208:THR:CG2	1:A:289:SER:HB3	2.38	0.54
1:A:238:SER:HA	1:A:267:VAL:O	2.09	0.53
1:B:229:GLN:NE2	1:B:326:GLU:OE2	2.42	0.53
1:A:230:ILE:HG13	1:A:333:LEU:CD2	2.38	0.53
1:A:208:THR:HA	1:A:237:PHE:CE2	2.44	0.52
1:A:237:PHE:HD1	1:A:238:SER:H	1.56	0.52
1:B:9:GLY:N	1:B:13:GLU:OE1	2.29	0.51
1:A:259:GLN:NE2	4:A:514:HOH:O	2.44	0.51
1:A:211:PRO:HB2	1:A:243:ASP:HB2	1.91	0.51
1:B:11:ARG:HG3	1:B:12:ALA:N	2.25	0.51
1:A:329:ILE:HG22	1:A:333:LEU:CD1	2.41	0.50
1:A:93:MET:O	1:A:95:PRO:HD3	2.11	0.50
1:A:315:ARG:NH2	4:A:510:HOH:O	2.45	0.49
1:B:283:LEU:HD13	1:B:337:ILE:HG12	1.94	0.49
1:B:270:PHE:HB2	1:B:275:TYR:HB2	1.94	0.49
1:A:224:PHE:CE1	1:A:252:ALA:HB1	2.48	0.49
1:A:267:VAL:HA	1:A:268:LYS:HZ3	1.78	0.48
1:A:322:VAL:HG21	1:A:329:ILE:HG12	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:156:TYR:OH	1:A:314:LEU:HG	2.14	0.47
1:A:220:VAL:O	1:A:223:LEU:HB3	2.14	0.47
1:A:267:VAL:HG13	1:A:268:LYS:HE2	1.95	0.47
1:A:325:ASP:O	1:A:329:ILE:HG13	2.14	0.47
1:A:208:THR:HG23	1:A:237:PHE:CZ	2.50	0.47
1:A:260:LEU:HD13	1:A:263:ARG:HD3	1.96	0.47
1:A:129:ILE:O	1:A:130:THR:HG23	2.15	0.47
1:A:229:GLN:NE2	1:A:326:GLU:OE1	2.47	0.47
1:A:320:LEU:HD23	1:A:320:LEU:HA	1.78	0.46
1:A:207:ILE:CD1	1:A:237:PHE:HD2	2.26	0.46
1:B:154:GLU:OE2	4:B:502:HOH:O	2.21	0.46
1:A:237:PHE:CE2	4:A:505:HOH:O	2.56	0.46
1:B:272:ILE:HA	1:B:275:TYR:HB3	1.97	0.45
1:A:155:GLU:H	1:A:155:GLU:CD	2.20	0.45
1:A:270:PHE:CE1	1:A:278:VAL:HG21	2.52	0.45
1:B:296:ALA:HB3	1:B:297:PRO:HD3	1.99	0.44
1:A:132:GLY:HA3	1:A:135:ASP:OD2	2.19	0.43
1:A:293:LEU:HD22	1:A:313:ARG:HB2	2.01	0.43
1:B:154:GLU:HG2	1:B:172:ASN:ND2	2.34	0.43
1:B:272:ILE:HG12	1:B:276:LEU:HG	2.01	0.43
1:A:247:THR:HA	1:A:250:VAL:HG12	2.00	0.42
1:A:27:ASP:OD1	1:A:29:GLU:HG2	2.20	0.42
1:A:250:VAL:HG13	1:A:251:LYS:HG3	2.01	0.42
1:B:11:ARG:CD	1:B:269:SER:OG	2.55	0.42
1:B:11:ARG:CG	4:B:715:HOH:O	2.56	0.42
1:A:230:ILE:HD11	1:A:330:VAL:HG13	2.01	0.42
1:A:91:LYS:HE2	1:A:91:LYS:HB3	1.88	0.42
1:A:152:ALA:HB1	1:A:351:LEU:HD22	2.01	0.41
1:A:226:ALA:O	1:A:330:VAL:HG22	2.21	0.41
1:A:166:GLN:HG3	1:A:373:ALA:HB1	2.02	0.41
1:A:247:THR:O	1:A:250:VAL:HG12	2.21	0.41
1:A:268:LYS:HE2	1:A:268:LYS:HB2	1.90	0.41
1:A:124:LEU:O	1:A:125:HIS:HB2	2.20	0.40
1:A:207:ILE:HA	1:A:286:VAL:O	2.21	0.40
1:A:136:GLU:OE2	1:B:375:PHE:HB2	2.22	0.40
1:A:207:ILE:O	1:A:237:PHE:CG	2.74	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	366/380 (96%)	341 (93%)	22 (6%)	3 (1%)	19	13
1	B	362/380 (95%)	351 (97%)	11 (3%)	0	100	100
All	All	728/760 (96%)	692 (95%)	33 (4%)	3 (0%)	34	30

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	268	LYS
1	A	212	GLU
1	A	211	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	318/328 (97%)	308 (97%)	10 (3%)	40	40
1	B	315/328 (96%)	312 (99%)	3 (1%)	76	81
All	All	633/656 (96%)	620 (98%)	13 (2%)	53	57

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	ARG
1	A	13	GLU
1	A	54	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	95	PRO
1	A	134	TYR
1	A	190	SER
1	A	228	LYS
1	A	339	PHE
1	A	349	PRO
1	A	362	GLU
1	B	11	ARG
1	B	134	TYR
1	B	339	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	GLN
1	A	234	ASN
1	A	300	GLN
1	B	210	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	401	-	4,4,4	0.15	0	6,6,6	0.22	0
2	SO4	B	401	-	4,4,4	0.22	0	6,6,6	0.23	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	370/380 (97%)	0.99	85 (22%) 0 0	18, 44, 119, 149	0
1	B	366/380 (96%)	-0.16	9 (2%) 57 56	19, 35, 59, 85	0
All	All	736/760 (96%)	0.42	94 (12%) 3 3	18, 38, 111, 149	0

All (94) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	215	LEU	11.1
1	A	214	ASN	8.8
1	A	216	LEU	8.8
1	A	217	GLU	8.4
1	A	339	PHE	8.0
1	A	249	ILE	7.8
1	A	213	THR	7.8
1	A	243	ASP	7.3
1	A	251	LYS	6.1
1	A	242	ALA	5.9
1	A	256	LEU	5.7
1	A	255	ASP	5.6
1	A	316	CYS	5.4
1	A	334	GLN	5.3
1	A	241	ASN	5.3
1	A	212	GLU	5.2
1	A	219	ASN	5.2
1	A	244	ASN	5.2
1	A	379	LEU	5.0
1	A	239	TYR	5.0
1	A	317	GLU	4.8
1	B	353	LEU	4.8
1	A	220	VAL	4.6
1	A	225	ASP	4.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	223	LEU	4.3
1	A	246	ASN	4.3
1	A	338	ASN	4.3
1	A	133	ALA	4.3
1	A	261	PRO	4.3
1	A	218	GLU	4.2
1	B	339	PHE	4.2
1	A	208	THR	4.2
1	A	245	GLY	4.1
1	A	230	ILE	4.1
1	A	323	ARG	4.1
1	A	320	LEU	3.9
1	A	233	VAL	3.9
1	A	268	LYS	3.8
1	A	131	GLU	3.8
1	A	237	PHE	3.7
1	A	253	MET	3.5
1	A	258	ALA	3.5
1	A	207	ILE	3.5
1	A	265	LEU	3.5
1	A	235	PHE	3.5
1	A	229	GLN	3.4
1	A	240	PRO	3.4
1	A	264	VAL	3.4
1	B	379	LEU	3.4
1	A	250	VAL	3.4
1	A	259	GLN	3.3
1	A	1	MET	3.3
1	A	260	LEU	3.3
1	A	221	ALA	3.3
1	A	337	ILE	3.2
1	A	196	TYR	3.2
1	A	209	TYR	3.2
1	A	254	LEU	3.2
1	A	232	ASP	3.0
1	A	248	ASN	3.0
1	A	302	PRO	3.0
1	B	338	ASN	3.0
1	A	226	ALA	2.9
1	A	247	THR	2.9
1	A	322	VAL	2.9
1	A	329	ILE	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	335	LYS	2.8
1	A	206	LEU	2.8
1	A	350	PRO	2.7
1	A	266	LEU	2.7
1	A	262	ASP	2.7
1	A	257	LYS	2.6
1	A	304	VAL	2.6
1	A	197	ASP	2.5
1	A	238	SER	2.5
1	A	353	LEU	2.5
1	A	328	GLU	2.4
1	A	224	PHE	2.4
1	A	195	LEU	2.4
1	B	232	ASP	2.4
1	A	211	PRO	2.4
1	B	11	ARG	2.3
1	A	324	LEU	2.3
1	A	205	PHE	2.2
1	A	308	ASP	2.2
1	B	269	SER	2.2
1	A	227	LEU	2.1
1	A	318	SER	2.1
1	B	216	LEU	2.1
1	A	333	LEU	2.1
1	A	321	ASP	2.0
1	A	132	GLY	2.0
1	B	101	LEU	2.0
1	A	286	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	LI	A	402	1/1	0.80	0.15	22,22,22,22	0
2	SO4	A	401	5/5	0.94	0.09	74,77,79,80	0
2	SO4	B	401	5/5	0.99	0.09	32,41,44,51	0
3	LI	B	402	1/1	0.99	0.25	40,40,40,40	0

6.5 Other polymers [i](#)

There are no such residues in this entry.