



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 10, 2020 – 01:11 PM BST

PDB ID : 3ZM8  
Title : Crystal structure of Podospora anserina GH26-CBM35 beta-(1,4)- mannanase  
Authors : Couturier, M.; Roussel, A.; Rosengren, A.; Leone, P.; Stalbrand, H.; Berrin, J.G.  
Deposited on : 2013-02-06  
Resolution : 2.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3168 (2.90-2.82)
Clashscore	141614	3438 (2.90-2.82)
Ramachandran outliers	138981	3348 (2.90-2.82)
Sidechain outliers	138945	3351 (2.90-2.82)
RSRZ outliers	127900	3103 (2.90-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	475	
2	B	2	

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3676 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called GH26 ENDO-BETA-1,4-MANNANASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	444	3493	2233	590	667	3	0	0	0

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MET	-	expression tag	UNP E2GHW2
A	-1	SER	-	expression tag	UNP E2GHW2
A	0	ILE	-	expression tag	UNP E2GHW2
A	450	GLY	-	expression tag	UNP E2GHW2
A	451	ASP	-	expression tag	UNP E2GHW2
A	452	GLU	-	expression tag	UNP E2GHW2
A	453	GLN	-	expression tag	UNP E2GHW2
A	454	LYS	-	expression tag	UNP E2GHW2
A	455	LEU	-	expression tag	UNP E2GHW2
A	456	ILE	-	expression tag	UNP E2GHW2
A	457	SER	-	expression tag	UNP E2GHW2
A	458	GLU	-	expression tag	UNP E2GHW2
A	459	GLU	-	expression tag	UNP E2GHW2
A	460	ASP	-	expression tag	UNP E2GHW2
A	461	LEU	-	expression tag	UNP E2GHW2
A	462	ASN	-	expression tag	UNP E2GHW2
A	463	SER	-	expression tag	UNP E2GHW2
A	464	ALA	-	expression tag	UNP E2GHW2
A	465	VAL	-	expression tag	UNP E2GHW2
A	466	ASP	-	expression tag	UNP E2GHW2
A	467	HIS	-	expression tag	UNP E2GHW2
A	468	HIS	-	expression tag	UNP E2GHW2
A	469	HIS	-	expression tag	UNP E2GHW2
A	470	HIS	-	expression tag	UNP E2GHW2
A	471	HIS	-	expression tag	UNP E2GHW2
A	472	HIS	-	expression tag	UNP E2GHW2

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace	
			Total	C	N				O
2	B	2	28	16	2	10	0	0	0


- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
3	A	1	1	1	0	0

- Molecule 4 is MERCURY (II) ION (three-letter code: HG) (formula: Hg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Hg		
4	A	1	1	1	0	0

- Molecule 5 is L(+)-TARTARIC ACID (three-letter code: TLA) (formula: C<sub>4</sub>H<sub>6</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	A	1	10	4	6	0	0


- Molecule 6 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
6	A	143	Total 143	O 143	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: GH26 ENDO-BETA-1,4-MANNANASE



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	97.49Å 97.49Å 268.72Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.30 – 2.85 47.96 – 2.70	Depositor EDS
% Data completeness (in resolution range)	100.0 (28.30-2.85) 99.7 (47.96-2.70)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.86 (at 2.69Å)	Xtrriage
Refinement program	BUSTER 2.11.4	Depositor
R, $R_{free}$	0.206 , 0.257 0.203 , 0.256	Depositor DCC
$R_{free}$ test set	1110 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	68.2	Xtrriage
Anisotropy	0.148	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 57.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3676	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	68.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.97% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HG, CA, TLA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.43	0/3594	0.65	0/4930

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3493	0	3327	31	0
2	B	28	0	25	0	0
3	A	1	0	0	0	0
4	A	1	0	0	0	0
5	A	10	0	4	0	0
6	A	143	0	0	0	0
All	All	3676	0	3356	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:90:ALA:CB	1:A:91:GLY:HA2	2.12	0.78
1:A:90:ALA:HB3	1:A:91:GLY:HA2	1.79	0.64
1:A:161:VAL:HG21	1:A:439:LEU:HD11	1.80	0.62
1:A:90:ALA:HB1	1:A:91:GLY:HA2	1.81	0.62
1:A:90:ALA:CB	1:A:91:GLY:CA	2.80	0.59
1:A:178:TRP:O	1:A:182:GLN:HG2	2.05	0.57
1:A:78:LEU:HD22	1:A:102:VAL:HG22	1.87	0.56
1:A:56:THR:HG22	1:A:106:ALA:H	1.70	0.56
1:A:232:ALA:HB3	1:A:253:THR:HG21	1.89	0.55
1:A:161:VAL:HG21	1:A:439:LEU:CD1	2.38	0.52
1:A:90:ALA:HB3	1:A:91:GLY:CA	2.38	0.52
1:A:71:ASP:HB3	1:A:89:PRO:O	2.10	0.51
1:A:66:ALA:HB3	1:A:121:LEU:HB2	1.97	0.47
1:A:373:TYR:OH	1:A:406:ASN:HB2	2.15	0.47
1:A:167:ILE:HG23	1:A:188:ALA:HB3	1.97	0.46
1:A:74:THR:O	1:A:85:GLU:HA	2.16	0.46
1:A:15:GLU:HG3	1:A:34:ARG:HB3	2.00	0.44
1:A:196:ASP:HB2	1:A:208:SER:HB2	1.99	0.44
1:A:25:ASP:HB3	1:A:36:TYR:CE1	2.54	0.43
1:A:284:LYS:HE3	1:A:331:HIS:HB3	2.01	0.43
1:A:191:ALA:HA	1:A:225:SER:O	2.19	0.42
1:A:376:LEU:HB3	1:A:386:ILE:HD12	2.01	0.42
1:A:43:GLY:HA2	1:A:115:ASN:HD22	1.84	0.42
1:A:94:PHE:CE1	1:A:123:ASP:HB2	2.54	0.42
1:A:236:LEU:HD23	1:A:244:TRP:HB2	2.01	0.42
1:A:46:LYS:HB3	1:A:114:VAL:HG13	2.02	0.41
1:A:218:HIS:CG	1:A:291:VAL:HG22	2.55	0.41
1:A:340:TRP:CE2	1:A:349:PRO:HD3	2.55	0.41
1:A:197:TYR:HE2	1:A:210:ALA:HB3	1.85	0.40
1:A:103:LEU:HD13	1:A:403:TYR:CE2	2.56	0.40
1:A:243:LYS:HD3	1:A:245:TRP:CZ2	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	442/475 (93%)	419 (95%)	22 (5%)	1 (0%)	47 75

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	414	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	362/391 (93%)	351 (97%)	11 (3%)	41 72

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	41	ASP
1	A	56	THR
1	A	84	SER
1	A	102	VAL
1	A	120	TYR
1	A	227	LEU
1	A	234	VAL
1	A	268	ASN
1	A	381	ARG
1	A	419	ASN
1	A	442	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	B	1	1,2	14,14,15	0.35	0	17,19,21	1.31	3 (17%)
2	NAG	B	2	2	14,14,15	0.30	0	17,19,21	0.71	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	NAG	O5-C1-C2	-4.06	104.88	111.29
2	B	1	NAG	C1-C2-N2	2.58	114.90	110.49
2	B	2	NAG	C1-O5-C5	2.09	115.02	112.19
2	B	1	NAG	C1-O5-C5	2.00	114.91	112.19


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	TLA	A	1452	-	3,9,9	0.69	0	6,12,12	1.43	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	TLA	A	1452	-	-	1/4/12/12	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
5	A	1452	TLA	C4-C3-C2	2.74	119.00	113.11

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1452	TLA	C1-C2-C3-C4

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	444/475 (93%)	0.43	32 (7%) <b>15</b> <b>11</b>	40, 67, 98, 118	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	119	TRP	4.5
1	A	68	ILE	3.8
1	A	93	SER	3.6
1	A	69	TYR	3.5
1	A	92	ASP	3.5
1	A	40	PHE	3.2
1	A	122	ILE	3.2
1	A	234	VAL	3.1
1	A	21	GLY	3.1
1	A	37	VAL	3.1
1	A	121	LEU	3.0
1	A	120	TYR	3.0
1	A	36	TYR	3.0
1	A	26	THR	2.9
1	A	262	ALA	2.7
1	A	322	ILE	2.6
1	A	261	LEU	2.6
1	A	24	VAL	2.5
1	A	34	ARG	2.5
1	A	47	ILE	2.5
1	A	19	LEU	2.5
1	A	45	ASP	2.5
1	A	25	ASP	2.5
1	A	18	ILE	2.5
1	A	265	GLN	2.3
1	A	44	SER	2.3
1	A	269	TYR	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	70	GLY	2.2
1	A	42	GLU	2.2
1	A	271	LEU	2.1
1	A	30	GLY	2.1
1	A	244	TRP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	B	2	14/15	0.88	0.21	82,86,91,92	0
2	NAG	B	1	14/15	0.94	0.23	85,89,94,94	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	TLA	A	1452	10/10	0.78	0.30	82,84,87,88	0
4	HG	A	1451	1/1	0.85	0.18	93,93,93,93	1
3	CA	A	1450	1/1	0.96	0.07	90,90,90,90	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.