



Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 06:42 pm BST

PDB ID : 5ZMC
Title : Structural Basis for Reactivation of -146C>T Mutant TERT Promoter by cooperative binding of p52 and ETS1/2
Authors : Xu, X.; Bharath, S.R.; Song, H.
Deposited on : 2018-04-02
Resolution : 2.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	16	
2	D	16	
3	B	111	
4	A	295	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2393 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(P*CP*GP*GP*GP*GP*AP*CP*CP*CP*GP*GP*AP*AP*GP*GP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	C	16	337	156	72	93	16	0	0	0

- Molecule 2 is a DNA chain called DNA (5'-D(P*GP*CP*CP*CP*TP*TP*CP*CP*GP*GP*GP*TP*CP*CP*CP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	D	16	319	151	53	99	16	0	0	0

- Molecule 3 is a protein called Protein C-ets-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	B	106	890	576	153	157	4	0	0	0

- Molecule 4 is a protein called Nuclear factor NF-kappa-B p100 subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	A	103	847	539	146	159	3	0	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	71.41Å 71.41Å 262.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	71.41 – 2.99 48.32 – 2.99	Depositor EDS
% Data completeness (in resolution range)	89.4 (71.41-2.99) 89.4 (48.32-2.99)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.00 (at 3.01Å)	Xtrriage
Refinement program	REFMAC 5.8.0216	Depositor
R, R_{free}	0.259 , 0.288 0.265 , 0.285	Depositor DCC
R_{free} test set	664 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	78.3	Xtrriage
Anisotropy	0.444	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 81.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	2393	wwPDB-VP
Average B, all atoms (Å ²)	101.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	0.56	0/380	0.84	0/586
2	D	0.41	0/354	0.79	0/542
3	B	0.47	0/915	0.65	0/1233
4	A	0.43	0/867	0.67	0/1167
All	All	0.46	0/2516	0.71	0/3528

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	337	0	177	15	0
2	D	319	0	180	6	0
3	B	890	0	885	13	0
4	A	847	0	828	6	0
All	All	2393	0	2070	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:-3:DG:H2''	1:C:-2:DG:C8	2.15	0.81
1:C:0:DA:H2'	1:C:1:DC:O4'	1.84	0.77
1:C:-3:DG:C2'	1:C:-2:DG:C8	2.73	0.72
1:C:-4:DG:H2''	1:C:-3:DG:O5'	1.91	0.71
2:D:110:DC:H2'	2:D:111:DT:H72	1.79	0.65
1:C:-5:DC:H2'	1:C:-4:DG:C8	2.31	0.65
4:A:263:TYR:CZ	4:A:306:PHE:HB2	2.34	0.62
3:B:361:TRP:CG	3:B:418:LEU:HD12	2.39	0.57
4:A:228:LEU:HD12	4:A:317:VAL:C	2.25	0.56
3:B:404:LYS:NZ	3:B:405:THR:O	2.39	0.56
3:B:387:GLU:O	3:B:390:SER:OG	2.25	0.55
3:B:403:HIS:CE1	3:B:415:VAL:HG11	2.42	0.54
2:D:110:DC:H2'	2:D:111:DT:C7	2.37	0.54
3:B:361:TRP:CD2	3:B:418:LEU:HD12	2.42	0.54
1:C:2:DC:H2'	1:C:3:DC:H6	1.73	0.54
3:B:424:TYR:CZ	4:A:298:LYS:HA	2.44	0.53
3:B:370:GLU:OE2	3:B:373:ARG:NH2	2.38	0.53
3:B:345:LEU:HD12	3:B:356:TRP:CE2	2.45	0.52
1:C:-3:DG:H2'	1:C:-2:DG:C8	2.43	0.51
2:D:119:DC:H2'	2:D:120:DC:C6	2.48	0.48
1:C:2:DC:H2'	1:C:3:DC:C6	2.48	0.48
4:A:263:TYR:CZ	4:A:306:PHE:CB	2.97	0.48
1:C:0:DA:H4'	1:C:1:DC:OP1	2.14	0.48
1:C:4:DG:H2''	1:C:5:DG:C8	2.50	0.47
2:D:109:DC:H2''	2:D:110:DC:H5''	1.96	0.46
3:B:384:MET:CE	3:B:389:LEU:N	2.78	0.46
2:D:119:DC:H2''	2:D:120:DC:H5'	1.99	0.45
4:A:259:GLU:OE2	4:A:312:LYS:HE2	2.17	0.44
3:B:389:LEU:HD23	3:B:389:LEU:C	2.37	0.44
1:C:3:DC:H2'	1:C:4:DG:C8	2.52	0.44
4:A:230:ILE:HG13	4:A:309:LEU:HD13	1.99	0.44
1:C:0:DA:C2'	1:C:1:DC:O4'	2.63	0.43
1:C:-2:DG:C2'	1:C:-1:DG:C8	3.01	0.43
2:D:113:DC:H2'	2:D:114:DC:C6	2.53	0.43
3:B:421:LEU:HD12	3:B:421:LEU:HA	1.87	0.43
1:C:-2:DG:H2''	1:C:-1:DG:C8	2.54	0.43
3:B:361:TRP:HB3	3:B:414:PHE:HB2	2.02	0.41
3:B:394:ARG:HG2	3:B:397:TYR:CE2	2.55	0.41
1:C:-4:DG:OP2	1:C:-4:DG:H8	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	B	104/111 (94%)	95 (91%)	9 (9%)	0	100	100
4	A	101/295 (34%)	94 (93%)	7 (7%)	0	100	100
All	All	205/406 (50%)	189 (92%)	16 (8%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	B	96/99 (97%)	95 (99%)	1 (1%)	76	91
4	A	93/256 (36%)	93 (100%)	0	100	100
All	All	189/355 (53%)	188 (100%)	1 (0%)	88	96

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	B	332	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	C	16/16 (100%)	0.07	0	100	87, 112, 135, 149	0
2	D	16/16 (100%)	0.26	0	100	73, 106, 131, 136	0
3	B	106/111 (95%)	0.65	6 (5%)	23	71, 86, 112, 139	0
4	A	103/295 (34%)	1.16	19 (18%)	1	84, 107, 143, 160	0
All	All	241/438 (55%)	0.80	25 (10%)	6	71, 96, 138, 160	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	A	305	VAL	3.9
4	A	311	ARG	3.9
4	A	267	GLU	3.8
4	A	313	ARG	3.6
4	A	228	LEU	3.4
4	A	319	ASP	3.4
4	A	226	SER	3.1
4	A	322	GLN	2.7
4	A	265	ASP	2.7
4	A	320	SER	2.6
4	A	308	GLN	2.6
4	A	309	LEU	2.5
4	A	314	GLY	2.5
4	A	263	TYR	2.5
4	A	254	GLN	2.4
4	A	312	LYS	2.4
3	B	368	PRO	2.4
3	B	429	LEU	2.4
4	A	307	LEU	2.3
4	A	252	LYS	2.3
3	B	406	ALA	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	B	437	PRO	2.2
3	B	409	ARG	2.2
3	B	372	ALA	2.1
4	A	306	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.