



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 2, 2024 – 01:25 PM JST

PDB ID : 8ZM8
Title : Crystal Structure of the first bromodomain of human BRD4 BD1 in complex with the inhibitor Y13221
Authors : Li, J.; Hu, Q.; Xu, H.; Zhao, X.; Zhang, C.; Zhu, R.; Wu, X.; Zhang, Y.; Xu, Y.
Deposited on : 2024-05-22
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.21
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.004 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2511 (3.00-3.00)
Clashscore	180529	2866 (3.00-3.00)
Ramachandran outliers	177936	2778 (3.00-3.00)
Sidechain outliers	177891	2781 (3.00-3.00)
RSRZ outliers	164620	2523 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	141	 2% 67% 8% 23%
1	B	141	 % 68% 9% 23%
1	C	141	 % 60% 15% 23%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2857 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BRD4_HUMAN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	107	900	590	144	160	6	0	0	0
1	B	109	920	601	150	163	6	0	0	0
1	C	108	912	597	147	162	6	0	0	0

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	28	MET	-	expression tag	UNP O60885
A	29	LYS	-	expression tag	UNP O60885
A	30	LYS	-	expression tag	UNP O60885
A	31	GLY	-	expression tag	UNP O60885
A	32	HIS	-	expression tag	UNP O60885
A	33	HIS	-	expression tag	UNP O60885
A	34	HIS	-	expression tag	UNP O60885
A	35	HIS	-	expression tag	UNP O60885
A	36	HIS	-	expression tag	UNP O60885
A	37	HIS	-	expression tag	UNP O60885
A	38	LEU	-	expression tag	UNP O60885
A	39	VAL	-	expression tag	UNP O60885
A	40	PRO	-	expression tag	UNP O60885
A	41	ARG	-	expression tag	UNP O60885
A	42	GLY	-	expression tag	UNP O60885
A	43	SER	-	expression tag	UNP O60885
B	28	MET	-	expression tag	UNP O60885
B	29	LYS	-	expression tag	UNP O60885
B	30	LYS	-	expression tag	UNP O60885
B	31	GLY	-	expression tag	UNP O60885
B	32	HIS	-	expression tag	UNP O60885
B	33	HIS	-	expression tag	UNP O60885
B	34	HIS	-	expression tag	UNP O60885

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	35	HIS	-	expression tag	UNP O60885
B	36	HIS	-	expression tag	UNP O60885
B	37	HIS	-	expression tag	UNP O60885
B	38	LEU	-	expression tag	UNP O60885
B	39	VAL	-	expression tag	UNP O60885
B	40	PRO	-	expression tag	UNP O60885
B	41	ARG	-	expression tag	UNP O60885
B	42	GLY	-	expression tag	UNP O60885
B	43	SER	-	expression tag	UNP O60885
C	28	MET	-	expression tag	UNP O60885
C	29	LYS	-	expression tag	UNP O60885
C	30	LYS	-	expression tag	UNP O60885
C	31	GLY	-	expression tag	UNP O60885
C	32	HIS	-	expression tag	UNP O60885
C	33	HIS	-	expression tag	UNP O60885
C	34	HIS	-	expression tag	UNP O60885
C	35	HIS	-	expression tag	UNP O60885
C	36	HIS	-	expression tag	UNP O60885
C	37	HIS	-	expression tag	UNP O60885
C	38	LEU	-	expression tag	UNP O60885
C	39	VAL	-	expression tag	UNP O60885
C	40	PRO	-	expression tag	UNP O60885
C	41	ARG	-	expression tag	UNP O60885
C	42	GLY	-	expression tag	UNP O60885
C	43	SER	-	expression tag	UNP O60885

- Molecule 2 is 2-[2-ethyl-4-(methoxymethyl)-1 {H}-imidazol-5-yl]-7-[2-(4-fluoranyl-2,6-dimethyl-phenoxy)-5-(2-oxidanylpropan-2-yl)phenyl]-5-methyl-furo[3,2-c]pyridin-4-one (three-letter code: A1L1X) (formula: C₃₂H₃₄FN₃O₅) (labeled as "Ligand of Interest" by depositor).


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: BRD4_HUMAN



- Molecule 1: BRD4_HUMAN



- Molecule 1: BRD4_HUMAN



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	142.07Å 142.07Å 134.14Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	123.03 – 3.00 123.03 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (123.03-3.00) 99.9 (123.03-3.00)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.32 (at 3.01Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.207 , 0.265 0.213 , 0.261	Depositor DCC
R_{free} test set	864 reflections (5.22%)	wwPDB-VP
Wilson B-factor (Å ²)	85.5	Xtrriage
Anisotropy	0.703	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 91.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2857	wwPDB-VP
Average B, all atoms (Å ²)	99.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: A1L1X

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.62	0/924	0.73	0/1256
1	B	0.62	0/944	0.71	0/1282
1	C	0.65	0/936	0.74	0/1271
All	All	0.63	0/2804	0.73	0/3809

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	900	0	896	4	0
1	B	920	0	917	4	0
1	C	912	0	913	11	0
2	A	41	0	0	0	0
2	B	41	0	0	1	0
2	C	41	0	0	0	0
3	C	2	0	0	0	0
All	All	2857	0	2726	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:94:LEU:HG	1:C:139:TYR:OH	2.05	0.56
1:A:71:LEU:HD21	1:A:111:LYS:HD3	1.89	0.53
1:C:61:ASN:OD1	1:C:62:GLN:N	2.43	0.51
1:C:157:PHE:CE1	1:C:161:ILE:HD13	2.46	0.51
1:B:81:TRP:CG	1:B:82:PRO:HD3	2.47	0.49
1:A:69:VAL:O	1:A:73:THR:OG1	2.31	0.48
1:C:81:TRP:CG	1:C:82:PRO:HD3	2.48	0.48
1:C:161:ILE:O	1:C:164:LEU:HB2	2.13	0.47
1:C:115:GLU:O	1:C:116:ASN:ND2	2.51	0.44
1:C:164:LEU:HD22	1:C:164:LEU:HA	1.89	0.43
1:C:61:ASN:HB3	1:C:64:GLN:HB2	1.99	0.43
1:A:81:TRP:N	1:A:82:PRO:CD	2.81	0.43
1:C:113:ARG:NH1	1:C:128:ASP:OD2	2.52	0.43
1:C:146:ILE:O	1:C:149:MET:HB2	2.19	0.42
1:B:146:ILE:HD13	2:B:201:A1LIX:CAC	2.50	0.42
1:A:77:HIS:HE1	1:A:79:PHE:CD2	2.38	0.41
1:C:103:THR:N	1:C:104:PRO:HD3	2.36	0.41
1:B:79:PHE:O	1:B:82:PRO:HD2	2.21	0.41
1:B:93:ASN:HD22	1:B:93:ASN:HA	1.81	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	105/141 (74%)	98 (93%)	7 (7%)	0	100	100
1	B	107/141 (76%)	103 (96%)	4 (4%)	0	100	100
1	C	106/141 (75%)	97 (92%)	9 (8%)	0	100	100
All	All	318/423 (75%)	298 (94%)	20 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	99/132 (75%)	94 (95%)	5 (5%)	20	53
1	B	101/132 (76%)	93 (92%)	8 (8%)	10	35
1	C	101/132 (76%)	92 (91%)	9 (9%)	8	31
All	All	301/396 (76%)	279 (93%)	22 (7%)	11	39

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	78	GLN
1	A	90	VAL
1	A	103	THR
1	A	105	MET
1	A	111	LYS
1	B	78	GLN
1	B	92	LEU
1	B	94	LEU
1	B	105	MET
1	B	121	ASN
1	B	131	THR
1	B	145	ASP
1	B	166	THR
1	C	59	GLN
1	C	94	LEU
1	C	96	ASP
1	C	99	LYS
1	C	102	LYS
1	C	103	THR
1	C	112	LYS
1	C	127	GLN
1	C	164	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	116	ASN
1	A	127	GLN
1	B	78	GLN
1	B	93	ASN
1	B	116	ASN
1	B	121	ASN
1	B	162	ASN
1	C	59	GLN
1	C	116	ASN
1	C	159	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	A1L1X	A	201	-	35,45,45	4.04	11 (31%)	48,68,68	1.13	1 (2%)
2	A1L1X	C	201	-	35,45,45	4.19	8 (22%)	48,68,68	1.24	5 (10%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	A1L1X	B	201	-	35,45,45	4.14	10 (28%)	48,68,68	1.09	4 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1L1X	A	201	-	-	3/17/23/23	0/5/5/5
2	A1L1X	C	201	-	-	3/17/23/23	0/5/5/5
2	A1L1X	B	201	-	-	5/17/23/23	0/5/5/5

All (29) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	201	A1L1X	CBB-CAN	-19.60	1.40	1.53
2	B	201	A1L1X	CBB-CAN	-19.27	1.40	1.53
2	A	201	A1L1X	CBB-CAN	-17.34	1.41	1.53
2	A	201	A1L1X	CBF-CAH	-6.65	1.34	1.49
2	A	201	A1L1X	CAB-CAC	-6.42	1.34	1.47
2	B	201	A1L1X	CAB-CAC	-6.32	1.34	1.47
2	C	201	A1L1X	CBF-CAH	-6.26	1.35	1.49
2	A	201	A1L1X	CBM-CBJ	6.23	1.57	1.50
2	B	201	A1L1X	CBF-CAH	-6.06	1.35	1.49
2	C	201	A1L1X	CAB-CAC	-5.82	1.35	1.47
2	A	201	A1L1X	CAA-CAF	-5.81	1.33	1.47
2	A	201	A1L1X	CAZ-CAX	-5.55	1.39	1.51
2	B	201	A1L1X	CAZ-CAX	-5.52	1.40	1.51
2	C	201	A1L1X	CAZ-CAX	-5.49	1.40	1.51
2	C	201	A1L1X	CAA-CAF	-5.46	1.34	1.47
2	C	201	A1L1X	CBM-CBJ	5.19	1.56	1.50
2	B	201	A1L1X	CBA-CAT	-5.17	1.40	1.51
2	B	201	A1L1X	CAA-CAF	-5.10	1.35	1.47
2	C	201	A1L1X	CBA-CAT	-5.01	1.41	1.51
2	A	201	A1L1X	CBA-CAT	-4.56	1.41	1.51
2	B	201	A1L1X	CBM-CBJ	4.52	1.55	1.50
2	C	201	A1L1X	CAJ-CAF	-4.17	1.42	1.49
2	A	201	A1L1X	CAJ-CAF	-4.05	1.42	1.49
2	B	201	A1L1X	CAJ-CAF	-4.00	1.42	1.49
2	B	201	A1L1X	CAI-CAB	-2.88	1.34	1.40
2	A	201	A1L1X	CAI-CAB	-2.76	1.34	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	201	A1L1X	FAY-CAV	-2.30	1.30	1.36
2	A	201	A1L1X	OBN-CBM	2.12	1.50	1.41
2	B	201	A1L1X	OAR-CAQ	-2.05	1.34	1.39

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	201	A1L1X	CAU-CAV-CAW	-2.95	119.80	123.52
2	C	201	A1L1X	CAQ-OAR-CAS	2.64	123.65	117.79
2	C	201	A1L1X	CAF-CAE-NAD	-2.55	119.70	122.96
2	A	201	A1L1X	CBD-CBB-CAN	-2.40	106.62	110.91
2	B	201	A1L1X	CAO-CAN-CBB	-2.22	118.72	121.55
2	B	201	A1L1X	CAF-CAE-NAD	-2.17	120.18	122.96
2	B	201	A1L1X	CAQ-OAR-CAS	2.15	122.57	117.79
2	C	201	A1L1X	CAT-CAU-CAV	2.11	120.78	119.26
2	C	201	A1L1X	CAQ-CAJ-CAF	2.05	124.08	121.57
2	B	201	A1L1X	CAU-CAV-CAW	-2.03	120.95	123.52

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	201	A1L1X	CBJ-CBM-OBN-CBO
2	C	201	A1L1X	CBJ-CBM-OBN-CBO
2	B	201	A1L1X	CAA-CAF-CAJ-CAM
2	B	201	A1L1X	CBJ-CBM-OBN-CBO
2	C	201	A1L1X	CAA-CAF-CAJ-CAM
2	B	201	A1L1X	NBI-CBH-CBK-CBL
2	A	201	A1L1X	CAA-CAF-CAJ-CAM
2	B	201	A1L1X	CAA-CAF-CAJ-CAQ
2	C	201	A1L1X	CAA-CAF-CAJ-CAQ
2	A	201	A1L1X	NBG-CBH-CBK-CBL
2	B	201	A1L1X	NBG-CBH-CBK-CBL


There are no ring outliers.



1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	201	A1L1X	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	107/141 (75%)	-0.29	3 (2%) 55 33	59, 82, 121, 164	0
1	B	109/141 (77%)	-0.29	2 (1%) 67 45	70, 88, 126, 144	0
1	C	108/141 (76%)	0.17	2 (1%) 66 44	76, 116, 159, 174	0
All	All	324/423 (76%)	-0.13	7 (2%) 62 40	59, 94, 146, 174	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	92	LEU	2.8
1	B	166	THR	2.8
1	A	117	ASN	2.3
1	C	72	LYS	2.2
1	A	166	THR	2.2
1	B	92	LEU	2.2
1	A	60	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	A1L1X	C	201	41/41	0.89	0.18	110,130,162,169	0
2	A1L1X	B	201	41/41	0.92	0.15	79,102,132,144	0
2	A1L1X	A	201	41/41	0.92	0.15	67,85,122,127	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around A1L1X B 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.