



Full wwPDB X-ray Structure Validation Report ⓘ

May 28, 2020 – 09:05 pm BST

PDB ID : 1ZNI
Title : INSULIN
Authors : Turkenburg, M.G.W.; Whittingham, J.L.; Dodson, G.G.; Dodson, E.J.; Xiao, B.; Bentley, G.A.
Deposited on : 1997-09-23
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	21	57% (green), 38% (yellow), 5% (orange)
1	C	21	57% (green), 43% (yellow)
2	B	30	63% (green), 30% (yellow), 7% (orange)
2	D	30	77% (green), 20% (yellow), . (orange)

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 929 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called INSULIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	21	163	99	25	35	4	0	0	0
1	C	21	167	102	25	36	4	0	2	0

- Molecule 2 is a protein called INSULIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	30	245	160	42	41	2	0	1	0
2	D	30	245	160	40	43	2	0	1	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Zn	0	0
			2	2		
3	D	1	Total	Zn	0	0
			1	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	3	Total	Cl	0	0
			3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	28	Total	O	0	0
			28	28		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	20	Total 20	O 20	0	0
5	C	23	Total 23	O 23	0	0
5	D	32	Total 32	O 32	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: INSULIN

Chain A: 



- Molecule 1: INSULIN

Chain C: 



- Molecule 2: INSULIN

Chain B: 



- Molecule 2: INSULIN

Chain D: 



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	80.70Å 80.70Å 37.60Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.00 – 1.50	Depositor
% Data completeness (in resolution range)	(Not available) (40.00-1.50)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC	Depositor
R, R_{free}	0.178 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	929	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.16	0/164	1.85	4/220 (1.8%)
1	C	1.23	1/177 (0.6%)	2.07	7/238 (2.9%)
2	B	1.13	0/258	1.93	5/347 (1.4%)
2	D	1.13	0/256	1.87	6/344 (1.7%)
All	All	1.16	1/855 (0.1%)	1.93	22/1149 (1.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	11	CYS	CA-CB	5.50	1.66	1.53

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	25	PHE	CB-CG-CD2	-13.17	111.58	120.80
2	B	25	PHE	CB-CG-CD1	12.87	129.81	120.80
1	C	11	CYS	CA-CB-SG	-9.23	97.38	114.00
1	C	19	TYR	CB-CG-CD1	-8.49	115.91	121.00
1	C	19	TYR	CB-CG-CD2	8.17	125.90	121.00
2	B	24	PHE	CB-CG-CD2	-8.05	115.17	120.80
2	D	26	TYR	CB-CG-CD2	-7.66	116.40	121.00
1	A	14	TYR	CB-CG-CD2	-6.78	116.93	121.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	17	LEU	CB-CG-CD2	6.72	122.43	111.00
1	C	14	TYR	CB-CG-CD1	-6.48	117.11	121.00
1	C	20	CYS	O-C-N	5.93	132.19	122.70
1	A	20	CYS	O-C-N	5.81	132.00	122.70
1	C	16	LEU	CB-CA-C	-5.76	99.26	110.20
1	A	14	TYR	CG-CD2-CE2	-5.72	116.72	121.30
2	B	6	LEU	CB-CG-CD1	-5.59	101.49	111.00
2	B	22	ARG	NE-CZ-NH2	-5.43	117.58	120.30
2	D	1	PHE	CB-CG-CD2	-5.35	117.05	120.80
2	D	26	TYR	CG-CD1-CE1	-5.32	117.05	121.30
2	D	20	GLY	CA-C-O	5.26	130.06	120.60
1	A	19	TYR	CG-CD2-CE2	5.22	125.48	121.30
2	D	8	GLY	CA-C-O	5.08	129.75	120.60
1	C	19	TYR	O-C-N	-5.01	114.68	122.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	20	GLY	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	163	0	149	5	0
1	C	167	0	155	7	0
2	B	245	0	233	5	0
2	D	245	0	232	1	0
3	B	2	0	0	0	0
3	D	1	0	0	0	0
4	B	3	0	0	0	0
5	A	28	0	0	3	0
5	B	20	0	0	0	0
5	C	23	0	0	1	0
5	D	32	0	0	2	0
All	All	929	0	769	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:3[A]:VAL:HG22	5:D:37:HOH:O	1.97	0.64
1:C:4:GLU:HG3	5:C:40:HOH:O	2.10	0.51
1:A:5:GLN:NE2	1:A:19:TYR:OH	2.44	0.51
1:A:21:ASN:ND2	5:A:37:HOH:O	2.45	0.49
1:C:3[A]:VAL:HG23	1:C:4:GLU:CD	2.33	0.49
1:C:12:SER:H	1:C:15:GLN:NE2	2.11	0.48
1:C:12:SER:H	1:C:15:GLN:HE21	1.62	0.47
1:C:3[A]:VAL:HG21	5:D:60:HOH:O	2.15	0.47
1:C:3[A]:VAL:HG23	1:C:4:GLU:OE2	2.15	0.47
1:A:5:GLN:HG3	5:A:49:HOH:O	2.15	0.45
5:A:34:HOH:O	2:B:2:VAL:HG13	2.16	0.45
2:B:29:LYS:O	2:B:30:ALA:C	2.55	0.44
2:B:21:GLU:OE2	2:B:22:ARG:HG3	2.18	0.44
2:B:15:LEU:HD23	2:B:15:LEU:HA	1.82	0.41
1:A:8:THR:HG22	2:B:2:VAL:HG21	2.03	0.40
1:A:13:LEU:O	1:A:17:GLU:HG3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	19/21 (90%)	19 (100%)	0	0	100	100
1	C	21/21 (100%)	19 (90%)	2 (10%)	0	100	100
2	B	29/30 (97%)	27 (93%)	1 (3%)	1 (3%)	3	0
2	D	29/30 (97%)	29 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	98/102 (96%)	94 (96%)	3 (3%)	1 (1%)	14 3

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	29	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	20/20 (100%)	19 (95%)	1 (5%)	24 4
1	C	22/20 (110%)	22 (100%)	0	100 100
2	B	26/25 (104%)	25 (96%)	1 (4%)	33 7
2	D	26/25 (104%)	23 (88%)	3 (12%)	5 0
All	All	94/90 (104%)	89 (95%)	5 (5%)	28 3

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	GLU
2	B	27	THR
2	D	2	VAL
2	D	21[A]	GLU
2	D	21[B]	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	GLN
1	A	21	ASN
1	C	15	GLN
1	C	18	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.