



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 4, 2024 – 03:50 AM EST

PDB ID : 1ZNO  
Title : Crystal Structure of VC702 from Vibrio Cholerae, Northeast Structural Genomics Consortium Target: VcP1  
Authors : Ni, S.; Forouhar, F.; Bussiere, D.E.; Robinson, H.; Kennedy, M.A.; Northeast Structural Genomics Consortium (NESG)  
Deposited on : 2005-05-11  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	183	
1	B	183	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MG	A	184	-	-	-	X

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2747 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hypothetical UPF0244 protein VC0702.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	S				Se
1	A	165	1299	819	230	244	1	5	0	0	0
1	B	165	1301	819	234	242	1	5	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	modified residue	UNP Q9KU27
A	10	MSE	MET	modified residue	UNP Q9KU27
A	55	MSE	MET	modified residue	UNP Q9KU27
A	97	MSE	MET	modified residue	UNP Q9KU27
A	114	MSE	MET	modified residue	UNP Q9KU27
A	133	MSE	MET	modified residue	UNP Q9KU27
B	1	MSE	MET	modified residue	UNP Q9KU27
B	10	MSE	MET	modified residue	UNP Q9KU27
B	55	MSE	MET	modified residue	UNP Q9KU27
B	97	MSE	MET	modified residue	UNP Q9KU27
B	114	MSE	MET	modified residue	UNP Q9KU27
B	133	MSE	MET	modified residue	UNP Q9KU27

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		
2	B	2	Total	Mg	0	0
			2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	62	Total O 62 62	0	0
3	B	82	Total O 82 82	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Hypothetical UPF0244 protein VC0702



- Molecule 1: Hypothetical UPF0244 protein VC0702



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	66.61Å 88.12Å 118.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.00 30.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.00-2.00) 99.9 (30.00-2.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.32 (at 2.00Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.230 , 0.263 0.246 , 0.243	Depositor DCC
$R_{free}$ test set	1226 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.7	Xtrriage
Anisotropy	0.604	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 46.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	2747	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.88% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.32	0/1320	0.56	0/1779
1	B	0.33	0/1322	0.57	0/1779
All	All	0.32	0/2642	0.57	0/3558

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1299	0	1291	56	0
1	B	1301	0	1298	43	0
2	A	1	0	0	0	0
2	B	2	0	0	0	0
3	A	62	0	0	0	0
3	B	82	0	0	3	2
All	All	2747	0	2589	94	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:156:HIS:HA	3:B:487:HOH:O	1.67	0.94
1:A:10:MSE:HE2	1:A:38:GLU:HB2	1.63	0.81
1:B:84:LEU:HG	1:B:97:MSE:SE	2.35	0.77
1:B:104:GLN:HG3	1:B:174:PHE:CE1	2.21	0.76
1:B:169:LEU:O	1:B:172:ILE:HD13	1.91	0.70
1:B:142:ILE:HD11	1:B:151:LEU:HB2	1.77	0.65
1:A:153:THR:HG22	1:B:172:ILE:HD11	1.79	0.64
1:B:154:ARG:HD2	3:B:455:HOH:O	1.98	0.63
1:A:11:ARG:HE	1:A:79:GLU:HB2	1.64	0.61
1:A:172:ILE:HD11	1:B:153:THR:HG22	1.83	0.61
1:A:97:MSE:C	1:A:98:ILE:HD12	2.21	0.61
1:B:169:LEU:HA	1:B:172:ILE:HD13	1.83	0.59
1:A:143:LYS:HA	1:A:143:LYS:NZ	2.18	0.59
1:A:167:LEU:O	1:A:171:LEU:HD13	2.03	0.59
1:B:15:ILE:CD1	1:B:26:VAL:HG11	2.34	0.58
1:B:178:GLU:H	1:B:178:GLU:CD	2.07	0.57
1:B:15:ILE:HD11	1:B:26:VAL:HG11	1.86	0.57
1:B:127:LYS:NZ	1:B:130:GLY:HA3	2.20	0.56
1:A:74:ARG:HB3	1:A:75:HIS:ND1	2.21	0.56
1:A:134:ASP:O	1:A:140:GLU:HB3	2.05	0.56
1:A:13:ILE:HG12	1:A:80:TYR:HB2	1.88	0.55
1:A:169:LEU:O	1:A:172:ILE:HG12	2.07	0.55
1:A:122:ARG:HD2	1:A:132:VAL:HG13	1.89	0.55
1:B:11:ARG:HE	1:B:79:GLU:HB2	1.72	0.54
1:A:16:ALA:HB1	1:A:67:ARG:HB3	1.89	0.53
1:B:55:MSE:SE	1:B:88:ILE:HD11	2.59	0.53
1:A:118:LEU:HD13	1:A:118:LEU:O	2.09	0.52
1:A:55:MSE:SE	1:A:88:ILE:HD11	2.59	0.51
1:A:64:ALA:CB	1:A:98:ILE:HD13	2.40	0.51
1:B:121:GLU:HG3	1:B:122:ARG:N	2.25	0.51
1:A:116:PRO:HD2	1:A:119:VAL:HG21	1.93	0.51
1:A:177:PRO:HD2	1:A:178:GLU:OE2	2.10	0.51
1:B:22:LYS:HE2	1:B:84:LEU:HD22	1.93	0.51
1:A:164:HIS:O	1:A:168:ILE:HG12	2.11	0.51
1:A:117:PRO:HD2	1:B:179:HIS:O	2.11	0.51
1:A:10:MSE:HG2	1:A:39:TRP:HA	1.92	0.50
1:A:178:GLU:CD	1:A:178:GLU:H	2.13	0.50
1:A:172:ILE:HB	1:A:173:PRO:HD3	1.94	0.50
1:A:64:ALA:HB1	1:A:98:ILE:HD13	1.94	0.50
1:B:16:ALA:HB1	1:B:67:ARG:HB3	1.94	0.49
1:A:133:MSE:HG3	1:A:143:LYS:HE3	1.94	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:115:LEU:HB2	1:A:120:LEU:HD21	1.94	0.49
1:B:12:LYS:HE3	3:B:421:HOH:O	2.13	0.49
1:A:143:LYS:O	1:A:147:GLY:HA2	2.13	0.49
1:A:149:ILE:HD12	1:A:159:ARG:HA	1.95	0.49
1:B:44:VAL:HG12	1:B:45:SER:N	2.28	0.49
1:B:142:ILE:HD12	1:B:145:LYS:HD3	1.95	0.49
1:B:169:LEU:HA	1:B:172:ILE:CD1	2.43	0.48
1:A:12:LYS:HG3	1:A:78:ALA:HA	1.96	0.47
1:A:142:ILE:HG23	1:A:145:LYS:HD2	1.95	0.47
1:A:122:ARG:HD2	1:A:132:VAL:CG1	2.44	0.47
1:B:39:TRP:N	1:B:39:TRP:CD1	2.82	0.47
1:B:97:MSE:HG3	1:B:167:LEU:CD2	2.45	0.47
1:A:142:ILE:O	1:A:145:LYS:HB2	2.15	0.47
1:B:10:MSE:HE2	1:B:38:GLU:HB3	1.97	0.47
1:B:149:ILE:HD13	1:B:162:VAL:HG21	1.97	0.46
1:B:78:ALA:HB3	1:B:81:TYR:CZ	2.51	0.46
1:A:129:LEU:O	1:A:132:VAL:HG23	2.15	0.46
1:B:150:GLY:HA2	1:B:157:LEU:O	2.16	0.45
1:A:88:ILE:N	1:A:88:ILE:HD13	2.32	0.45
1:A:44:VAL:HG12	1:A:45:SER:N	2.31	0.45
1:A:54:PRO:O	1:A:88:ILE:HD13	2.17	0.45
1:B:18:GLN:NE2	1:B:45:SER:HB2	2.32	0.45
1:A:48:SER:O	1:A:49:GLU:HB2	2.17	0.44
1:B:169:LEU:CA	1:B:172:ILE:HD13	2.47	0.44
1:B:123:LEU:HD23	1:B:123:LEU:O	2.17	0.44
1:B:97:MSE:HG3	1:B:167:LEU:HD23	1.99	0.44
1:A:142:ILE:CG2	1:A:145:LYS:HD2	2.48	0.44
1:A:33:VAL:HG23	1:A:168:ILE:HD12	1.99	0.43
1:B:88:ILE:N	1:B:88:ILE:HD13	2.34	0.43
1:A:144:GLN:O	1:A:144:GLN:HG3	2.19	0.43
1:B:177:PRO:HD2	1:B:178:GLU:OE2	2.19	0.43
1:A:143:LYS:HZ2	1:A:148:ALA:N	2.17	0.43
1:B:169:LEU:C	1:B:172:ILE:HD13	2.40	0.43
1:A:123:LEU:C	1:A:129:LEU:HB2	2.39	0.42
1:A:169:LEU:HD22	1:B:113:LEU:HD12	2.01	0.42
1:A:91:ASN:HA	1:A:120:LEU:HD11	2.00	0.42
1:A:143:LYS:HA	1:A:143:LYS:HZ3	1.85	0.42
1:A:78:ALA:HB3	1:A:81:TYR:CZ	2.54	0.42
1:A:133:MSE:C	1:A:135:GLU:H	2.23	0.42
1:A:129:LEU:C	1:A:131:ASP:H	2.23	0.41
1:B:135:GLU:N	1:B:135:GLU:OE2	2.52	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:18:GLN:HG2	1:A:43:GLY:O	2.21	0.41
1:A:10:MSE:HG3	1:A:38:GLU:OE1	2.20	0.41
1:A:97:MSE:HE2	1:A:166:ALA:HB3	2.02	0.41
1:A:11:ARG:HH21	1:A:79:GLU:HB2	1.86	0.41
1:B:172:ILE:HB	1:B:173:PRO:HD3	2.03	0.41
1:A:150:GLY:HA2	1:A:157:LEU:O	2.21	0.41
1:A:48:SER:HB3	1:A:67:ARG:CZ	2.51	0.40
1:B:101:SER:OG	1:B:104:GLN:HG2	2.21	0.40
1:A:172:ILE:HD11	1:B:153:THR:CG2	2.50	0.40
1:A:151:LEU:HD22	1:A:151:LEU:HA	1.92	0.40
1:B:27:ARG:HE	1:B:41:PHE:HB2	1.85	0.40
1:B:48:SER:O	1:B:49:GLU:HB2	2.22	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:483:HOH:O	3:B:483:HOH:O[3_556]	1.25	0.95
3:B:486:HOH:O	3:B:487:HOH:O[3_556]	2.09	0.11

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	159/183 (87%)	145 (91%)	12 (8%)	2 (1%)	10	5
1	B	159/183 (87%)	153 (96%)	5 (3%)	1 (1%)	22	17
All	All	318/366 (87%)	298 (94%)	17 (5%)	3 (1%)	14	10

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	122	ARG
1	A	37	GLN
1	B	133	MSE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	141/150 (94%)	133 (94%)	8 (6%)	17	14
1	B	140/150 (93%)	134 (96%)	6 (4%)	25	23
All	All	281/300 (94%)	267 (95%)	14 (5%)	20	18

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38	GLU
1	A	39	TRP
1	A	73	GLN
1	A	88	ILE
1	A	97	MSE
1	A	113	LEU
1	A	143	LYS
1	A	151	LEU
1	B	39	TRP
1	B	88	ILE
1	B	97	MSE
1	B	113	LEU
1	B	120	LEU
1	B	178	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	GLN
1	A	24	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	53	GLN
1	A	73	GLN
1	A	103	GLN
1	A	144	GLN
1	B	18	GLN
1	B	24	ASN
1	B	73	GLN
1	B	144	GLN
1	B	179	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	160/183 (87%)	0.75	27 (16%) <b>5</b> <b>4</b>	14, 30, 69, 88	0
1	B	160/183 (87%)	0.53	16 (10%) <b>14</b> <b>13</b>	13, 27, 72, 80	0
All	All	320/366 (87%)	0.64	43 (13%) <b>8</b> <b>7</b>	13, 29, 71, 88	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	130	GLY	5.6
1	A	136	VAL	5.1
1	A	39	TRP	4.8
1	B	142	ILE	4.6
1	B	132	VAL	4.4
1	A	141	ASN	4.2
1	B	135	GLU	3.8
1	A	123	LEU	3.7
1	A	11	ARG	3.6
1	A	102	ASP	3.6
1	B	39	TRP	3.6
1	B	123	LEU	3.5
1	A	37	GLN	3.5
1	A	34	PHE	3.4
1	A	36	ASP	3.4
1	B	131	ASP	3.2
1	A	35	PRO	3.2
1	B	113	LEU	3.2
1	A	144	GLN	3.1
1	A	132	VAL	3.1
1	A	129	LEU	3.0
1	A	88	ILE	2.7
1	A	143	LYS	2.7
1	B	127	LYS	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	182	SER	2.6
1	A	181	PRO	2.6
1	B	124	ARG	2.5
1	B	102	ASP	2.5
1	A	142	ILE	2.5
1	A	74	ARG	2.5
1	A	135	GLU	2.4
1	A	131	ASP	2.4
1	B	74	ARG	2.3
1	A	146	GLY	2.2
1	A	134	ASP	2.2
1	B	122	ARG	2.2
1	A	130	GLY	2.2
1	A	113	LEU	2.2
1	A	120	LEU	2.1
1	B	126	ALA	2.1
1	B	125	GLN	2.1
1	A	119	VAL	2.1
1	B	182	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MG	A	184	1/1	0.67	0.60	58,58,58,58	1
2	MG	B	184	1/1	0.87	0.49	40,40,40,40	1
2	MG	B	185	1/1	0.96	0.12	28,28,28,28	0



## 6.5 Other polymers [i](#)

There are no such residues in this entry.