



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 8, 2020 – 11:34 AM BST

PDB ID : 4ZO6
Title : Crystal Structure of mutant (D270A) beta-glucosidase from *Listeria innocua* in complex with cellobiose
Authors : Nakajima, M.; Yoshida, R.; Miyanaga, A.; Abe, K.; Takahashi, Y.; Sugimoto, N.; Toyozumi, H.; Nakai, H.; Kitaoka, M.; Taguchi, H.
Deposited on : 2015-05-06
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	731	 % 90% 8% ..
1	B	731	 % 91% 8% .
2	C	2	 100%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 12144 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lin1840 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	724	Total 5600	C 3544	N 929	O 1102	S 25	0	1	0
1	B	724	Total 5592	C 3539	N 928	O 1101	S 24	0	0	0

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	2	GLU	LYS	engineered mutation	UNP Q92AS9
A	270	ALA	ASP	engineered mutation	UNP Q92AS9
A	724	LEU	-	expression tag	UNP Q92AS9
A	725	GLU	-	expression tag	UNP Q92AS9
A	726	HIS	-	expression tag	UNP Q92AS9
A	727	HIS	-	expression tag	UNP Q92AS9
A	728	HIS	-	expression tag	UNP Q92AS9
A	729	HIS	-	expression tag	UNP Q92AS9
A	730	HIS	-	expression tag	UNP Q92AS9
A	731	HIS	-	expression tag	UNP Q92AS9
B	2	GLU	LYS	engineered mutation	UNP Q92AS9
B	270	ALA	ASP	engineered mutation	UNP Q92AS9
B	724	LEU	-	expression tag	UNP Q92AS9
B	725	GLU	-	expression tag	UNP Q92AS9
B	726	HIS	-	expression tag	UNP Q92AS9
B	727	HIS	-	expression tag	UNP Q92AS9
B	728	HIS	-	expression tag	UNP Q92AS9
B	729	HIS	-	expression tag	UNP Q92AS9
B	730	HIS	-	expression tag	UNP Q92AS9
B	731	HIS	-	expression tag	UNP Q92AS9

- Molecule 2 is an oligosaccharide called beta-D-glucopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	C	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 3 is beta-D-glucopyranose (three-letter code: BGC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			12	6	6		
3	A	1	Total	C	O	0	0
			12	6	6		
3	A	1	Total	C	O	0	0
			12	6	6		
3	A	1	Total	C	O	0	0
			12	6	6		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 12 6 6	0	1

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Mg 1 1	0	0
5	A	1	Total Mg 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	447	Total O 447 447	0	0
6	B	420	Total O 420 420	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Lin1840 protein



- Molecule 1: Lin1840 protein



- Molecule 2: beta-D-glucopyranose-(1-4)-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	89.79Å 95.35Å 215.38Å 90.00° 96.30° 90.00°	Depositor
Resolution (Å)	32.56 – 2.00 32.56 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.6 (32.56-2.00) 96.7 (32.56-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.91 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.183 , 0.255 0.192 , 0.260	Depositor DCC
R_{free} test set	5906 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	15.2	Xtrriage
Anisotropy	0.136	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 49.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	12144	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.83	0/5702	0.92	6/7727 (0.1%)
1	B	0.82	0/5694	0.90	11/7717 (0.1%)
All	All	0.83	0/11396	0.91	17/15444 (0.1%)

There are no bond length outliers.

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	655	ARG	NE-CZ-NH2	9.28	124.94	120.30
1	A	175	ASP	CB-CG-OD1	7.23	124.81	118.30
1	B	655	ARG	NE-CZ-NH1	-7.19	116.70	120.30
1	A	364	ARG	NE-CZ-NH2	-6.78	116.91	120.30
1	B	149	ARG	NE-CZ-NH2	-6.38	117.11	120.30
1	B	149	ARG	NE-CZ-NH1	6.07	123.33	120.30
1	B	364	ARG	NE-CZ-NH2	-5.97	117.32	120.30
1	B	326	ASP	CB-CG-OD1	5.90	123.61	118.30
1	B	534	ARG	NE-CZ-NH1	5.80	123.20	120.30
1	A	258	ARG	NE-CZ-NH1	-5.68	117.46	120.30
1	B	589	ASP	CB-CG-OD1	5.41	123.17	118.30
1	B	426	ARG	NE-CZ-NH1	5.37	122.99	120.30
1	A	702	ARG	NE-CZ-NH1	5.37	122.98	120.30
1	A	655	ARG	NE-CZ-NH1	5.31	122.96	120.30
1	B	368	ARG	NE-CZ-NH2	5.31	122.95	120.30
1	A	501	VAL	CB-CA-C	-5.16	101.59	111.40
1	B	254	ARG	NE-CZ-NH2	-5.12	117.74	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5600	0	5554	35	0
1	B	5592	0	5546	25	0
2	C	23	0	21	2	0
3	A	48	0	47	1	0
4	A	12	0	16	2	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	447	0	0	7	0
6	B	420	0	0	1	0
All	All	12144	0	11184	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:804[A]:GOL:H31	6:A:903:HOH:O	1.84	0.77
4:A:804[A]:GOL:O3	6:A:901:HOH:O	2.00	0.72
1:A:723:VAL:HG12	1:A:724:LEU:HD13	1.76	0.68
2:C:1:BGC:H6C2	2:C:2:BGC:O2	2.01	0.60
1:B:385:LEU:HD21	1:B:501:VAL:HG21	1.84	0.58
1:A:698:PRO:HD2	6:A:1325:HOH:O	2.02	0.58
1:A:244:ILE:HD11	1:B:653:ILE:HD13	1.88	0.56
1:A:250:LYS:NZ	1:A:295:GLU:OE1	2.39	0.55
1:A:258:ARG:HD3	6:A:1265:HOH:O	2.06	0.55
1:A:244:ILE:CD1	1:B:652:SER:O	2.55	0.55
1:B:451:LYS:CD	1:B:496:LEU:HD22	2.37	0.55
1:A:616:LYS:HE3	1:A:616:LYS:O	2.08	0.54
1:B:271:TRP:C	1:B:271:TRP:CD1	2.81	0.53
1:B:9:LEU:O	1:B:13:MET:HG3	2.08	0.53
1:A:15:LEU:HD23	1:A:312:LYS:HE2	1.91	0.52
1:B:409:TRP:CZ2	2:C:2:BGC:H6C1	2.45	0.52
1:B:368:ARG:NH1	1:B:534:ARG:HD3	2.25	0.52
1:A:567:HIS:HD2	1:A:568:LEU:O	1.95	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:9:LEU:O	1:A:13:MET:HG3	2.12	0.49
1:A:454:VAL:HG12	1:A:498:LYS:HG3	1.95	0.49
1:A:61:LEU:CD2	1:A:271:TRP:CZ3	2.96	0.48
1:A:147:TRP:CZ3	1:A:197:GLY:HA2	2.47	0.48
1:A:244:ILE:HD11	1:B:652:SER:O	2.15	0.47
1:A:630:ILE:HG23	1:A:668:LEU:HD11	1.96	0.47
1:A:719:VAL:O	6:A:902:HOH:O	2.21	0.47
1:A:81:ARG:NH1	1:A:82:LEU:HD21	2.30	0.47
1:A:220:LEU:N	1:A:221:PRO:CD	2.78	0.46
1:B:220:LEU:N	1:B:221:PRO:CD	2.80	0.45
1:A:494:GLN:HG3	6:A:1239:HOH:O	2.15	0.45
1:A:409:TRP:CE2	3:A:803:BGC:H4	2.52	0.45
1:B:368:ARG:HD3	1:B:538:ASP:OD2	2.16	0.45
1:B:639:LYS:HE2	1:B:667:ALA:HB2	1.99	0.45
1:B:658:LYS:HE2	1:B:709:SER:O	2.17	0.44
1:B:700:LEU:HD11	1:B:716:GLU:HB3	1.99	0.44
1:A:100:PHE:CG	1:A:118:MET:HG2	2.53	0.44
1:A:271:TRP:CD1	1:A:271:TRP:C	2.91	0.44
1:B:336:LYS:HG2	1:B:341:LEU:HD12	1.98	0.44
1:B:61:LEU:HD21	1:B:271:TRP:CZ3	2.53	0.44
1:A:244:ILE:CD1	1:B:653:ILE:HD13	2.48	0.44
1:A:659:GLU:HB2	1:A:661:LYS:HE2	2.00	0.43
1:B:353:ASP:OD1	1:B:353:ASP:C	2.56	0.43
1:A:723:VAL:O	1:A:724:LEU:HB2	2.19	0.43
1:A:645:TYR:HB2	1:A:704:PHE:HB2	2.01	0.43
1:A:658:LYS:HE2	1:A:709:SER:O	2.17	0.43
1:A:478:ALA:O	1:B:559:GLY:HA3	2.18	0.43
1:A:235:MET:HA	1:A:268:ILE:O	2.19	0.43
1:A:2:GLU:HB2	1:A:5:LYS:CE	2.49	0.42
1:B:258:ARG:HD3	6:B:1199:HOH:O	2.18	0.42
1:A:454:VAL:CG1	1:A:498:LYS:HG3	2.49	0.42
1:A:76:TYR:CD2	1:A:86:LEU:HD11	2.55	0.41
1:B:147:TRP:CZ3	1:B:197:GLY:HA2	2.56	0.41
1:B:350:LYS:HA	1:B:350:LYS:HD3	1.88	0.41
1:B:406:LEU:HA	1:B:529:GLY:HA2	2.03	0.41
1:A:636:ILE:HD12	6:A:1343:HOH:O	2.21	0.41
1:A:233:LEU:HD12	1:A:233:LEU:N	2.36	0.41
1:A:531:GLU:OE1	1:A:534:ARG:HD2	2.21	0.41
1:B:100:PHE:CG	1:B:118:MET:HG2	2.55	0.41
1:B:235:MET:HA	1:B:268:ILE:O	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	723/731 (99%)	702 (97%)	21 (3%)	0	100	100
1	B	722/731 (99%)	704 (98%)	18 (2%)	0	100	100
All	All	1445/1462 (99%)	1406 (97%)	39 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	605/611 (99%)	590 (98%)	15 (2%)	47	49
1	B	604/611 (99%)	585 (97%)	19 (3%)	40	40
All	All	1209/1222 (99%)	1175 (97%)	34 (3%)	43	44

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	GLU
1	A	9	LEU
1	A	129	ASP
1	A	151	MET
1	A	250	LYS
1	A	308	ILE
1	A	320	LEU
1	A	430	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	438	GLU
1	A	570	THR
1	A	579	LYS
1	A	614	LEU
1	A	616	LYS
1	A	683	GLU
1	A	724	LEU
1	B	11	SER
1	B	78	LYS
1	B	129	ASP
1	B	151	MET
1	B	210	MET
1	B	284	ARG
1	B	308	ILE
1	B	320	LEU
1	B	333	LEU
1	B	350	LYS
1	B	383	ARG
1	B	491	LYS
1	B	610	LYS
1	B	613	SER
1	B	616	LYS
1	B	680	LEU
1	B	691	GLN
1	B	692	LEU
1	B	724	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	182	ASN
1	A	455	GLN
1	A	567	HIS
1	B	182	ASN
1	B	625	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BGC	C	1	2	12,12,12	0.63	0	17,17,17	1.83	5 (29%)
2	BGC	C	2	2	11,11,12	0.93	1 (9%)	15,15,17	2.97	8 (53%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	C	1	2	-	1/2/22/22	0/1/1/1
2	BGC	C	2	2	-	1/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	2	BGC	C2-C3	2.04	1.55	1.52

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	BGC	C6-C5-C4	6.51	128.25	113.00
2	C	2	BGC	O5-C5-C6	-4.65	99.91	107.20
2	C	2	BGC	C3-C4-C5	-4.54	102.15	110.24
2	C	2	BGC	O5-C1-C2	-4.49	103.83	110.77
2	C	1	BGC	C1-O5-C5	3.56	120.38	113.66
2	C	1	BGC	O5-C5-C4	3.28	115.65	109.69

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	BGC	O4-C4-C3	-3.15	103.07	110.35
2	C	2	BGC	O6-C6-C5	-2.83	101.57	111.29
2	C	1	BGC	O5-C5-C6	-2.63	99.89	106.44
2	C	2	BGC	O4-C4-C5	2.60	115.76	109.30
2	C	2	BGC	C1-O5-C5	-2.47	108.85	112.19
2	C	1	BGC	O2-C2-C3	-2.32	104.98	110.35
2	C	2	BGC	C1-C2-C3	2.06	112.20	109.67

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	2	BGC	O5-C5-C6-O6
2	C	1	BGC	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	BGC	1	0
2	C	2	BGC	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	BGC	A	803	-	12,12,12	1.26	2 (16%)	17,17,17	3.86	9 (52%)
4	GOL	A	804[B]	-	5,5,5	0.30	0	5,5,5	0.35	0
3	BGC	A	802	-	12,12,12	0.82	0	17,17,17	2.08	6 (35%)
3	BGC	A	806	-	12,12,12	0.58	0	17,17,17	1.24	2 (11%)
3	BGC	A	801	-	12,12,12	0.74	0	17,17,17	2.11	6 (35%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	A	804[A]	-	5,5,5	0.23	0	5,5,5	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BGC	A	803	-	-	0/2/22/22	0/1/1/1
4	GOL	A	804[B]	-	-	2/4/4/4	-
3	BGC	A	802	-	-	0/2/22/22	0/1/1/1
3	BGC	A	806	-	-	2/2/22/22	0/1/1/1
3	BGC	A	801	-	-	2/2/22/22	0/1/1/1
4	GOL	A	804[A]	-	-	2/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	803	BGC	O1-C1	2.31	1.46	1.39
3	A	803	BGC	C3-C2	2.23	1.58	1.52

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	803	BGC	C6-C5-C4	8.31	132.46	113.00
3	A	803	BGC	C3-C4-C5	-6.90	97.93	110.24
3	A	803	BGC	O5-C1-C2	-6.76	98.23	110.28
3	A	803	BGC	O4-C4-C5	5.48	122.92	109.30
3	A	801	BGC	O3-C3-C4	-4.69	99.50	110.35
3	A	803	BGC	O1-C1-C2	4.20	120.85	109.03
3	A	802	BGC	C1-C2-C3	-4.03	101.96	110.31
3	A	802	BGC	C1-O5-C5	-3.90	106.31	113.66
3	A	803	BGC	O6-C6-C5	-3.73	98.48	111.29
3	A	801	BGC	C1-O5-C5	3.47	120.20	113.66
3	A	803	BGC	C1-O5-C5	-3.36	107.32	113.66
3	A	802	BGC	O5-C1-C2	-3.36	104.28	110.28
3	A	801	BGC	O1-C1-O5	-3.23	100.69	110.38
3	A	801	BGC	C3-C4-C5	2.84	115.30	110.24
3	A	803	BGC	O5-C5-C6	-2.80	99.47	106.44
3	A	802	BGC	O2-C2-C1	2.61	115.21	109.16
3	A	801	BGC	O2-C2-C3	2.55	116.25	110.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	806	BGC	O3-C3-C2	-2.50	104.56	110.35
3	A	801	BGC	O5-C5-C4	2.38	114.02	109.69
3	A	802	BGC	O4-C4-C3	-2.19	105.29	110.35
3	A	802	BGC	C4-C3-C2	-2.10	107.16	110.82
3	A	806	BGC	O5-C5-C6	2.06	111.55	106.44
3	A	803	BGC	O3-C3-C2	2.05	115.09	110.35

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	804[A]	GOL	C1-C2-C3-O3
3	A	801	BGC	O5-C5-C6-O6
3	A	801	BGC	C4-C5-C6-O6
3	A	806	BGC	C4-C5-C6-O6
3	A	806	BGC	O5-C5-C6-O6
4	A	804[A]	GOL	O2-C2-C3-O3
4	A	804[B]	GOL	O1-C1-C2-O2
4	A	804[B]	GOL	O1-C1-C2-C3

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	803	BGC	1	0
4	A	804[A]	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	724/731 (99%)	-0.33	7 (0%) 82 81	7, 14, 29, 47	0
1	B	724/731 (99%)	-0.30	10 (1%) 75 74	7, 16, 32, 51	0
All	All	1448/1462 (99%)	-0.32	17 (1%) 79 78	7, 15, 31, 51	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	351	ASN	5.0
1	A	576	PRO	3.9
1	B	576	PRO	3.5
1	B	580	GLY	3.5
1	A	580	GLY	3.0
1	A	38	GLU	2.8
1	A	579	LYS	2.7
1	B	579	LYS	2.7
1	B	352	ASN	2.4
1	B	577	GLU	2.3
1	A	39	LEU	2.3
1	B	350	LYS	2.2
1	B	353	ASP	2.2
1	B	356	LYS	2.1
1	A	313	GLY	2.1
1	A	42	PRO	2.1
1	B	37	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BGC	C	1	12/12	0.87	0.20	33,38,43,58	0
2	BGC	C	2	11/12	0.92	0.12	16,26,30,36	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	BGC	A	803	12/12	0.86	0.15	16,24,29,33	0
3	BGC	A	802	12/12	0.93	0.15	16,28,33,36	0
4	GOL	A	804[B]	6/6	0.94	0.17	13,16,16,18	6
3	BGC	A	801	12/12	0.94	0.10	13,18,21,25	0
4	GOL	A	804[A]	6/6	0.94	0.17	15,17,17,17	6
3	BGC	A	806	12/12	0.95	0.08	16,17,19,21	0
5	MG	B	802	1/1	0.98	0.05	15,15,15,15	0
5	MG	A	805	1/1	0.98	0.06	15,15,15,15	0

6.5 Other polymers [i](#)

There are no such residues in this entry.