



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 17, 2024 – 10:10 PM JST

PDB ID : 8ZRZ
Title : The 1.26 angstrom resolution structure of Bacillus cereus beta-amylase in complex with maltose
Authors : Mikami, B.; Hirata, A.
Deposited on : 2024-06-05
Resolution : 1.26 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.002 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.38.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1447 (1.28-1.24)
Clashscore	180529	1571 (1.28-1.24)
Ramachandran outliers	177936	1538 (1.28-1.24)
Sidechain outliers	177891	1537 (1.28-1.24)
RSRZ outliers	164620	1447 (1.28-1.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	516	 89% 10%
2	C	2	 50% 50%
2	D	2	 100%
2	F	2	 100%
3	E	2	 100%
4	B	4	 50% 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GLC	B	3[B]	X	-	-	-
7	GOL	A	604	-	X	-	-
7	GOL	A	605	-	X	-	-
7	GOL	A	606[A]	-	X	-	-
7	GOL	A	606[B]	-	X	-	-
7	GOL	A	607	-	X	-	-
7	GOL	A	608	-	X	-	-
7	GOL	A	609	-	X	-	-
7	GOL	A	610	-	X	-	-
7	GOL	A	611	-	X	-	-

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 5480 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-amylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	516	4371	2819	706	824	22	0	46	0

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	C	2	23	12	11	0	0	0
2	F	2	23	12	11	0	0	0
2	D	2	23	12	11	0	0	0

- Molecule 3 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
3	E	2	23	12	11	0	0	0

- Molecule 4 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
4	B	4	Total	C	O	0	2	0
			58	30	28			

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Ca	0	0
			1	1		

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	1
			12	6	6		
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is beta-D-glucopyranose (three-letter code: BGC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total C O 12 6 6	0	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	867	Total O 887 887	0	20

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta-amylase




- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose




- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



- Molecule 3: alpha-D-glucopyranose-(1-4)-beta-D-glucopyranose



- Molecule 4: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain B:  50% 50%

GLC1
GLC2
GLC3
GLC4

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	56.97Å 89.84Å 66.03Å 90.00° 103.09° 90.00°	Depositor
Resolution (Å)	10.00 – 1.26 10.00 – 1.26	Depositor EDS
% Data completeness (in resolution range)	93.4 (10.00-1.26) 95.3 (10.00-1.26)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.63 (at 1.26Å)	Xtrriage
Refinement program	SHELXL-97 2018/3	Depositor
R, R_{free}	0.124 , 0.165 0.121 , 0.158	Depositor DCC
R_{free} test set	8555 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	12.4	Xtrriage
Anisotropy	0.611	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 85.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	5480	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, GLC, BGC, CA, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.75	2/4618 (0.0%)	1.20	27/6267 (0.4%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	3	ASN	C-N	7.54	1.46	1.33
1	A	4	GLY	C-N	5.76	1.47	1.34

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	331[A]	CYS	CA-CB-SG	10.78	133.40	114.00
1	A	331[B]	CYS	CA-CB-SG	10.78	133.40	114.00
1	A	380[A]	ARG	NE-CZ-NH1	6.68	123.64	120.30
1	A	380[B]	ARG	NE-CZ-NH1	6.68	123.64	120.30
1	A	189	ARG	NE-CZ-NH1	6.43	123.52	120.30
1	A	36	ARG	NE-CZ-NH2	-6.37	117.11	120.30
1	A	329[A]	PHE	CB-CG-CD2	-5.91	116.66	120.80
1	A	329[B]	PHE	CB-CG-CD2	-5.91	116.66	120.80
1	A	332[A]	LEU	C-N-CA	5.61	135.72	121.70
1	A	332[B]	LEU	C-N-CA	5.61	135.72	121.70
1	A	378	TYR	CB-CG-CD1	5.45	124.27	121.00
1	A	158[A]	ASP	CB-CA-C	5.40	121.19	110.40
1	A	158[B]	ASP	CB-CA-C	5.40	121.19	110.40
1	A	331[A]	CYS	CB-CA-C	5.33	121.07	110.40
1	A	331[B]	CYS	CB-CA-C	5.33	121.07	110.40
1	A	178	TYR	CA-CB-CG	5.31	123.50	113.40
1	A	80	MET	CG-SD-CE	-5.29	91.74	100.20
1	A	146	TYR	CB-CG-CD1	5.25	124.15	121.00
1	A	454[A]	TYR	CB-CG-CD1	5.23	124.14	121.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	454[B]	TYR	CB-CG-CD1	5.23	124.14	121.00
1	A	71	PHE	CB-CG-CD1	5.18	124.43	120.80
1	A	266[B]	LYS	CB-CA-C	5.17	120.75	110.40
1	A	266[C]	LYS	CB-CA-C	5.17	120.75	110.40
1	A	266[D]	LYS	CB-CA-C	5.17	120.75	110.40
1	A	136[A]	ASP	CB-CG-OD1	5.07	122.86	118.30
1	A	136[B]	ASP	CB-CG-OD1	5.07	122.86	118.30
1	A	403	TYR	CZ-CE2-CD2	5.06	124.36	119.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4371	0	4263	40	0
2	C	23	0	21	1	0
2	D	23	0	21	0	0
2	F	23	0	21	0	0
3	E	23	0	21	0	0
4	B	58	0	47	3	0
5	A	1	0	0	0	0
6	A	5	0	0	0	0
7	A	54	0	36	7	0
8	A	12	0	12	2	0
9	A	887	0	0	15	0
All	All	5480	0	4442	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:330[A]:THR:N	9:A:702:HOH:O	2.07	0.81

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:2[A]:GLC:H62	4:B:3[A]:GLC:O1	1.85	0.77
1:A:331[A]:CYS:HB2	1:A:369:ALA:HB2	1.67	0.76
1:A:493:LYS:HE2	9:A:1284:HOH:O	1.85	0.76
1:A:225:LEU:HA	9:A:1271:HOH:O	1.85	0.75
1:A:331[B]:CYS:HB2	9:A:946:HOH:O	1.93	0.69
1:A:487[B]:SER:OG	1:A:493:LYS:HD2	1.93	0.68
9:A:1104:HOH:O	4:B:3[A]:GLC:H3	1.92	0.68
7:A:605:GOL:H12	9:A:916:HOH:O	1.97	0.63
1:A:352[A]:ASN:ND2	7:A:608:GOL:O2	2.27	0.62
1:A:502:TRP:NE1	8:A:602:BGC:H6C1	2.14	0.62
1:A:266[C]:LYS:HD2	1:A:323:ALA:HB2	1.82	0.62
1:A:266[C]:LYS:HE2	1:A:323:ALA:N	2.15	0.62
1:A:331[B]:CYS:HB3	9:A:701:HOH:O	2.00	0.60
1:A:375:GLU:HB2	1:A:410:LYS:HD3	1.82	0.60
1:A:508:LYS:HD3	9:A:1473[B]:HOH:O	2.00	0.59
7:A:606[A]:GOL:H12	9:A:728:HOH:O	2.03	0.57
1:A:502:TRP:HE1	8:A:602:BGC:H6C1	1.69	0.57
1:A:2[B]:VAL:CG1	1:A:454[B]:TYR:CE1	2.88	0.56
1:A:2[B]:VAL:CG1	1:A:454[B]:TYR:HE1	2.19	0.56
1:A:2[B]:VAL:HG12	1:A:454[B]:TYR:CE1	2.41	0.55
1:A:320:PHE:CD2	1:A:327[A]:VAL:HG22	2.40	0.55
1:A:347:LYS:HE3	1:A:384[B]:MET:SD	2.48	0.54
1:A:225:LEU:HD23	9:A:1271:HOH:O	2.09	0.52
1:A:162[A]:LYS:HG3	1:A:283:PRO:O	2.09	0.52
1:A:219[A]:LYS:HD3	9:A:809:HOH:O	2.11	0.49
1:A:224:LYS:HG3	7:A:610:GOL:C1	2.43	0.48
1:A:126:ASN:OD1	1:A:128[B]:GLU:HG2	2.14	0.48
1:A:329[B]:PHE:O	1:A:366:GLY:HA2	2.13	0.48
7:A:609:GOL:H12	9:A:1081:HOH:O	2.12	0.48
1:A:51:TRP:CZ2	1:A:91:CYS:SG	3.07	0.48
1:A:57:LYS:HG3	9:A:775:HOH:O	2.14	0.47
1:A:219[A]:LYS:HA	7:A:604:GOL:C2	2.46	0.46
1:A:194:ALA:HB3	1:A:253[A]:TYR:OH	2.16	0.46
1:A:76:LYS:HD2	9:A:1198:HOH:O	2.15	0.45
1:A:185:GLY:O	1:A:188[B]:SER:OG	2.29	0.45
1:A:508:LYS:HB3	1:A:508:LYS:HE3	1.66	0.43
1:A:249:TYR:CE2	2:C:1:GLC:H4	2.53	0.43
1:A:450[B]:ASP:OD2	1:A:453:GLN:N	2.48	0.43
1:A:248[A]:MET:HE3	9:A:1495:HOH:O	2.20	0.42
1:A:219[B]:LYS:HA	7:A:604:GOL:C2	2.51	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	561/516 (109%)	546 (97%)	14 (2%)	1 (0%)	44 16

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	398	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	487/440 (111%)	479 (98%)	8 (2%)	58 23

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7[A]	MET
1	A	7[B]	MET
1	A	219[A]	LYS
1	A	219[B]	LYS
1	A	331[A]	CYS
1	A	331[B]	CYS
1	A	502	TRP
1	A	504	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GLC	B	1	4	12,12,12	0.53	0	17,17,17	1.07	1 (5%)
4	GLC	B	2[A]	-	11,11,12	0.71	0	15,15,17	2.00	4 (26%)
4	GLC	B	2[B]	-	11,11,12	0.70	0	15,15,17	1.73	3 (20%)
4	GLC	B	3[A]	4	12,12,12	0.62	0	17,17,17	1.97	5 (29%)
4	GLC	B	3[B]	-	12,12,12	1.22	2 (16%)	17,17,17	2.99	7 (41%)
4	GLC	B	4	4	11,11,12	0.64	0	15,15,17	0.81	1 (6%)
2	GLC	C	1	2	12,12,12	0.63	0	17,17,17	1.07	1 (5%)
2	GLC	C	2	2	11,11,12	0.83	1 (9%)	15,15,17	1.01	1 (6%)
2	GLC	D	1	2	12,12,12	0.69	0	17,17,17	1.00	1 (5%)
2	GLC	D	2	2	11,11,12	0.65	0	15,15,17	1.52	3 (20%)
3	BGC	E	1	3	12,12,12	0.60	0	17,17,17	1.16	1 (5%)
3	GLC	E	2	3	11,11,12	0.78	0	15,15,17	1.10	1 (6%)
2	GLC	F	1	2	12,12,12	0.73	0	17,17,17	1.24	2 (11%)
2	GLC	F	2	2	11,11,12	0.64	0	15,15,17	1.35	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '·' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GLC	B	1	4	-	0/2/22/22	0/1/1/1
4	GLC	B	2[A]	-	-	0/2/19/22	0/1/1/1
4	GLC	B	2[B]	-	-	0/2/19/22	0/1/1/1
4	GLC	B	3[A]	4	-	0/2/22/22	0/1/1/1
4	GLC	B	3[B]	-	1/1/5/5	0/2/22/22	0/1/1/1
4	GLC	B	4	4	-	0/2/19/22	0/1/1/1
2	GLC	C	1	2	-	0/2/22/22	0/1/1/1
2	GLC	C	2	2	-	0/2/19/22	0/1/1/1
2	GLC	D	1	2	-	0/2/22/22	0/1/1/1
2	GLC	D	2	2	-	2/2/19/22	0/1/1/1
3	BGC	E	1	3	-	0/2/22/22	0/1/1/1
3	GLC	E	2	3	-	0/2/19/22	0/1/1/1
2	GLC	F	1	2	-	0/2/22/22	0/1/1/1
2	GLC	F	2	2	-	0/2/19/22	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	3[B]	GLC	O5-C1	-2.67	1.36	1.42
2	C	2	GLC	O5-C1	-2.41	1.39	1.43
4	B	3[B]	GLC	O3-C3	2.26	1.48	1.43

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	3[B]	GLC	O1-C1-O5	6.86	130.98	110.38
4	B	3[B]	GLC	O5-C5-C4	-5.47	99.75	109.69
4	B	3[B]	GLC	C4-C3-C2	-5.11	101.91	110.82
4	B	2[A]	GLC	C1-C2-C3	4.12	114.73	109.67
4	B	2[B]	GLC	C1-C2-C3	4.12	114.73	109.67
4	B	2[A]	GLC	O6-C6-C5	3.90	124.67	111.29
4	B	3[A]	GLC	O4-C4-C3	3.61	118.69	110.35
4	B	3[B]	GLC	O1-C1-C2	3.45	118.76	109.03
4	B	3[B]	GLC	C1-O5-C5	3.22	119.74	113.66
4	B	3[B]	GLC	O3-C3-C2	-3.21	102.93	110.35
4	B	3[A]	GLC	O5-C5-C4	-3.18	103.92	109.69
4	B	2[A]	GLC	O3-C3-C2	-3.12	104.02	109.99
4	B	2[B]	GLC	O3-C3-C2	-3.12	104.02	109.99
4	B	3[A]	GLC	C4-C3-C2	3.06	116.17	110.82
4	B	3[A]	GLC	C6-C5-C4	-2.96	106.08	113.00
3	E	1	BGC	O1-C1-O5	-2.94	101.55	110.38
2	C	1	GLC	C4-C3-C2	-2.73	106.05	110.82
2	D	2	GLC	C1-O5-C5	2.69	115.83	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	1	GLC	C1-O5-C5	2.62	118.61	113.66
4	B	2[A]	GLC	O5-C5-C6	-2.49	103.29	107.20
4	B	2[B]	GLC	O5-C5-C6	-2.49	103.29	107.20
2	F	1	GLC	O4-C4-C3	-2.46	104.67	110.35
2	D	2	GLC	C1-C2-C3	-2.45	106.65	109.67
2	C	2	GLC	O5-C5-C6	-2.39	103.45	107.20
2	D	1	GLC	C3-C4-C5	2.35	114.42	110.24
2	F	2	GLC	O3-C3-C2	-2.30	105.59	109.99
4	B	3[B]	GLC	C6-C5-C4	2.18	118.11	113.00
4	B	1	GLC	O3-C3-C4	-2.15	105.38	110.35
3	E	2	GLC	C1-C2-C3	-2.09	107.09	109.67
4	B	3[A]	GLC	O4-C4-C5	-2.09	104.10	109.30
2	D	2	GLC	O5-C5-C4	-2.09	105.75	110.83
4	B	4	GLC	C1-C2-C3	-2.04	107.15	109.67

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	B	3[B]	GLC	C1

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	2	GLC	O5-C5-C6-O6
2	D	2	GLC	C4-C5-C6-O6


There are no ring outliers.


3 monomers are involved in 4 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	3[A]	GLC	3	0
4	B	2[A]	GLC	1	0
2	C	1	GLC	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 1 is monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
7	GOL	A	610	-	5,5,5	4.44	5 (100%)	5,5,5	5.91	4 (80%)
7	GOL	A	604	-	5,5,5	4.31	5 (100%)	5,5,5	3.48	3 (60%)
6	SO4	A	603	-	4,4,4	0.80	0	6,6,6	0.89	0
7	GOL	A	605	-	5,5,5	4.35	5 (100%)	5,5,5	3.71	3 (60%)
7	GOL	A	609	-	5,5,5	4.43	5 (100%)	5,5,5	3.32	3 (60%)
7	GOL	A	607	-	5,5,5	4.47	5 (100%)	5,5,5	5.51	3 (60%)
7	GOL	A	611	-	5,5,5	4.53	5 (100%)	5,5,5	5.92	3 (60%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	GOL	A	606[B]	-	5,5,5	4.50	5 (100%)	5,5,5	4.18	3 (60%)
7	GOL	A	606[A]	-	5,5,5	4.56	5 (100%)	5,5,5	4.49	3 (60%)
8	BGC	A	602	-	12,12,12	0.52	0	17,17,17	1.33	1 (5%)
7	GOL	A	608	-	5,5,5	4.28	5 (100%)	5,5,5	4.13	3 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	A	610	-	-	1/4/4/4	-
7	GOL	A	604	-	-	4/4/4/4	-
7	GOL	A	605	-	-	3/4/4/4	-
7	GOL	A	609	-	-	3/4/4/4	-
7	GOL	A	607	-	-	2/4/4/4	-
7	GOL	A	606[B]	-	-	1/4/4/4	-
7	GOL	A	611	-	-	3/4/4/4	-
7	GOL	A	606[A]	-	-	3/4/4/4	-
8	BGC	A	602	-	-	2/2/22/22	0/1/1/1
7	GOL	A	608	-	-	2/4/4/4	-

All (45) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	607	GOL	C3-C2	-7.32	1.21	1.51
7	A	611	GOL	C3-C2	-7.31	1.21	1.51
7	A	610	GOL	C3-C2	-7.20	1.22	1.51
7	A	606[A]	GOL	C3-C2	-7.17	1.22	1.51
7	A	606[B]	GOL	C3-C2	-6.89	1.23	1.51
7	A	605	GOL	C3-C2	-6.88	1.23	1.51
7	A	608	GOL	C3-C2	-6.78	1.23	1.51
7	A	609	GOL	C3-C2	-6.74	1.24	1.51
7	A	604	GOL	C3-C2	-6.71	1.24	1.51
7	A	606[A]	GOL	O1-C1	4.81	1.62	1.42
7	A	609	GOL	O1-C1	4.74	1.62	1.42
7	A	604	GOL	O1-C1	4.66	1.62	1.42
7	A	611	GOL	O1-C1	4.63	1.62	1.42
7	A	605	GOL	O1-C1	4.61	1.61	1.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	606[B]	GOL	O1-C1	4.58	1.61	1.42
7	A	608	GOL	O1-C1	4.41	1.61	1.42
7	A	607	GOL	O1-C1	4.27	1.60	1.42
7	A	610	GOL	C1-C2	-4.11	1.34	1.51
7	A	606[B]	GOL	O3-C3	4.03	1.59	1.42
7	A	606[A]	GOL	O3-C3	3.87	1.58	1.42
7	A	605	GOL	O3-C3	3.78	1.58	1.42
7	A	610	GOL	O3-C3	3.71	1.58	1.42
7	A	604	GOL	O3-C3	3.71	1.58	1.42
7	A	611	GOL	O3-C3	3.66	1.57	1.42
7	A	609	GOL	O3-C3	3.60	1.57	1.42
7	A	608	GOL	O3-C3	3.51	1.57	1.42
7	A	607	GOL	O3-C3	3.47	1.57	1.42
7	A	610	GOL	O1-C1	3.44	1.56	1.42
7	A	606[B]	GOL	C1-C2	-3.12	1.38	1.51
7	A	609	GOL	C1-C2	-3.03	1.39	1.51
7	A	608	GOL	C1-C2	-2.92	1.39	1.51
7	A	607	GOL	O2-C2	-2.91	1.34	1.43
7	A	609	GOL	O2-C2	-2.85	1.34	1.43
7	A	606[A]	GOL	C1-C2	-2.76	1.40	1.51
7	A	607	GOL	C1-C2	-2.75	1.40	1.51
7	A	604	GOL	O2-C2	-2.72	1.35	1.43
7	A	605	GOL	C1-C2	-2.71	1.40	1.51
7	A	611	GOL	O2-C2	-2.70	1.35	1.43
7	A	611	GOL	C1-C2	-2.68	1.40	1.51
7	A	606[B]	GOL	O2-C2	-2.65	1.35	1.43
7	A	606[A]	GOL	O2-C2	-2.61	1.35	1.43
7	A	608	GOL	O2-C2	-2.30	1.36	1.43
7	A	604	GOL	C1-C2	-2.26	1.42	1.51
7	A	605	GOL	O2-C2	-2.15	1.37	1.43
7	A	610	GOL	O2-C2	-2.06	1.37	1.43

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	611	GOL	O3-C3-C2	10.39	160.02	110.20
7	A	610	GOL	O1-C1-C2	10.34	159.77	110.20
7	A	607	GOL	O3-C3-C2	9.60	156.22	110.20
7	A	606[A]	GOL	O3-C3-C2	7.74	147.33	110.20
7	A	611	GOL	O2-C2-C3	7.54	142.34	109.12
7	A	608	GOL	O3-C3-C2	7.06	144.06	110.20
7	A	606[B]	GOL	O3-C3-C2	6.67	142.17	110.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	607	GOL	O2-C2-C3	6.53	137.89	109.12
7	A	604	GOL	O3-C3-C2	6.38	140.81	110.20
7	A	610	GOL	O3-C3-C2	6.10	139.43	110.20
7	A	605	GOL	O3-C3-C2	5.96	138.76	110.20
7	A	606[A]	GOL	O2-C2-C3	5.78	134.58	109.12
7	A	606[B]	GOL	O2-C2-C3	5.28	132.39	109.12
7	A	605	GOL	O2-C2-C3	5.12	131.66	109.12
7	A	610	GOL	O2-C2-C3	5.04	131.30	109.12
7	A	609	GOL	O2-C2-C3	4.77	130.15	109.12
7	A	609	GOL	O3-C3-C2	4.66	132.55	110.20
7	A	608	GOL	O1-C1-C2	4.26	130.64	110.20
7	A	608	GOL	O2-C2-C3	3.99	126.70	109.12
7	A	607	GOL	O1-C1-C2	3.92	129.01	110.20
7	A	606[B]	GOL	O1-C1-C2	3.70	127.95	110.20
7	A	604	GOL	O2-C2-C3	3.28	123.57	109.12
7	A	611	GOL	O1-C1-C2	3.19	125.49	110.20
7	A	609	GOL	O1-C1-C2	3.01	124.63	110.20
7	A	606[A]	GOL	O1-C1-C2	2.66	122.98	110.20
7	A	605	GOL	O1-C1-C2	2.40	121.69	110.20
7	A	604	GOL	O1-C1-C2	2.39	121.68	110.20
8	A	602	BGC	O5-C5-C6	-2.19	100.99	106.44
7	A	610	GOL	O2-C2-C1	2.02	118.04	109.12

There are no chirality outliers.

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	611	GOL	C1-C2-C3-O3
7	A	604	GOL	O1-C1-C2-C3
7	A	604	GOL	C1-C2-C3-O3
7	A	605	GOL	O1-C1-C2-C3
7	A	606[A]	GOL	O1-C1-C2-C3
7	A	606[A]	GOL	C1-C2-C3-O3
7	A	608	GOL	C1-C2-C3-O3
7	A	609	GOL	O1-C1-C2-C3
7	A	604	GOL	O1-C1-C2-O2
7	A	604	GOL	O2-C2-C3-O3
7	A	611	GOL	O1-C1-C2-C3
7	A	607	GOL	C1-C2-C3-O3
7	A	610	GOL	O1-C1-C2-C3
8	A	602	BGC	O5-C5-C6-O6
7	A	606[B]	GOL	O2-C2-C3-O3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
7	A	608	GOL	O2-C2-C3-O3
7	A	606[A]	GOL	O1-C1-C2-O2
7	A	605	GOL	O1-C1-C2-O2
7	A	605	GOL	O2-C2-C3-O3
7	A	609	GOL	O2-C2-C3-O3
8	A	602	BGC	C4-C5-C6-O6
7	A	609	GOL	O1-C1-C2-O2
7	A	611	GOL	O1-C1-C2-O2
7	A	607	GOL	O1-C1-C2-C3

There are no ring outliers.

7 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	610	GOL	1	0
7	A	604	GOL	2	0
7	A	605	GOL	1	0
7	A	609	GOL	1	0
7	A	606[A]	GOL	1	0
8	A	602	BGC	2	0
7	A	608	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	516/516 (100%)	-0.60	1 (0%) 92 92	7, 17, 30, 55	46 (8%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	4	GLY	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

SUGAR-RSR INFOmissingINFO

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	GOL	A	606[A]	6/6	0.75	0.14	33,39,46,57	6
7	GOL	A	606[B]	6/6	0.75	0.14	38,41,42,42	6
7	GOL	A	609	6/6	0.77	0.14	35,47,61,61	0
7	GOL	A	604	6/6	0.78	0.13	39,82,90,94	0
7	GOL	A	608	6/6	0.81	0.14	49,56,63,74	6
7	GOL	A	605	6/6	0.81	0.15	65,73,78,86	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	GOL	A	611	6/6	0.85	0.16	102,109,112,124	0
7	GOL	A	610	6/6	0.88	0.11	24,33,34,36	6
7	GOL	A	607	6/6	0.89	0.09	47,48,56,63	6
8	BGC	A	602	12/12	0.90	0.11	19,26,38,40	12
6	SO4	A	603	5/5	0.95	0.08	21,23,26,27	5
5	CA	A	601	1/1	1.00	0.03	18,18,18,18	1

6.5 Other polymers [i](#)

There are no such residues in this entry.